arXiv:math/0602574v3 [math.GT] 6 Feb 2007

WARSAW UNIVERSITY
FACULTY OF MATHEMATICS, INFORMATICS AND MECHANICS

Characterization and topological rigidity of Nöbeling manifolds

PHD THESIS

ANDRZEJ NAGÓRKO

Supervisor:
PROF. DR HAB. HENRYK TORUŃCZYK

Warsaw, March 2006

Author's declaration

Aware of legal responsibility I herby declare that I have written this thesis myself and all its contents have been obtained by legal means.

Date

Author's signature

Supervisor's declaration

This thesis is ready to be reviewed.

Date

Supervisor's signature

Thesis supervisor
Henryk Toruńczyk

Author
Andrzej Nagórko

Characterization and topological rigidity of Nöbeling manifolds

Abstract

We develop a theory of Nöbeling manifolds similar to the theory of Hilbert space manifolds. We show that it reflects the theory of Menger manifolds developed by M. Bestvina [8] and is its counterpart in the realm of complete spaces. In particular we prove the Nöbeling manifold characterization conjecture.

We define the n-dimensional universal Nöbeling space ν^n to be the subset of R^{2n+1} consisting of all points with at most n rational coordinates. To enable comparison with the infinite dimensional case we let ν^∞ denote the Hilbert space. We define an n-dimensional Nöbeling manifold to be a Polish space locally homeomorphic to ν^n. The following theorem for $n = \infty$ is the characterization theorem of H. Toruńczyk [36]. We establish it for $n < \infty$, where it was known as the Nöbeling manifold characterization conjecture.

Characterization theorem *An n-dimensional Polish $ANE(n)$-space is a Nöbeling manifold if and only if it is strongly universal in dimension n.*

The following theorem was proved by D. W. Henderson and R. Schori [23] for $n = \infty$. We establish it in the finite dimensional case.

Topological rigidity theorem. *Two n-dimensional Nöbeling manifolds are homeomorphic if and only if they are n-homotopy equivalent.*

We also establish the open embedding theorem, the Z-set unknotting theorem, the local Z-set unknotting theorem and the sum theorem for Nöbeling manifolds.

Key words and phrases.

Nöbeling manifold classification, Nöbeling space characterization, Z-set unknotting theorem, open embedding theorem, carrier, nerve theorem, regular cover, semiregular cover

2000 Mathematics Subject Classification.

Primary 55M10, 54F45; Secondary 54C20.

Promotor
Henryk Toruńczyk

Autor
Andrzej Nagórko

Charakteryzacja i Topologiczna Sztywność Rozmaitości Nöbelinga

Streszczenie

Celem niniejszej rozprawy jest rozwinięcie skończenie wymiarowego odpowiednika klasycznego działu topologii nieskończenie wymiarowej: teorii rozmaitości modelowanych na przestrzeni Hilberta. W szczególności dowodzimy skończenie wymiarowej wersji twierdzenia Toruńczyka o charakteryzacji.

Twierdzenie o charakteryzacji. *Polska n-wymiarowa $ANE(n)$-przestrzeń jest rozmaitością Nöbelinga jedynie jeśli jest mocno uniwersalna w wymiarze n.*

Dowodzimy też skończenie wymiarowej wersji twierdzenia Hendersona klasyfikującego rozmaitości modelowane na przestrzeni Hilberta.

Twierdzenie o sztywności. *Dwie n-wymiarowe rozmaitości Nöbelinga są homeomorficzne jedynie jeśli są n-homotopijnie równoważne.*

Poza tym dowodzimy twierdzenia o jednakowym położeniu Z-zbiorów, jego lokalnej wersji, twierdzenia o otwartym zanurzeniu i twierdzenia o sumie dla rozmaitości Nöbelinga.

Contents

Abstract iii
Streszczenie v
Table of Contents vii
Acknowledgments ix
Citations to Previously Published Work ix
Dedication xi

I Introduction and preliminaries 1

1 Introduction 3

2 Preliminaries 6
2.1 Covers and interior covers 6
2.2 Absolute extensors 8
2.3 Nerves of covers and barycentric stars 10
2.4 Strong universality 11
2.5 n-Homotopy equivalence 15
2.6 Z-sets 20

II Reducing the proof of the main results to the construction of n-regular and n-semiregular $\mathcal{N}_n$-covers 25

3 Approximation within an $\mathcal{N}_n$-cover 27
3.1 $Z(\mathcal{F})$-sets 27
3.2 Approximation within a cover 28
3.3 Z-collections 30

4 Constructing closed $\mathcal{N}_n$-covers 32
4.1 Adjustment of a collection 32
4.2 Limits of sequences of adjustments 33
4.3 Construction of a closed $\mathcal{N}_n$-swelling 34

5 Carrier and nerve theorems **37**
- 5.1 Regular covers . . . 37
- 5.2 Carrier theorem . . . 38
- 5.3 Nerve theorem . . . 39

6 Anticanonical maps and semiregularity **41**
- 6.1 A nerve theorem and the notion of semiregularity . . . 41
- 6.2 A construction of regular covers . . . 44
- 6.3 A construction of semiregular covers . . . 47

7 The homeomorphism extension theorem **49**

8 Proof of the main results **51**

III Constructing n-semiregular and n-regular $\mathcal{N}_n$-covers **57**

9 Basic constructions in $\mathcal{N}_n$-spaces **59**
- 9.1 Adjustment to a Z-collection . . . 59
- 9.2 Fitting closed $\mathcal{N}_n$-neighborhoods . . . 60
- 9.3 Patching of holes . . . 62

10 Core of a cover **64**
- 10.1 The existence of an n-core . . . 65
- 10.2 An n-core of a limit of a sequence of deformations . . . 68
- 10.3 Proof of theorem 10.1 . . . 69
- 10.4 Retraction onto a core and a proof of theorem 6.17 . . . 76

11 Proof of theorem 6.7 **78**
- 11.1 Patching of small holes . . . 78
- 11.2 $\circledcirc$-Contractibility . . . 82
- 11.3 Proof of theorem 6.7 for $k = 0$ . . . 87
- 11.4 $\Cup$-Contractibility . . . 88
- 11.5 Patching of large holes . . . 89
- 11.6 Proof of theorem 6.7 for $k > 0$ . . . 94

Bibliography **98**

Index **101**

Acknowledgments

Completing this thesis was an overwhelming experience and would be impossible withoth the help of others. Foremost I would like to thank my family for support and encouragement during these years.

I am grateful to my thesis supervisor, professor H. Toruńczyk, who helped with his enormous knowledge and was always available to talk and review my writings. I am grateful to reviewers of the first version of this thesis, professors W. Marciszewski and S. Spież, who helped to improve this text a lot.

I would like to thank the organizers of the Topology seminar at the Institute of Mathematics of Polish Academy of Sciences - professors J. Krasinkiewicz and S. Spież - who gave me an opportunity to give a long series of talks on the subject title. I'm grateful to professors T. Dobrowolski and M. Sobolewski who also attended these talks.

I'm grateful to D. Michalik, R. Górak, M. Sawicki and professors E. Pol and A. Chigogidze for helping me out with various questions I have had.

The starting point and the original inspiration for this work was a paper by M. Levin, K. Kawamura and E. D. Tymchatyn, who proved the characterization theorem for one-dimensional Nöbeling spaces [26]. I had the pleasure of meeting professors M. Levin and K. Kawamura at the International Conference and Workshops on Geometric Topology in June 2005, where the results of this thesis were presented.

Citations to Previously Published Work

Contents of chapter 5 were published as a part of

> "Carrier and nerve theorems in the extension theory", A. Nagórko, Proc. Amer. Math. Soc. 135(2):551-558, 2007.

To my father

Part I

Introduction and preliminaries

Chapter 1

Introduction

,,Perhaps this [characterization conjecture for Nöbeling manifolds]
is one of the most exciting problems of general topology".
A. N. Dranishnikov [16]

The aim of this paper is to develop a theory of Nöbeling manifolds similar to the theory of Hilbert space manifolds. In particular, the Nöbeling manifold characterization conjecture is proven.

We define the n-dimensional universal *Nöbeling space* ν^n to be the set of all points of $\mathbb{R}^{2n+1}$ with at most n rational coordinates. To enable comparison with the infinite dimensional case we let ν^∞ denote the Hilbert space $\mathbb{R}^\infty$. We define an *n-dimensional Nöbeling manifold* to be a Polish space locally homeomorphic to ν^n. The following theorem for $n = \infty$ is the characterization theorem of H. Toruńczyk [36]. We establish it for $n < \infty$, where it was known as the Nöbeling manifold characterization conjecture [13, 15, 27, 39].

Characterization theorem. *An n-dimensional Polish space is a Nöbeling manifold if and only if it is an absolute neighborhood extensor in dimension n that is strongly universal in dimension n.*

D. W. Henderson and R. Schori proved [23] the following theorem for $n = \infty$. We give a proof of the finite dimensional case.

Topological rigidity theorem. *Two n-dimensional Nöbeling manifolds are homeomorphic if and only if they are n-homotopy equivalent.*

Topological rigidity and characterization theorems imply a characterization theorem for Nöbeling spaces: every n-dimensional strongly universal Polish absolute extensor in dimension n is homeomorphic to the n-dimensional Nöbeling space. Another consequence of topological rigidity theorem is the open embedding theorem. For $n = \infty$ it was proved by Henderson [22]. We prove it in the finite dimensional case.

Open embedding theorem. *Every n-dimensional Nöbeling manifold is homeomorphic to an open subset of the n-dimensional Nöbeling space.*

The topological rigidity theorem has the following generalization. Again, it was known for $n = \infty$ by results of R. D. Anderson and J. D. McCharen [4] and we prove it in the finite dimensional case.

Z-Set unknotting theorem. *A homeomorphism between Z-sets in n-dimensional Nöbeling manifolds extends to an ambient homeomorphism if and only if it extends to an n-homotopy equivalence.*

We also prove a local version of the Z-set unknotting theorem, for $n < \infty$. It was proved for $n = \infty$ independently by W. Barit [5] and by Cz. Bessaga and A. Pełczyński [6].

Local Z-set unknotting theorem. *For every open cover $\mathcal{U}$ of a Nöbeling manifold there exists an open cover $\mathcal{V}$ such that every homeomorphism between Z-subsets of the manifold that is $\mathcal{V}$-close to the inclusion extends to a homeomorphism of the entire manifold that is $\mathcal{U}$-close to the identity.*

By an example of R. B. Sher a space that is an union of two Hilbert cubes that meet in a Hilbert cube needs not to be a Hilbert cube [33]. We show that no such example exists in the theory of Nöbeling spaces.

Sum theorem. *If a space X is an union of two closed n-dimensional Nöbeling manifolds whose intersection is also an n-dimensional Nöbeling manifold, then X is an n-dimensional Nöbeling manifold.*

As we shall see in chapter 8 the proof of the above mentioned theorems can rather easily be reduced to a proof of the main theorem, stated below. We say that a space is an *abstract Nöbeling manifold* if it is an n-dimensional Polish absolute neighborhood extensor in dimension n that is strongly universal in dimension n, i.e. if it satisfies the conditions of the characterization theorem.

Main theorem. *A homeomorphism between Z-sets in abstract Nöbeling manifolds extends to an ambient homeomorphism if and only if it extends to an n-homotopy equivalence.*

The theory developed in this paper reflects the theory of Menger manifolds developed by M. Bestvina [8]. He proved finite dimensional analogues of the characterization of the Hilbert cube given by Toruńczyk [35] and of the topological rigidity of Hilbert cube manifolds given by J. West [38] and T. A. Chapman [12]. Proofs given in the present paper does not admit direct generalizations neither to the compact case nor to the complete, infinite dimensional case.

Special cases of the characterization and the Z-set unknotting theorems, for the case of Nöbeling spaces rather than that of Nöbeling manifolds, were established by several authors. The one-dimensional cases were proved by K. Kawamura, M. Levin and E. D. Tymchatyn [26]. S. Ageev announced a proof of the characterization theorem and of a special case of the Z-set unknotting theorem [1] but recently he declared it contained a gap [2]. The Z-set unknotting theorem for a certain model of a Nöbeling space was proved by Levin [28]. Levin's result and those of this thesis were presented at the International Conference and Workshops on Geometric Topology, 3-10 July 2005. Very recently, Levin posted a preprint [29] containing a proof of the characterization theorem for Nöbeling spaces of all dimensions.

How the paper is organized

The text is divided into three parts. In the first part we recall the general results used. In the second part we reduce the proof of the theorems menitoned above to the proofs of two theorems (6.7 and 6.17) that give sufficient conditions for the existence of covers of abstract Nöbeling manifolds that resemble triangulations of simplicial complexes. In the third part these two theorems are proved.

Chapter 2

Preliminaries

To facilitate reading of this paper we recall the general results and notions used throughout it. It makes the paper self-contained in a sense that, with few exceptions, in the next chapters there are no references to external sources.

The image and the domain of a function f are denoted by $\operatorname{im} f$ and by $\operatorname{dom} f$ respectively. We write $f(A)$ for $f(A \cap \operatorname{dom} f)$ for each map f and each set A, i.e. we do not require A to be a subset of the domain of f. The unit interval is denoted by $[0,1]$ and the symbol I is used exclusively to denote a set of indices of a collection of sets. If $\mathcal{F} = \{F_i\}_{i \in I}$ is an indexed collection of sets and J is a non-empty subset of the indexing set I, then we let $F_J = \bigcap_{j \in J} F_j$. We use the following symbols to denote standard topological operations: $\operatorname{Cl}_X A = \operatorname{Cl} A$ denotes the *closure* of a subset A of a space X, $\operatorname{Int}_X A = \operatorname{Int} A$ denotes the *interior* of a subset A of a space X and ∂X denotes the *geometrical boundary* of an euclidean manifold X.

We say that a space is *Polish* is it is separable and completely metrizable. A *map* is a continuous function between two topological spaces. We say that sets *intersect* if their intersection is non-empty.

2.1 Covers and interior covers

We say that a collection $\mathcal{F}$ of sets is *indexed*, whenever there is a chosen *indexing set* I and an *indexing function* that assigns to each element i of I an element F_i of $\mathcal{F}$. We shall write $\mathcal{F} = \{F_i\}_{i \in I}$ for short. Note that the indexing function is onto, but needs not to be one-to-one. Nonetheless, we will distinguish between elements of $\mathcal{F}$ that are equal, but have different indices. Unless otherwise stated, we shall assume that every collection of sets is indexed.

Definition. A *cover* of a topological space X is a collection of subsets of X whose union equals to X. An *interior cover* of X is a collection of subsets of X whose interiors cover X.

Symbols $\mathcal{F}, \mathcal{G}$ will usually denote closed covers and symbols $\mathcal{U}, \mathcal{V}$ will usually denote open covers.

Definition. Elements of two indexed collections of sets are *corresponding* if they have equal indices. Covers with equal indexing sets are *isomorphic* if for each set of intersecting elements in one cover the corresponding elements in the other cover intersect.

In more formal terms, a cover $\mathcal{F} = \{F_i\}_{i \in I}$ is isomorphic to a cover $\mathcal{G} = \{G_i\}_{i \in I}$ if for each non-empty $J \subset I$ the intersection $F_J = \bigcap_{j \in J} F_j$ is non-empty if and only if the intersection $G_J = \bigcap_{j \in J} G_j$ is non-empty. For every $i \in I$ elements F_i of $\mathcal{F}$ and G_i of $\mathcal{G}$ are corresponding.

Definition 2.1. Let $\mathcal{F} = \{F_i\}_{i \in I}$ and $\mathcal{G} = \{G_i\}_{i \in I}$ be collections of subsets of a space X, indexed by the same indexing set I.

(a) We say that $\mathcal{G}$ *is a swelling of* $\mathcal{F}$, if it is isomorphic to $\mathcal{F}$ and if every element of $\mathcal{F}$ is a subset of the corresponding element of $\mathcal{G}$.

(b) We say that $\mathcal{F}$ *is a shrinking of* $\mathcal{G}$, if $\mathcal{G}$ is a swelling of $\mathcal{F}$.

(c) For each subset A of X, we let $\{F_i \cap A\}_{i \in I}$ be *the restriction of* $\mathcal{F}$ *to* A and denote it by $\mathcal{F}/A$.

(d) For each subset A of X, we say that $\mathcal{F}$ *is equal to* $\mathcal{G}$ *on* A, if $F_i \cap A = G_i \cap A$ for each i in I, i.e. if the restrictions of $\mathcal{F}$ and $\mathcal{G}$ to A are equal.

(e) We say that $\mathcal{F}$ *refines a collection* $\mathcal{H}$ of subsets of X, if every element of $\mathcal{F}$ is a subset of an element of $\mathcal{H}$.

(f) For each point x in X, we let $\operatorname{st}_{\mathcal{F}} x$ be the union of all elements of $\mathcal{F}$ that contain x. We say that $\operatorname{st}_{\mathcal{F}} x$ is a *star of* x *in* $\mathcal{F}$.

(g) For each subset A of X, we let $\operatorname{st}_{\mathcal{F}} A$ be the union of A with all elements of $\mathcal{F}$ that intersect A. We say that $\operatorname{st}_{\mathcal{F}} A$ is a *star of* A *in* $\mathcal{F}$.

(h) For each collection $\mathcal{H}$ of subsets of X, we let $\operatorname{st}_{\mathcal{H}} \mathcal{F} = \{\operatorname{st}_{\mathcal{H}} F_i\}_{i \in I}$. We say that $\operatorname{st}_{\mathcal{H}} \mathcal{F}$ is a *star of* $\mathcal{F}$ *in* $\mathcal{H}$.

(i) For each collection $\mathcal{H}$ of subsets of X and each $m > 0$, we let $\operatorname{st}^m_{\mathcal{H}} \mathcal{F} = \operatorname{st}_{\mathcal{H}} \operatorname{st}^{m-1}_{\mathcal{H}} \mathcal{F}$ and $\operatorname{st}^0_{\mathcal{H}} \mathcal{F} = \mathcal{F}$. We say that $\operatorname{st}^m \mathcal{F}$ is *the mth star of* $\mathcal{F}$ *in* $\mathcal{H}$.

(j) For each $m > 0$, we let $\operatorname{st}^m \mathcal{F} = \operatorname{st}_{\mathcal{F}} \operatorname{st}^{m-1} \mathcal{F}$ and $\operatorname{st}^0 \mathcal{F} = \mathcal{F}$. We say that $\operatorname{st}^m \mathcal{F}$ is *the mth star of* $\mathcal{F}$.

Note that $\mathcal{F}$ and $\operatorname{st}_{\mathcal{H}} \mathcal{F}$ have the same indexing set, hence it makes sense to talk about corresponding elements of $\mathcal{F}$ and $\operatorname{st}_{\mathcal{H}} \mathcal{F}$.

Lemma 2.2. *If k and m are nonnegative integers and $\mathcal{F}$ is a collection of sets, then*

$$\operatorname{st}^k_{\mathcal{F}} \operatorname{st}^m \mathcal{F} = \operatorname{st}^{k+m} \mathcal{F} \text{ and } \operatorname{st}^k \operatorname{st}^m \mathcal{F} = \operatorname{st}^{2km+m+k} \mathcal{F}.$$

2.2 Absolute extensors

Many characterizations of absolute extensors for various classes of topological spaces made it into quite a mess in notation. We are going to settle for the following definitions.

Definition. We say that a space X is an *absolute extensor in dimension* n if it is a metric space and if every map into X from a closed subset of an n-dimensional metric space extends over the entire space. The class of absolute extensors in dimension n is denoted by $AE(n)$ and its elements are called $AE(n)$*-spaces.*

Definition. We say that a metric space X is an *absolute neighborhood extensor in dimension* n if it is a metric space and if every map into X from a closed subset A of an n-dimensional metric space extends over an open neighborhood of A. The class of absolute neighborhood extensors in dimension n is denoted by $ANE(n)$ and its elements are called $ANE(n)$*-spaces.*

The following characterization of the class of absolute neighborhood extensors in dimension n is proven in [25], with references to original papers by K. Borsuk and J. Dugundji.

Definition 2.3. We say that a map g is $\mathcal{U}$*-close* to a map f, or that g is a $\mathcal{U}$*-approximation* of f, if $\mathcal{U}$ is an open cover of the common codomain of f and g and the collection $\{f^{-1}(U) \cap g^{-1}(U)\}_{U \in \mathcal{U}}$ covers the common domain of f and g.

Theorem 2.4. *The following conditions are equivalent for every metric space X and every integer $n < \infty$:*

(1) X is an $ANE(n)$-space.

(2) X is locally k-connected for every $k < n$. In other words, for each point x in X and every neighborhood U of x there exists a smaller neighborhood V of x such that every map from the boundary of a $(k+1)$-dimensional ball into V extends to a map of the entire ball into U.

*(3) for each open cover $\mathcal{U}$ of X there is an open cover $\mathcal{V}$ of X such that any two $\mathcal{V}$-close maps into X, defined on an at most $(n-1)$-dimensional Polish space, are $\mathcal{U}$-homotopic. Recall that two maps are $\mathcal{U}$-*homotopic *if they are homotopic by a homotopy whose paths refine $\mathcal{U}$.*

The third condition of theorem 2.4 implies the following proposition, as the space of maps (taken with the sup topology) from a k-dimensional sphere into a separable metric space is separable.

Proposition 2.5. *If a space is a separable absolute neighborhood extensor in dimension n, then its homotopy groups of dimensions less than n are countable.*

A characterization of the class of absolute extensors in dimension n was given by J. Dugundji [17] in a form of the following theorem.

Theorem 2.6. *A space is an absolute extensor in dimension* n *if and only if it is an absolute neighborhood extensor in dimension* n *and its homotopy groups of dimensions less than* n *vanish.*

Theorem 2.7 is a closed sum theorem for absolute neighborhood extensors in dimension n.

Theorem 2.7 ([25, p. 49])**.** *A space is an absolute neighborhood extensor in dimension* n *whenever it can be represented as an union of two closed* $ANE(n)$*-subspaces that meet in an* $ANE(n)$*-subspace.*

We cite the following theorem after [13, Proposition 4.1.7, p. 131], with $(*)_{n-1}$ renamed to $(*)_n$.

Theorem 2.8. *Every open cover* $\mathcal{U}$ *of an* $ANE(n)$*-space* Y *has an open refinement* $\mathcal{V}$ *such that the following condition holds:*

$(*)_n$ *For any at most* n*-dimensional metric space* B*, any closed subspace* A *of* B*, and any two* $\mathcal{V}$*-close maps* $f, g\colon A \to Y$*, if* f *has an extension* $\tilde{f}\colon B \to Y$*, then* g *has an extension* $\tilde{g}\colon B \to Y$ *that is* $\mathcal{U}$*-close to* $\tilde{f}$*.*

Lemma 2.9. *For every open neighborhood* U *of a closed* $AE(n)$*-subset of an at most* n*-dimensional* $ANE(n)$*-space there exists an open neighborhood* V *such that every map from a closed subset of an at most* n*-dimensional metric space into* V *extends over the entire space to a map into* U*.*

Proof. By theorem 2.4, U is an absolute neighborhood extensor in dimension n. By theorem 2.8, there exists an open cover $\mathcal{V}$ of U, for which $(*)_n$ is satisfied with $\mathcal{U} = \{U\}$. Let A be a closed subset of U that is an $AE(n)$-space. Since U is at most n-dimensional, there exists a retraction $r\colon U \to A$. Let

$$V = \bigcup_{W \in \mathcal{V}} r^{-1}(W) \cap W.$$

It is an open neighborhood of A. We shall prove that it satisfies the assertion.

Let f be a map into V from a closed subset B of an at most n-dimensional metric space X. By the definition of V, maps $r \circ f$ and f are $\mathcal{V}$-close. Since A is an absolute extensor in dimension n, the map $r \circ f$ extends over X to a map into A. Hence, by $(*)_n$, f extends over X to a map into U. We are done. $\square$

2.3 Nerves of covers and barycentric stars

We define simplicial complexes in the usual way [25, p. 99]. We do not allow infinite-dimensional simplices. Unless otherwise stated, simplicial complexes are endowed with the metric topology [25, p. 100]. We do not assume that they are countable or locally finite.

Theorem 2.10 ([25, p. 107]). *Every simplicial complex is topologically complete. Every countable simplicial complex is separable.*

Definition. Let K be a simplicial complex. A *barycentric star* $\operatorname{bst} v$ of a vertex v of K is the union of all simplices of the first barycentric subdivision of K that contain v. We let $\operatorname{bst} L$ be the union of barycentric stars of vertices of a subcomplex L of K. We treat it as a subcomplex of the barycentric subdivision of K and define *the mth barycentric star* $\operatorname{bst}^m L$ *of* L recursively by the formula $\operatorname{bst}^m L = \operatorname{bst}(\operatorname{bst}^{m-1} L)$.



Figure 2.1: The first and the second barycentric star of a vertex v of a triangle.

Definition. The nerve of an point-finite indexed collection $\mathcal{F}$ of sets is a simplicial complex whose set of vertices is in one-to-one correspondence with non-empty elements of $\mathcal{F}$ and whose vertices span a simplex if and only if the corresponding elements in $\mathcal{F}$ intersect.

Lemma 2.11. *A collection of barycentric stars of vertices of a simplicial complex is its closed locally finite cover. The nerve of a cover of a simplicial complex K by barycentric stars of its vertices is isomorphic to K. The cover of the nerve of a cover $\mathcal{F}$ by barycentric stars of its vertices is isomorphic to $\mathcal{F}$.*

Proof. Let K be a simplicial complex. Observe that

(1) *a barycentric star of a vertex of K intersects a simplex in K if and only if the vertex lies in the simplex,*

(2) vertices of K span a simplex if and only if their barycentric stars intersect.

Choose a point in K and let δ be a simplex that contains it. By (1), the union of barycentric stars of vertices contained in $K \setminus \delta$ is disjoint from δ. By [25, p. 99], every subcomplex of a simplicial complex is closed, hence the complement of this union is an open neighborhood of δ (and of the chosen point). By the definition, it meets only barycentric stars of vertices contained in δ, hence the cover of K by barycentric stars of its vertices is locally finite.

Let $\mathcal{B}$ be a cover of K by barycentric stars of its vertices. By (2), a set of vertices of K spans a simplex if and only if their barycentric stars intersect. By the definition, the elements of $\mathcal{B}$ intersect if and only if the corresponding vertices of the nerve of $\mathcal{B}$ span a simplex. Hence the second assertion of the lemma.

By the definition, a set of elements of a collection $\mathcal{F}$ of sets has non-empty intersection if and only if the corresponding vertices in the nerve of $\mathcal{F}$ span a simplex. By (2), these vertices span a simplex if and only if their barycentric stars intersect. Hence the third assertion of the lemma. ☐

Lemma 2.12. *A cover of a simplicial complex by barycentric stars of its vertices has the property that every intersection of its elements is either empty or contractible.*

Proof. Let K be a simplicial complex and let $\mathcal{B}$ be a cover of K by barycentric stars of its vertices. By lemma 2.11, the nerve of $\mathcal{B}$ is isomorphic to K. Hence if elements of $\mathcal{B}$ intersect, then the collection $\{v_k\}$ of corresponding vertices of K span a simplex δ in K. By the definition, a barycentric star $\operatorname{bst} v_k$ is a full subcomplex of the first barycentric subdivision of K that contains all barycenters of faces that contain v_k (with a zero dimensional face $\{v_k\}$ included). Hence the intersection $\bigcap \operatorname{bst} v_k$ is a full subcomplex of the first barycentric subdivision of K that contains all barycenters of faces that contain δ. Observe that this subcomplex is a cone with apex equal to the the barycenter of δ. By [34, p. 116], it is contractible. We are done. ☐

2.4 Strong universality

In the present paper, we shall endow all function spaces with the limitation topology, which embraces a concept of closeness introduced in definition 2.3.

Definition. We let $C(X,Y)$ denote the set of all maps from a space X into a space Y. For each map $f\colon X \to Y$ and for each open cover $\mathcal{U}$ of Y, we let $B(f,\mathcal{U})$ denote the set of all maps that are $\mathcal{U}$-close to f. The *limitation topology* on $C(X,Y)$ is defined by a basis of neighborhoods $\{B(f,\mathcal{U})\}$, where f runs over all elements of $C(X,Y)$ and $\mathcal{U}$ runs over all open covers of Y.

It should be emphasized that this topology needs not to be metrizable and the neighborhoods $B(f,\mathcal{U})$ defined above need not to be open in it [11].

Lemma 2.13 ([36]). *If X and Y are completely metrizable, then $C(X,Y)$ endowed with the limitation topology has the Baire property. Moreover, every closed subset of $C(X,Y)$ has the Baire property.*

See [11, p. 429] for a detailed discussion.

Definition. We say that *a map f is approximable by embeddings* to express that "for every open cover $\mathcal{U}$ of the codomain of f there exists an embedding, which is $\mathcal{U}$-close to f". Likewise for closed embeddings, Z-embeddings, etc.

In other words, a map f from X into Y is approximable by elements of a subset of $C(X,Y)$ if it lies in its closure.

Definition. We say that a Polish space X is *strongly universal in dimension n* if every map from an at most n-dimensional Polish space into X is approximable by closed embeddings.

In other words, a Polish space Y is strongly universal in dimension n, if for each at most n-dimensional Polish space X, the set of closed embeddings is a dense subset of $C(X,Y)$.

Proposition 2.14 ([13, p. 34]). *If X is a Polish space, then the set of closed embeddings of X into a space Y is a G_δ-set in $C(X,Y)$ endowed with the limitation topology.*

The following two theorems imply that every n-dimensional Nöbeling manifold is strongly universal in dimension n, which is the "easy" part of the characterization theorem.

Theorem 2.15. *The n-dimensional Nöbeling space is strongly universal in dimension n.*

Following the spirit of the literature, we omit the proof. Hint: by lemma 1.4(b) from [36], it suffices to show that every map from a countable simplicial complex into ν^n is approximable by closed embeddings.

Theorem 2.16. *If a Polish $ANE(n)$-space X is strongly universal in dimension n, then every open subset of X is strongly universal in dimension n. If every point of a Polish $ANE(n)$-space X has a neighborhood that is strongly universal in dimension n, then X is strongly universal in dimension n.*

Theorem 2.16 appears to be a folk theorem, well known to the experts, but unpublished.

Proof. We base the proof on the observation that a map g is a closed embedding whenever there is a closed interior cover $\mathcal{G}$ of the codomain such that for every $G \in \mathcal{G}$ the restriction $g_{|g^{-1}(G)}$ is a closed embedding.

Consider the following condition for each collection $\mathcal{F}$ of subsets of a space Y.

$(*_{\mathcal{F},Y})$ *for each $F \in \mathcal{F}$, every map from at most n-dimensional Polish space into F is approximable by closed embeddings into Y (with a control by covers of Y).*

We will show that if $(*_{\mathcal{F},Y})$ is satisfied for some interior cover $\mathcal{F}$ of an $ANE(n)$-space Y, then Y is strongly universal in dimension n.

Let $\mathcal{F}$ be an interior cover of an $ANE(n)$-space Y. Let $\mathcal{G}$ denote a closed interior cover of Y whose second star refines $\mathcal{F}$. Let X denote an at most n-dimensional Polish space and let f be a map from X into Y. Let B denote the interior, with respect to the limitation topology, of the set of maps from X into Y that are $\mathcal{G}$-close to f. It contains f (see [11]). It has the Baire property by lemma 2.13. Fix $G \in \mathcal{G}$ and let B_G denote a subset of B that contains maps whose restrictions to $f^{-1}(\mathrm{st}_{\mathcal{G}}\, G)$ are closed embeddings. By proposition 2.14, it is a G_δ-set in B. We'll prove that it is dense in B. Let g denote an element of B and let $\mathcal{U}$ be an open cover of Y. We will show that g has a $\mathcal{U}$-approximation by a map that belongs to B_G. Since B is open, we may assume that every map $\mathcal{U}$-close to g belongs to B. Let $\mathcal{V}$ denote a cover that satisfies condition $(*)_n$ of theorem 2.8 for $\mathcal{U}$. Since g is $\mathcal{G}$-close to f and the second star of $\mathcal{G}$ refines $\mathcal{F}$, there is $F \in \mathcal{F}$ such that $g(f^{-1}(\mathrm{st}_{\mathcal{G}}\, G)) \subset F$. By $(*_{\mathcal{F},Y})$ there is a $\mathcal{V}$-approximation of $g_{|f^{-1}(\mathrm{st}_{\mathcal{G}}\, G)}$ by a closed embedding into Y and by the definition of $\mathcal{V}$ it extends over X to a map that is $\mathcal{U}$-close to g. Therefore B_G is dense in B.

By the Baire property of B the intersection $\bigcap_{G\in\mathcal{G}} B_G$ is dense in B. Therefore f is approximable by maps that are $\mathcal{G}$-close to f and whose restrictions to $f^{-1}(\mathrm{st}_{\mathcal{G}}\, G)$ are closed embeddings for every $G \in \mathcal{G}$. Let g denote such a map. Observe that $g^{-1}(G) \subset f^{-1}(\mathrm{st}_{\mathcal{G}}\, G)$, since g is $\mathcal{G}$-close to f. Therefore g is a closed embedding on $g^{-1}(G)$ for every $G \in \mathcal{G}$. Hence it is a closed embedding of X into Y. Therefore we proved that if $(*_{\mathcal{F},Y})$ is satisfied for an interior cover $\mathcal{F}$ of an $ANE(n)$-space Y, then Y is strongly universal in dimension n.

To finish the proof it suffices to show that if we assume hypthesis of either of the implications of the theorem, then there exists an interior cover $\mathcal{F}$ of X that satisfies condition $(*_{\mathcal{F},X})$.

Let X be a Polish $ANE(n)$-space that is strongly universal in dimension n. Let U be an open subset of X. For each $x \in U$ let F_x be a closed (in X) neighborhood of x such that $F_x \subset U$. We will show that the interior cover $\mathcal{F} = \{F_x\}_{x\in U}$ of U satisfies $(*_{\mathcal{F},U})$. Pick $F_x \in \mathcal{F}$ and let $\mathcal{U}$ be an open cover of U. Let $\mathcal{V} = \{X \setminus F_x\} \cup \mathcal{U}$. It is a cover of X. By strong universality of X, every map from at most n-dimensional Polish space into F_x is $\mathcal{V}$-approximable by closed embeddings into X. But every $\mathcal{V}$-approximation of a map into F_x must have image in U. Hence $\mathcal{F}$ satisfies condition $(*_{\mathcal{F},U})$. By theorem 2.4, U is an $ANE(n)$-space. Hence U is stringly universal in dimension n. This finishes the proof of the first implication.

Let X be a Polish $ANE(n)$-space such that each point x in X has a neighborhood N_x that is strongly universal in dimension n. This neighborhood may be neither open nor closed. Let F_x be a closed neighborhood of x and let U_x be an open neighborhood of x such that $F_x \subset U_x \subset \mathrm{Cl}_X U_x \subset N_x$. We will show that the interior cover $\mathcal{F} = \{F_x\}_{x \in X}$ of X satisfies $(*_{\mathcal{F},X})$. Pick $F_x \in \mathcal{F}$ and let $\mathcal{U}$ be an open cover of X. Let f be a map from at most n-dimensional Polish space into F_x. We have to show that f has a $\mathcal{U}$-approximation by a closed embedding into X. Without a loss of generality we may assume that $\mathcal{U}$ refines $\{U_x, X \setminus F_x\}$. Let $\mathcal{V} = \{U \cap N_x\}_{U \in \mathcal{U}}$. Since N_x is strongly universal in dimension n, f has a $\mathcal{V}$-approximation by a closed embedding into N_x. Since $\mathcal{V}$ refines $\{U_x, X \setminus F_x\}$ and the image of f is contained in F_x, the image of this approximation is contained in U_x. This image is closed in $\mathrm{Cl}_X U_x$, as $\mathrm{Cl}_X U_x$ is a subset of N_x, so it must be closed in X. Hence $(*_{\mathcal{F},X})$ is satisfied and X is strongly universal in dimension n. This finishes the proof of the second implication. ☐

The characterization theorem states that every n-dimensional Polish strongly universal $ANE(n)$-space is a Nöbeling manifold. It is convenient to have a shorthand for the class of spaces that possess these properties. To this end we adapt the notion of an $\mathcal{N}_n$-space from [26], with the exception that we do not require that an $\mathcal{N}_n$-space has vanishing homotopy groups of dimensions less than n.

Definition 2.17. We say that a space is an *$\mathcal{N}_n$-space* if it is an n-dimensional Polish $ANE(n)$-space that is strongly universal in dimension n. By $\mathcal{N}_n$ we denote the class of $\mathcal{N}_n$-spaces.

In the language used in the introduction, the class of $\mathcal{N}_n$-spaces is the class of abstract n-dimensional Nöbeling manifolds.

Theorem 2.18. *Every non-empty open subset of an $\mathcal{N}_n$-space is an $\mathcal{N}_n$-space. If every point of a separable space X has a neighborhood that is an $\mathcal{N}_n$-space, then X is an $\mathcal{N}_n$-space.*

Proof. Let U be a non-empty open subset of an $\mathcal{N}_n$-space X. By strong universality of X, one can embed an n-dimensional space into U, hence U itself is n-dimensional. By [18, theorem 4.3.23], every open subset of a Polish space is Polish, hence U is Polish. By theorem 2.4, U is an absolute extensor in dimension n. By theorem 2.16, U is strongly universal in dimension n. Hence U is an $\mathcal{N}_n$-space.

If every point in X has an $\mathcal{N}_n$-neighborhood, then X is clearly n-dimensional. Complete metrizability is a local property, hence X is Polish. By theorem 2.4, X is an $ANE(n)$-space. By theorem 2.16, X is strongly universal in dimension n. Hence X is an $\mathcal{N}_n$-space. ☐

We finish this section with a technical lemma that will be used in chapter 4.

Lemma 2.19. *For each open cover $\mathcal{U}$ of a Polish space X there exists a sequence $\mathcal{U}_1, \mathcal{U}_2, \ldots$ of open covers of X such that for each sequence $f_1, f_2, \ldots$ of maps from X into X such that $\operatorname{im} f_{k-1} \subset \operatorname{dom} f_k$ and such that f_k is $\mathcal{U}_k$-close to the inclusion $\operatorname{dom} f_k \subset X$, the pointwise limit $\lim_{k\to\infty} f_k$ exists, is continuous and is $\mathcal{U}$-close to the identity.*

Proof. Let α be a map from X into $(0, \infty)$. We say that a map $g\colon X \to X$ is α-close to a map $f\colon X \to X$, if $d(f(x), g(x)) \leq \alpha(f(x))$ for each $x \in X$. By lemma 2.1.6 of [13] there is an open cover $\mathcal{V}_\alpha$ of X such that if a map g is $\mathcal{V}_\alpha$-close to a map f, then it is α-close to it. We define a map $\alpha_{\mathcal{U}}\colon X \to (0, \infty)$ by the formula $\alpha_{\mathcal{U}}(x) = (1/2) \sup\{d(x, X \setminus U)\colon U \in \mathcal{U}\}$ (the formula is taken from page 35 of [13]). Again by lemma 2.1.6 of [13] if g is $\alpha_{\mathcal{U}}$-close to g, then it is $\mathcal{U}$-close to it. We let

$$\mathcal{U}_k = \mathcal{V}_{(1/2^{k+1})\alpha_{\mathcal{U}}}.$$

By the triangle inequality for the Hausdorff metric for each U the map $x \mapsto d(x, X \setminus U)$ is 1-Lipschitz. Hence $\alpha_{\mathcal{U}}$ is 1-Lipschitz. Let $f_k\colon X \to X$, $k \geq 0$, be a sequence of maps such that f_k is $\mathcal{U}_k$-close to f_{k-1}. A direct computation shows that the distance between $f_k(x)$ and $f_0(x)$ is at most $(1 - 1/2^k)\alpha_{\mathcal{U}}(f_0(x))$. Therefore by completeness of X the limit $f_\infty = \lim_{k\to\infty} f_k$ exists and the map f_∞ is $\alpha_{\mathcal{U}}$-close, hence $\mathcal{U}$-close, to f_0.

□

2.5 n-Homotopy equivalence

First we introduce a notion of a weak n-homotopy equivalence. Note that by a characterization theorem of J. H. C. Whitehead (to be given on page 17), the phrase "n-homotopy equivalences" in theorems stated in the introduction can be replaced by "weak n-homotopy equivalences".

Definition. We say that a map is a *weak n-homotopy equivalence* if it induces isomorphisms on homotopy groups of dimensions less than n, regardless of the choice of the base point.

Definition. We say that maps $f, g\colon X \to Y$ are *n-homotopic* if for every map Φ from a complex of dimension less than n into X, the compositions $f \circ \Phi$ and $g \circ \Phi$ are homotopic in the usual sense.

Lemma 2.20. *If a map is n-homotopic to a weak n-homotopy equivalence, then it is a weak n-homotopy equivalence.*

Proof. Let $\varphi\colon X \to Y$ ba a weak n-homotopy equivalence and let ψ be a map that is n-homotopic to φ. Fix a base point in X and let $0 \leq k < n$. We have to show that ψ

induces an isomorphism on kth homotopy groups of X and Y. Let K be a wedge of k-dimensional spheres and let $f\colon K \to Y$ be a map that induces an epimorphism on kth homotopy groups of K and Y. By the definition of n-homotopy, $\varphi \circ f$ and $\psi \circ f$ are homotopic. Let γ be a patch of the base-point under this homotopy. Let φ_* be an isomorphism of $\pi_k(X)$ and $\pi_k(Y)$ induced by φ. The homomorphism ψ_* induced by ψ on $\pi_k(X)$ and $\pi_k(Y)$ is equal to an isomorphism φ_* conjugated with γ (see [21, p. 341] for precise definitions). Hence ψ_* is an isomorphism as well. $\square$

Lemma 2.21. *Every $ANE(n)$-space Y admits an open cover $\mathcal{U}$ such that every map that is $\mathcal{U}$-close to a weak n-homotopy equivalence of a metric space X and Y is also a weak n-homotopy equivalence.*

In particular, the set of weak n-homotopy equivalences of X and Y is an open subset of $C(X, Y)$.

Proof. By theorem 2.4, there exists an open cover $\mathcal{U}$ of Y such that every two $\mathcal{U}$-close maps defined on a metric space of dimension less than n are homotopic. In particular, every two $\mathcal{U}$-close maps into Y are n-homotopic. By lemma 2.20, the cover $\mathcal{U}$ satisfies our claim. $\square$

Lemma 2.22. *If $\varphi\colon X \to Y$ is a weak n-homotopy equivalence, K is a simplicial complex whose dimension is less than n and f is a map from K into Y, then there exists a map $f'\colon K \to X$ whose composition with φ is homotopic to f. If L is a subcomplex of K and if we are given a map from L into X whose composition with φ is equal to $f_{|L}$, then we may require that f' extends this map and that the homotopy of f and $\varphi \circ f'$ is $\operatorname{rel} L$. If K is a subcomplex of an at most n-dimensional simplicial complex M and f extends to a map g from M into Y, then f' extends to a map from M into X. In particular, there exists a map from M into X whose composition with φ is n-homotopic to g.*

Proof. The proof of the first two implications follows exactly the proof of proposition 6.4 from page 77 of [7]. To prove the third implication, consider an extension $g\colon M \to X$ of f. Without a loss of generality we may assume that there are no simplices of dimension less than n in $M \setminus K$ (by enlarging K if needed) and that M is endowed with a weak topology (by theorem 6.4 on page 81 of [7], M endowed with a weak topology is homotopy equivalent to M endowed with a metric topology). Hence all we have to do is to show how to extend f' over a single n-dimensional simplex δ of M. The map f' is defined on $\partial\delta$. By the assumptions, $\varphi \circ f'_{|\partial\delta}$ is homotopic to $f_{|\partial\delta}$ and $f_{|\partial\delta}$ extends over δ to a map into X. Hence $\varphi \circ f'_{|\partial\delta}$ represents a trivial element of $\pi_{n-1}(Y)$. Since φ is a weak n-homotopy equivalence, $f'_{|\partial\delta}$ must represent a trivial element of $\pi_{n-1}(X)$. Hence $f'_{|\partial\delta}$ extends over δ to a map into X. Hence f' admits an extension to a map g' from M into X. It follows from the cellular approximation theorem [21, Corollary 4.12], that $\varphi \circ g'$ is n-homotopic to g. $\square$

Definition. We say that a map $\varphi\colon X \to Y$ is an *n-homotopy equivalence* if there exists a map $\psi\colon Y \to X$ such that $\psi \circ \varphi$ is n-homotopic to id_X and $\varphi \circ \psi$ is n-homotopic to id_Y. We say that X is *n-homotopy equivalent* to Y if there exists an n-homotopy equivalence of X and Y.

Whitehead's characterization. *A map of two n-dimensional $ANE(n)$-spaces is a weak n-homotopy equivalence if and only if it is an n-homotopy equivalence.*

The theorem was stated by Whitehead [40] with slightly stronger assumptions, although his proof works in this case as well. For completeness, we give a proof based on the argument from [7]. A proof of an even more general case appeared in [14].

Proof. The proof is a modification of the proof of theorem 6.1 from page 78 of [7]. Let φ be a map from an at most n-dimensional $ANE(n)$-space X into an at most n-dimensional $ANE(n)$-space Y. We shall write $f \overset{n}{\sim} g$ to denote that f and g are n-homotopic.

Assume that φ is an n-homotopy equivalence. By the definition, there exists a map $\psi\colon Y \to X$ such that $\psi \circ \varphi \overset{n}{\sim} id_X$ and $\varphi \circ \psi \overset{n}{\sim} id_Y$. The identity on X is a weak n-homotopy equivalence, hence by lemma 2.20, $\psi \circ \varphi$ is a weak n-homotopy equivalence. Hence for each $k < n$, $\psi \circ \varphi$ induces an isomorphism on $\pi_k(X)$, so φ induces a monomorphism of $\pi_k(X)$ and $\pi_k(Y)$. Again by lemma 2.20, $\varphi \circ \psi$ induces an isomorphism on $\pi_k(Y)$, hence φ induces an epimorphism of $\pi_k(X)$ and $\pi_k(Y)$. Hence φ is a weak n-homotopy equivalence.

Assume that φ is a weak n-homotopy equivalence. By theorem 4.1 on page 156 in [25] and by theorem 2.4, every n-dimensional $ANE(n)$-space is n-homotopy dominated by an at most n-dimensional complex. Hence we get diagrams $K \xrightarrow{g_K} X \xrightarrow{f_K} K$ and $L \xrightarrow{g_L} Y \xrightarrow{f_L} L$ where K, L are at most n-dimensional complexes, $g_K \circ f_K \overset{n}{\sim} id_X$ and $g_L \circ f_L \overset{n}{\sim} id_Y$. By lemma 2.22 there is a map $g'_L\colon L \to X$ such that $\varphi \circ g'_L \overset{n}{\sim} g_L$. Let $\psi = g'_L \circ f_L$. We'll show that φ and ψ are n-homotopy inverses of each other. Observe that $\varphi \circ \psi = \varphi \circ g'_L \circ f_L \overset{n}{\sim} f_L \circ g_L \overset{n}{\sim} id_Y$. By lemma 2.20, ψ is a weak n-homotopy equivalence. Hence by lemma 2.22, there is a map $g'_K\colon K \to Y$ such that $\psi \circ g'_K \overset{n}{\sim} g_K$. Therefore $\varphi \overset{n}{\sim} \varphi \circ g_K \circ f_K \overset{n}{\sim} \varphi \circ \psi \circ g'_K \circ f_K \overset{n}{\sim} g'_K \circ f_K$. So $\psi \circ \varphi \overset{n}{\sim} \psi \circ g'_K \circ f_K \overset{n}{\sim} g_K \circ f_K \overset{n}{\sim} id_X$ and φ is an n-homotopy equivalence. ☐

Theorem 2.23. *If X is an at most n-dimensional $ANE(n)$-space and A is a closed $ANE(n)$-subset of X such that the inclusion $A \subset X$ is a weak n-homotopy equivalence, then A is a retract of X. Moreover, every retraction of X onto A is a weak n-homotopy equivalence.*

Proof. [1] By Whitehead's characterization, the inclusion $\imath\colon A \subset X$ is an n-homotopy equivalence. By the definition of an n-homotopy equivalence, there exists a map

---

[1] I'm grateful to A. Chigogidze for showing me this simple proof.

$\psi\colon X \to A$ whose restriction to A is n-homotopic to the identity on A. By proposition 4.1.8 from [13], the identity on A extends over X to a retraction.

Let r be any retraction of X onto A. The composition $r \circ \imath$ is equal to $\imath$, which is a weak n-homotopy equivalence by the assumptions. Hence r must be a weak n-homotopy equivalence as well. □

Lemma 2.24. *If A_1, A_2 and $A_1 \cap A_2$ are at most n-dimensional closed $ANE(n)$-subspaces of $A_1 \cup A_2$ and the inclusion $A_1 \cap A_2 \subset A_2$ is a weak n-homotopy equivalence, then the inclusion $A_1 \subset A_1 \cup A_2$ is a weak n-homotopy equivalence.*

Proof. By theorem 2.23, there exists a retraction of A_2 onto $A_1 \cap A_2$. It extends to a retraction $r\colon A_1 \cup A_2 \to A_1$, since both A_1 and A_2 are closed in $A_1 \cup A_2$. Therefore the inclusion $A_1 \subset A_1 \cup A_2$ induces monomorphisms on all homotopy groups.

By theorem 2.4, there exists an open cover $\mathcal{U}$ of $A_1 \cup A_2$ such that every two $\mathcal{U}$-close maps from an at most $(n-1)$-dimensional Polish space are homotopic. Let $\mathcal{V}$ be an open cover of $A_1 \cup A_2$ that satisfies condition $(*)_n$ of theorem 2.8. Like we argued in the proof of lemma 2.9, the restriction of r to a small closed neigbhorhood of A_1 in $A_1 \cup A_2$ is $\mathcal{V}$-close to the identity. By the choice of $\mathcal{V}$, this restriction admits an extension to a map s from $A_1 \cup A_2$ into $A_1 \cup A_2$ that is $\mathcal{U}$-close to the identity.

Fix $k < n$ and consider a map $\varphi\colon S^k \to A_1 \cup A_2$. Let $\psi = s \circ \varphi$. By the choice of $\mathcal{U}$, ψ is homotopic to φ. By the construction of s, there exist open subsets $U, V \subset S^k$ such that $U \subset \psi^{-1}(A_1)$, $V \subset \psi^{-1}(A_2)$ and $U \cup V = S^k$. Hence we may fix subcomplexes C and D of S^k such that $C \cup D = S^k$ and $\psi(C) \subset A_1$ and $\psi(D) \subset A_2$. By Whitehead's characterization, the inclusion $A_1 \cap A_2 \subset A_2$ is an n-homotopy equivalence. By lemma 2.22, the map $\psi_{|D}$ is homotopic $\operatorname{rel} C \cap D$ with a map into $A_1 \cap A_2$. Obviously this homotopy extends to a homotopy $\operatorname{rel} C$. Therefore the inclusion $A_1 \subset A_1 \cup A_2$ induces epimorphisms on homotopy groups of dimensions less than n. We are done. □

Corollary 2.25. *If A_1, A_2 and $A_1 \cap A_2$ are at most n-dimensional closed $ANE(n)$-subspaces of a space X and inclusions $A_1 \subset X$ and $A_1 \cap A_2 \subset A_2$ are weak n-homotopy equivalences, then the inclusion $A_1 \cup A_2 \subset X$ is a weak n-homotopy equivalence.*

Proof. By lemma 2.24, the inclusion $A_1 \subset A_1 \cup A_2$ is a weak n-homotopy equivalence. So, by Whitehead's characterization, there exists a weak n-homotopy equivalence $A_1 \cup A_2 \to A_1$. We are done, by the assumption that $A_1 \subset X$ is a weak n-homotopy equivalence. □

Lemma 2.26. *If X is a separable $ANE(n)$-space and f is a map of an at most n-dimensional locally finite countable simplicial complex K into X, then there exists an at most n-dimensional locally finite countable simplicial complex L, containing K as a subcomplex, and an extension of f to a weak n-homotopy equivalence of L and X.*

Proof. This is a standard argument that follows the argument given in [20]. The only difference is that we want L to be a locally finite countable simplicial complex. Let $K_0 = K$ and let $f_0 = f$. For $m = 1, 2, \ldots, n$ we shall construct a locally finite countable simplicial complex K_m that contains K_{m-1} as a subcomplex and a map $f_m \colon K_m \to X$ that is an extension of f_{m-1} to a weak m-homotopy equivalence (note that we allow the dimension of K_m to be greater than n).

Fix $0 \leq m < n$ and assume that we already constructed $f_m \colon K_m \to X$. By the separability of X, the space of maps from S^m into X is separable. By theorem 2.4, the m-dimensional homotopy group $\pi_m(X)$ is countable. Hence we may enlarge K_m to a locally finite countable simplicial complex K'_m and extend f_m to a map $f'_m \colon K'_m \to X$ that is a weak m-homotopy equivalence and induces an epimorphism $\pi_m(f'_m)$ on mth homotopy groups. Let $\{\varphi_k \colon S^m \to K'_m\}_{k\geq 1}$ be a set of representants of the kernel of $\pi_m(f'_m)$. By the simplicial approximation theorem [21, Theorem 2C.1], we may assume that φ_k's are simplicial. Let $K''_m = K'_m \times [1, \infty)$, let $p \colon K''_m \to K'_m$ be a projection along $[1, \infty)$ and let $i_k \colon K'_m \to K'_m \times \{k\}$ be a map such that $p \circ i_k$ is the identity for each $k \geq 1$. We identify $K'_m \times \{1\}$ with K'_m so that K_m is a subcomplex of K''_m. Observe that $f''_m = f'_m \circ p$ is an extension of f'_m over K''_m that is a weak m-homotopy equivalence that induces an epimorphism on mth homotopy groups. The set $\{i_k \circ \varphi_k\}_{k\geq 1}$ is a set of representants of the kernel of $\pi_m(f''_m)$. Let $\Phi_k \colon B^{m+1} \to K''_m$ be a partial map defined on the boundary S^m of B^{m+1} to be equal to φ_k. Let K_{m+1} be a mapping cylinder of an disjoint union of Φ_k's. It is a locally finite countable simplicial complex. Since each φ_k represents an element of the kernel of $\pi_m(f''_m)$, map f''_m extends to a map $f_{m+1} \colon K_{m+1} \to X$. This map is a weak $(m+1)$-homotopy equivalence [20].

By the cellular approximation theorem [21, Corollary 4.12], the restriction of f_n to the n-dimensional skeleton of K_n is a weak n-homotopy equivalence. It is a map that we were looking for. ☐

Now we are ready to prove that every n-dimensional absolute neighborhood extensor in dimension n is n-homotopy equivalent to an open subset of ν^n. This theorem together with characterization and rigidity theorems implies the open embedding theorem for Nöbeling manifolds.

Proposition 2.27. *For every open subset U of $\mathbb{R}^{2n+1}$ the inclusion of $U \cap \nu^n$ into U is a weak n-homotopy equivalence.*

Proof. The following statement can be proved by the standard "general position" argument.

(*) *Every map from an at most n-dimensional finite simplicial complex into $\mathbb{R}^{2n+1}$ is approximable by maps into $\mathbb{R}^{2n+1} \setminus H$, where H is an arbitrary n-dimensional hyperplane. Moreover, if the map restricted to a subcomplex L has its image disjoint from H, then such approximations exist* rel L.

Let K be a finite simplicial complex, let L be a subcomplex of K and let ψ be a map from L into ν^n. Let C_ψ be a set of maps from K into $\mathbb{R}^{2n+1}$ that are equal to ψ on L. It is a closed subset of $C(X,Y)$. By (*), a subset of C_ψ consisting of maps into the complement of an n-dimensional hyperplane that is disjoint from the image of ψ is dense in C_ψ. By lemma 2.13, C_ψ has the Baire property, hence the set of maps from K into ν^n that are equal to ψ on L is dense in C_ψ.

Let φ be a map from a k-dimensional sphere into U, for $k \leq n$. By letting $K = S^k$ and $L = \emptyset$ we see that φ can be approximated by a map into $U \cap \nu^n$ and by theorem 2.4, if the approximation is close enough, then it is homotopic to φ in U. Therefore the inclusion induces epimorphisms on kth homotopy groups, for $k \leq n$.

Let φ be a map from a k-dimensional sphere into $U \cap \nu^n$ and assume that φ extends over a $(k+1)$-dimensional ball to a map Φ into U. By letting $K = B^{k+1}$ and $L = S^k$, we see that Φ can be approximated rel S^k by a map into $U \cap \nu^n$. Therefore the inclusion induces monomorphisms on kth homotopy groups, for $k < n$. We are done. □

Corollary 2.28. *The n-dimensional Nöbeling space is an absolute extensor in dimension n.*

Proof. Every point of ν^n has arbitrarily small neighborhoods homeomorphic to ν^n. By proposition 2.27, the Nöbeling space ν^n has vanishing homotopy groups of dimensions less than n. Therefore theorems 2.6 and 2.4 imply the assertion. □

Proposition 2.29. *For every separable $ANE(n)$-space X there exists an open subset U of ν^n and a weak n-homotopy equivalence $U \to X$.*

Proof. By lemma 2.26, for every separable $ANE(n)$-space X there exists an n-dimensional countable locally finite simplicial complex K and a weak n-homotopy equivalence $K \to X$. Realize K in $\mathbb{R}^{2n+1}$ in such a way that every ball around origin of the space intersects only finitely many vertices of K. With such realization it is clear that K has an open neighborhood U (in $\mathbb{R}^{2n+1}$) that is homotopy equivalent to K. By proposition 2.27, the set $U \cap \nu^n$ is weak n-homotopy equivalent to U. The proof is finished, since U is homotopy equivalent to K and K is weak n-homotopy equivalent to X. □

2.6 Z-sets

The definition of a Z-set was given by Anderson to describe a general concept of a negligible set [3]. There are many flavors of the definition to be found in the literature and most of them are equivalent in the class of Nöbeling manifolds [13, p. 191]. We are going to settle for the following one, proposed by Toruńczyk.

Definition. We say that a subset A of a space X is a *Z-set* in X if it is closed in X and if the identity on X is approximable by maps into $X \setminus A$. A countable union of

Z-sets is said to be a *Z_σ-set.* A map is said to be a *Z-embedding* if it is an embedding and its image is a Z-set in the codomain.

Theorem 2.30 ([13, Proposition 5.1.2]). *A closed subset A of an $\mathcal{N}_n$-space X is a Z-set if and only if every map from I^n into X is approximable by maps into the complement of A.*

Proposition 2.31 ([13, Corollary 5.1.6]). *Every compact subset of an $\mathcal{N}_n$-space is a Z-set.*

Proposition 2.32. *If a closed subset of an $\mathcal{N}_n$-space X is a countable union of Z-sets in X, then it is a Z-set in X.*

Proof. Let $A = \bigcup_{n\geq 1} A_n$ be a closed countable union of Z-sets in an $\mathcal{N}_n$-space X. Let $G_n = \{f \in C(I^n, X)\colon \operatorname{im} f \cap A_n = \emptyset\}$. By theorem 2.30, G_n is a dense subset of $C(I^n, X)$. By the assumption that A_n is closed, G_n is an open subset of $C(I^n, X)$. By lemma 2.13, the intersection $\bigcap_{n\geq 1} G_n$ is a dense subset of $C(I^n, X)$. By theorem 2.30, A is a Z-set in X. □

Proposition 2.33. *Let A be a closed subset of an $\mathcal{N}_n$-space X. If A is a Z-set, then for each open subset U of X, $A \cap U$ is a Z-set in U. If each point of A has a neighborhood U in X such that $A \cap U$ is a Z-set in U, then A is a Z-set in X.*

Proof. Let A be a Z-set in X. Every open subset U of X is a countable union $\bigcup_{n\geq 1} F_n$ of closed subsets of X. Directly from the definition, the intersection $F_n \cap A$ is a Z-set in X. Hence the set $C(I^n, X \setminus (F_n \cap A))$ is a dense subset of $C(I^n, X)$. The set $C(I^n, U)$ is an open subset of $C(I^n, X)$. Hence $C(I^n, U \setminus (F_n \cap A))$ is a dense subset of $C(I^n, U)$. By theorem 2.30, F_n is a Z-set in U. By proposition 2.32, $A = \bigcup_{n\geq 1} F_n$ is a Z-set in U.

Let A be a closed subset of X such that each point x of A has a neighborhood U_x such that $A \cap U_x$ is a Z-set in U_x. Every open subset U of X admits an open cover $\mathcal{U}$ such that if a map $U \to U$ is $\mathcal{U}$-close to the identity on U, then it admits a countinuous extension by the identity on $X \setminus U$ (we will elaborate on this in the proof of lemma 2.38). Hence it follows from the definition that if a closed subset of X is a Z-set in an open subset U of X, then it is a Z-set in X. Since X is separable metric, there exists a countable cover $\mathcal{U}$ of X that is a subset of $\{U_x\}_{x\in X}$. Let $\mathcal{F}$ be a closed shrinking of $\mathcal{U}$. By proposition 2.32, $A = \bigcup_{F\in\mathcal{F}} A \cap F$ is a Z-set in X. □

Remark 2.34. Propositions 2.31 and 2.33 imply that every closed locally compact subset of an $\mathcal{N}_n$-space is a Z-set.

Theorem 2.35 ([13, Proposition 5.1.7]). *If X is an $\mathcal{N}_n$-space, then every map f from at most n-dimensional Polish space into X is approximable by Z-embeddings. Moreover, if f restricted to a closed subset A of its domain is a closed Z-embedding, then such approximations exist* rel A.

We'll need the following corollaries of theorem 2.35.

Corollary 2.36. *Let h denote a homeomorphism of a Z-subset of an $\mathcal{N}_n$-space X and a Z-subset of an $\mathcal{N}_n$-space Y. If h extends to an n-homotopy equivalence of X and Y, then h^{-1} extends to an n-homotopy equivalence of Y and X. Moreover, we may require the extension of h^{-1} to be a closed embedding.*

Proof. By theorem 2.35 the n-homotopy equivalence of X and Y can be approximated $\operatorname{rel} Z_1$ by a closed embedding $h_1 \colon X \to Y$. By theorem 2.4 a sufficiently close approximation of an n-homotopy equivalence is an n-homotopy equivalence, therefore we may assume that h_1 is an n-homotopy equivalence. By theorem 2.23 there exists a retraction $r_1 \colon Y \to h_1(X)$. The map $h_1^{-1} \circ r_1$ is an extension of h^{-1} to an n-homotopy equivalence of Y and X. □

Corollary 2.37. *If D is a Z_σ-set in an $\mathcal{N}_n$-space Y, then every map f from an n-dimensional Polish space into Y is approximable by closed embeddings with images disjoint from D. If f restricted to a closed subset A of its domain is a Z-embedding with image disjoint from D, then such approximations exist* $\operatorname{rel} A$.

Proof. Let $C(X, Y)$ denote the space of maps from X into Y endowed with the limitation topology. It is easy to verify that the set G of Z-embeddings is a G_δ in $C(X, Y)$. Let $D = \bigcup Z_n$ and assume that every Z_n is a Z-set in Y. Let $G_n = \{f \in G \colon f(X) \cap Z_n = \emptyset\}$. Let F denote the set of maps equal to f on A. By theorem 2.35 the set $F \cap G_n$ is dense in F. Therefore by Toruńczyk's lemma, the set $F \cap \bigcap G_n$ is dense in F and we are done. □

Lemma 2.38. *Every open subset U of a metric space Y admits an open cover $\mathcal{U}$ such that if V is an open neighborhood of $Y \setminus U$, f is a map from a metric space into Y whose restriction to $f^{-1}(V)$ is a (closed) embedding into V and a map g is a $\mathcal{U}$-approximation of the restriction of f to $f^{-1}(U)$ by a (closed) embedding into U, then $f_{|f^{-1}(Y \setminus U)} \cup g$ is a (closed) embedding into Y.*

Proof. Let A be the complement of U in Y. Let $\mathcal{U}$ be a cover (we allow it to be uncountable) of U by a collection of balls such that each ball from $\mathcal{U}$ has a radius equal to one-fifth of the distance from its center to A. Let V be an open neighborhood of A and let f be a map from a metric space X into Y whose restriction to $f^{-1}(V)$ is an embedding into V. Let $g \colon f^{-1}(U) \to U$ be a $\mathcal{U}$-approximation of the restriction of f to $f^{-1}(U)$ by an embedding into U. We shall prove that $h = f_{|f^{-1}(A)} \cup g$ is an embedding of X into Y.

The image of $f_{|f^{-1}(A)}$ is a subset of A, the image of g lies in the complement of A and both of these maps are one-to-one, hence h is one-to-one. Let $x_0, x_1, \ldots$ be a sequence in X. If $h(x_0) \in U$, then $x_n \to x_0$ iff $g(x_n) \to g(x_0)$ (by which we understand that $g(x_n)$ is defined for almost all n and converges to $g(x_0)$) iff $h(x_n) \to h(x_0)$

(because U is open). If $h(x_0) \in A$, then

$$(*) \qquad \frac{2}{3}d(f(x_n), f(x_0)) \leq d(h(x_n), h(x_0)) \leq \frac{3}{2}d(f(x_n), f(x_0)),$$

by the choice of $\mathcal{U}$. Hence if $h(x_0) \in A$, then $x_n \to x_0$ iff $f(x_n) \to f(x_0)$ (because f is an embedding on $f^{-1}(V)$) iff $h(x_n) \to h(x_0)$ (by (*)). Hence

$$\lim_{n\to\infty} x_n = x_0 \Leftrightarrow \lim_{n\to\infty} h(x_n) = h(x_0),$$

so h is a homeomorphism.

Assume that the restriction of f to $f^{-1}(V)$ is a closed embedding into V and that g is a closed embedding into U. To show that h is a closed embedding it suffices to show that $h(X)$ is closed in Y. If $y \in A$, then $h(x_n) \to y$ iff $f(x_n) \to y$, by (*). Hence $A \cap \mathrm{Cl}_X h(X) = A \cap \mathrm{Cl}_X f(X) = A \cap f(X) = A \cap h(X)$. If $y \in U$, then $h(x_n) \to y$ iff $g(x_n) \to y$. Hence $U \cap \mathrm{Cl}_X h(X) = U \cap \mathrm{Cl}_X g(X) = U \cap g(X) = U \cap h(X)$. Therefore $h(X)$ is equal to its closure. We are done. □

Proposition 2.39. *Assume that V is an open subset of an $\mathcal{N}_n$-space Y and f is a map from an at most n-dimensional Polish space X into Y. If f is a closed embedding into V on $f^{-1}(V)$ and A is a closed (in Y) subset of V, then f is approximable* rel $f^{-1}(A)$ *by closed embeddings. Moreover, if Z is a subset of X and $f_{|Z}$ is a Z-embedding, then such approximations exist* rel $f^{-1}(A) \cup Z$. *Even more, if Z_Y is a Z_σ-set in Y and Z_Y is disjoint from $f(Z)$, then we may additionally require that the approximations have images disjoint from Z_Y.*

Proof. Let $\mathcal{V}$ be an open cover of Y. Let $U = Y \setminus A$ and let $\mathcal{U}$ be an open cover of U obtained via lemma 2.38 applied to U and Y. Without a loss of generality, we may assume that $\mathcal{U}$ refines $\mathcal{V}$. By proposition 2.33, the restriction of f to $Z \cap f^{-1}(U)$ is a Z-embedding into U. By theorem 2.16, U is strongly universal in dimension n, hence by corollary 2.37 applied with $D = Z_Y \cap U$, there exists a $\mathcal{U}$-approximation g of $f_{|f^{-1}(U)}$ rel $Z \cap f^{-1}(U)$ by a closed embedding into U. By lemma 2.38 and by the choice of $\mathcal{U}$, the map $g \cup f_{|f^{-1}(U)}$ is a closed embedding into Y and is $\mathcal{V}$-close to f. □

Definition. A *closed $\mathcal{N}_n$-neighborhood* of a set A is a closed set that is an $\mathcal{N}_n$-space and contains A in its interior.

Remark 2.40. Proposition 2.39 and theorem 2.16 imply that every closed subset of an $\mathcal{N}_n$-space has arbitrarily small closed $\mathcal{N}_n$-neighborhoods.

Part II

Reducing the proof of the main results to the construction of n-regular and n-semiregular $\mathcal{N}_n$-covers

Chapter 3

Approximation within an $\mathcal{N}_n$-cover

The aim of this section is to prove theorem 3.6 about approximation within closed $\mathcal{N}_n$-covers. Its proof relies on proposition 2.31, which states that every compact subset of an $\mathcal{N}_n$-space is a Z-set. This property and the theorem fail in the case of Hilbert cube manifolds [33]. Corollary 3.8 is a sum theorem for abstract Nöbeling manifolds.

Recall that $\mathcal{N}_n$ is the class of abstract n-dimensional Nöbeling manifolds (definition 2.17). We define $\mathcal{N}_n$-covers in the following way.

Definition. Let $\mathcal{C}$ be a class of topological spaces. We say that a collection is a *$\mathcal{C}$-collection* if the intersection of each non-empty collection of its elements either belongs to $\mathcal{C}$ or is empty. If the collection covers the underlying space, then we say that it is a *$\mathcal{C}$-cover*.

3.1 $Z(\mathcal{F})$-sets

In order to effectively work with closed $\mathcal{N}_n$-covers we need the following basic generalization of the concept of a Z-set.

Definition. Let $\mathcal{F}$ be a collection of subsets of a space X and let A be a subset of X. We say that A is a *$Z(\mathcal{F})$-set in X* if it is a Z-set in X and if for each subset $\mathcal{E}$ of $\mathcal{F}$ the intersection $A \cap \bigcap \mathcal{E}$ is a Z-set in $\bigcap \mathcal{E}$. We say that A is a *$Z(\mathcal{F})$-set* if it is a $Z(\mathcal{F})$-set in $\bigcup \mathcal{F}$. We say that a map is a *$Z(\mathcal{F})$-embedding into X* if it is an embedding and its image is a $Z(\mathcal{F})$-set in X. We say that it is a *$Z(\mathcal{F})$-embedding* if it is a $Z(\mathcal{F})$-embedding into $\bigcup \mathcal{F}$.

Remark 3.1. By proposition 2.33, if $\mathcal{U}$ is a collection of open subsets of a space X, then every Z-set in X is a $Z(\mathcal{U})$-set in X.

The following propositions are direct corollaries of propositions 2.31, 2.32 and 2.33.

Proposition 3.2. *If A is a compact subset of a space X and $\mathcal{F}$ is a closed $\mathcal{N}_n$-cover of X, then A is a $Z(\mathcal{F})$-set.*

Proposition 3.3. *If a closed subset of a space X is a countable union of $Z(\mathcal{F})$-sets in X, then it is a $Z(\mathcal{F})$-set in X.*

Recall that by definition 2.1, $\mathcal{F}/U = \{F_i \cap U\}_{i\in I}$ for each collection $\mathcal{F} = \{F_i\}_{i\in I}$ and each set U.

Proposition 3.4. *Let A be a closed subset of a space X and let $\mathcal{F}$ be a closed cover of X. If A is a $Z(\mathcal{F})$-set, then for each open subset U of X, $A \cap U$ is a $Z(\mathcal{F}/U)$-set. If each point of A has a neighborhood U in X such that $A \cap U$ is a $Z(\mathcal{F}/U)$-set, then A is a $Z(\mathcal{F})$-set.*

Remark 3.5. Propositions 3.2 and 3.4 imply that if $\mathcal{F}$ is a closed $\mathcal{N}_n$-cover of a space X, then every closed locally compact subset of X is a $Z(\mathcal{F})$-set.

3.2 Approximation within a cover

Definition. Assume that $\mathcal{U}$ and $\mathcal{F}$ are covers of a space Y and that $\mathcal{U}$ is open. We say that a map $g\colon X \to Y$ is a *$\mathcal{U}$-approximation within $\mathcal{F}$* of a map $f\colon X \to Y$ if it is $\mathcal{U}$-close to f and $gf^{-1}(F) \subset F$ for each $F \in \mathcal{F}$. We say that f is *approximable within $\mathcal{F}$* by embeddings (closed embeddings, Z-embeddings, etc.) if for each open cover $\mathcal{U}$ it admits a $\mathcal{U}$-approximation within $\mathcal{F}$ that is an embedding (closed embedding, Z-embedding, etc.).

Note the asymmetry: the assumption that g is a $\mathcal{U}$-approximation of f within $\mathcal{F}$ doesn't imply that f is a $\mathcal{U}$-approximation of g within $\mathcal{F}$.

Theorem 3.6. *Let $\mathcal{G}$ be a closed countable star-finite locally finite $\mathcal{N}_n$-collection in a space Y. If $\mathcal{G}$ covers Y, then every map f from an n-dimensional Polish space into Y is approximable within $\mathcal{G}$ by $Z(\mathcal{G})$-embeddings. Moreover, if f restricted to a closed subset A of its domain is a $Z(\mathcal{G})$-embedding, then the approximating $Z(\mathcal{G})$-embeddings can be taken to coincide with f on A.*

Proof. Let $\mathcal{G} = \{G_i\}_{i\in I}$ and let $G_J = \bigcap_{j\in J} G_j$ for each $J \subset I$. On this occasion, we allow J to be the empty set and let $G_\emptyset = Y$. If G_J is non-empty, then the space $C(I^n, G_J)$ of maps from I^n into G_J is separable (cf. proposition 2.5). By the assumption that $f(A)$ is a $Z(\mathcal{G})$-set, the intersection of $f(A)$ with G_J is a Z-set in G_J, hence we can pick a countable dense subset of $C(I^n, G_J)$ whose elements have images disjoint from $f(A)$. Let D denote the union of images of maps belonging to these countable dense subsets, over all $J \subset I$. By theorem 2.30,

(1) every closed embedding with image disjoint from D is a $Z(\mathcal{G})$-embedding.

Since $\mathcal{G}$ is countable and locally finite, G_J is non-empty only for countably many J's, hence D is sigma-compact. By the assumptions, G_J is an $\mathcal{N}_n$-space, hence by proposition 2.31, the intersection of D with G_J is a Z_σ-set in G_J. By (1) and by corollary 2.37,

(2) for every $J \subset I$ every map from n-dimensional Polish space into G_J is approximable by closed embeddings with images disjoint from $D \cap G_J$. Moreover, if the map restricted to a closed subset A of its domain is a closed embedding with image disjoint from $D \cap G_J$, then such approximations exist rel A.

Let $\mathcal{U}$ be an open cover of Y. We will show that the given map f can be $\mathcal{U}$-approximated within $\mathcal{G}$ by a $Z(\mathcal{G})$-embedding. Let $\mathcal{I} = \{J \subset I\colon J \neq \emptyset, G_J \neq \emptyset\}$. Order $\mathcal{I}$ into a sequence $\{J_k\}_{k\in\mathbb{N}}$ non increasing in the order by inclusion, i.e. such that if $J_l \supsetneq J_k$, then $l < k$. Such ordering exists because $\mathcal{G}$ is star-finite. By theorem 2.8, there is a sequence of pairs of covers $(\mathcal{V}_k, \mathcal{U}_k)$ such that for each k

(3) $\mathcal{V}_k$ and $\mathcal{U}_k$ are open covers of G_{J_k} that satisfy condition $(*)_n$ of theorem 2.8,

(4) $\operatorname{st}\mathcal{U}_k \prec \mathcal{U}$ and for every l if $J_l \supsetneq J_k$, then $\operatorname{st}\mathcal{U}_l \prec \mathcal{V}_k$.

The construction of a sequence $(\mathcal{U}_k, \mathcal{V}_k)$ starts with k's that correspond to sets J_k of cardinality one and continues backwards by a recursive application of theorem 2.8.

Let f denote the given map from an n-dimensional Polish space X into Y. Let $F_i = f^{-1}(G_i)$ for every $i \in I$. We let $g_0 = f_{|A}$ and recursively define a sequence of maps g_k in such a way that g_k extends g_{k-1} over F_{J_k} and

(5) $g_k(F_{J_k}) \subset G_{J_k}$ and g_k is $\operatorname{st}\mathcal{U}_k$-close to f on F_{J_k},

(6) g_k is a closed embedding with image disjoint from D.

In kth step of the construction we extend g_{k-1} over F_{J_k}. By (5) and by the order of J_k's, $g_{k-1}(F_{J_k}) \subset G_{J_k}$. By the construction, the intersection of F_{J_k} with the domain of g_{k-1} is equal to $A \cup \bigcup_{l<k, J_l \supset J_k} F_{J_l}$. By the choice of g_0, g_{k-1} is equal to f on the set A. By (5), g_{k-1} is $\operatorname{st}\mathcal{U}_l$-close to f on F_{J_l} for each $l < k$ such that $J_l \supset J_k$. By (4), $\operatorname{st}\mathcal{U}_l$ refines $\mathcal{V}_k$, hence g_{k-1} is $\mathcal{V}_k$-close to f on the intersection of F_{J_k} with the domain of g_{k-1}. By (3), the restriction of g_{k-1} to F_{J_k} extends over F_{J_k} to a map from F_{J_k} into G_{J_k} that is $\mathcal{U}_k$-close to f. Let $\widetilde{g}_{k-1}$ be an union of any such extension with the restriction of g_{k-1} to $B = g_{k-1}^{-1}(G_{J_k})$. By (6), the restriction of $\widetilde{g}_{k-1}$ to B has image disjoint from D. By (2), $\widetilde{g}_{k-1}$, defined on $B \cup F_{J_k}$, is approximable rel B by a closed (*) embedding into G_{J_k} that is $\operatorname{st}\mathcal{U}_k$-close to f and has image disjoint from D. We let g_k to be equal to this embedding on F_{J_k} and equal to g_{k-1} on $\operatorname{dom} g_{k-1} \setminus F_{J_k}$. The definition of B guarantees that g_k is a closed embedding. Conditions (5) and (6) are satisfied directly from the construction.

Let $g = \bigcup_{k\in\mathbb{N}} g_k$. It is a closed embedding because $\mathcal{G}$ is locally finite. By (6), it has image disjoint from D. Hence, by (1), it is a $Z(\mathcal{G})$-embedding. By (4) it is $\mathcal{U}$-close to f and by (5) it is an approximation of f within $\mathcal{G}$. Since $g_{|A}$ is equal to f, we are done. □

We can draw two important corollaries from theorem 3.6.

Corollary 3.7. *If $\mathcal{F}$ is a closed countable star-finite locally finite $\mathcal{N}_n$-cover of a space X and Z is a $Z(\mathcal{F})$-set, then every map of an n-dimensional Polish space into X is approximable within $\mathcal{F}$ by closed embeddings with images disjoint from Z.*

Proof. Let f be the given map. We can assume that the domain of f is separated from Z, so $f \cup id_Z$ is a well-defined map. By theorem 3.6 it is approximable within $\mathcal{F}$ and $\operatorname{rel} Z$ by a closed embedding. The restriction of such approximation to the domain of f is an embedding that we were looking for. □

The second corollary follows from theorem 2.7 and theorem 3.6 applied to the cover $\mathcal{G} = \{X, Y\}$ of $X \cup Y$.

Corollary 3.8. *If X, Y and $X \cap Y$ are $\mathcal{N}_n$-spaces and $X \cap Y$ is closed both in X and in Y, then $X \cup Y$ is an $\mathcal{N}_n$-space.*

3.3 Z-collections[1]

We prove that under stronger assumptions we can require in the statement of theorem 3.6 not only that the constructed map g approximates the given map f within $\mathcal{G}$, but also that f approximates g within $\mathcal{G}$. The stronger assumption is that $\mathcal{G}$ is a Z-collection in Y.

Definition. We say that a collection $\mathcal{F}$ of closed subsets of a topological space X is a *Z-collection* if every element of $\mathcal{F}$ is a Z-set in X and if for every $\mathcal{A} \subsetneq \mathcal{B} \subset \mathcal{F}$ the set $\bigcap \mathcal{B}$ is a Z-set in $\bigcap \mathcal{A}$.

Observe that a locally finite Z-collection cannot, by the definition, cover the underlying space. Hence, we need the following, modified statement.

Theorem 3.9. *If the assumption that $\mathcal{G}$ covers Y in theorem 3.6 is replaced by the assumption that Y is an $\mathcal{N}_n$-space and if we assume that $\mathcal{G}$ is a Z-collection, then the approximating map g can be taken so that $g^{-1}(G_i) = f^{-1}(G_i)$ for each $i \in I$.*

Proof. The proof is an easy modification of the proof of theorem 3.6. The necessary changes are to change condition (5) to

(5') $f^{-1}(G_{J_l}) = g_k^{-1}(G_{J_l})$ for each $l \leq k$ and g_k is $\operatorname{st}\mathcal{U}_k$-close to f on F_{J_k},

and to change the sentence marked by (*) on the margin to

> By (2), $\widetilde{g}_{k-1}$, defined on $B \cup F_{J_k}$, is approximable $\operatorname{rel} B$ by a closed embedding into G_{J_k} that is $\operatorname{st}\mathcal{U}_k$-close to f, has image disjoint from D and maps $F_{J_k} \setminus B$ into $G_{J_k} \setminus \bigcup_{l<k} G_{J_l}$.

---

[1] we shall not use the theorem proved in ths section until the third part of the paper.

By the assumption $\mathcal{G}$ is a Z-collection, hence the intersection of $\bigcup_{l<k} G_{J_l}$ with G_{J_k} is a Z-set in G_{J_k}. Hence, by corollary 2.37, such approximation exists. The assumption that the approximation maps $F_{J_k} \setminus B$ into $G_{J_k} \setminus \bigcup_{l<k} G_{J_l}$ guarantees that $f^{-1}(G_{J_l}) = g_k^{-1}(G_{J_l})$ for each $l < k$. $\square$

We will show how to construct Z-collections in $\mathcal{N}_n$-spaces in section 9.1 in chapter 9.

Chapter 4

Constructing closed $\mathcal{N}_n$-covers

4.1 Adjustment of a collection

By theorem 2.18, every open subset of an $\mathcal{N}_n$-space is an $\mathcal{N}_n$-space. Obviously the property of being an $\mathcal{N}_n$-space is not inherited by closed subsets of $\mathcal{N}_n$-spaces. However by the strong universality of $\mathcal{N}_n$-spaces, the inclusion of an arbitrary open subset of an $\mathcal{N}_n$-space is approximable by closed embeddings and images thereof are closed $\mathcal{N}_n$-subsets of the space.

Let U be an open and A a closed neighborhood of a point x in an $\mathcal{N}_n$-space X such that $A \subset U$. By proposition 2.39, the inclusion of U into X can be approximated $\operatorname{rel} A$ by a closed embedding. The image of such an approximation (an *adjustment* of U) is a closed $\mathcal{N}_n$-neighborhood of x. In the sequel, we shall apply this technique to collections of sets, in a way described by the following definition.

Definition. Let $\mathcal{F} = \{F_i\}_{i \in I}$ and $\mathcal{G} = \{G_i\}_{i \in I}$ be arbitrary collections of subsets of a space X and let $\mathcal{U}$ be an open cover of X.

(a) We say that $\mathcal{G}$ is a *$\mathcal{U}$-adjustment* of $\mathcal{F}$ if for each $J \subset I$ there is a homeomorphism of $F_J = \bigcap_{j \in J} F_j$ onto $G_J = \bigcap_{j \in J} G_j$ that is $\mathcal{U}$-close to the inclusion $F_J \subset X$.

(b) If each of these homeomorphisms can be taken to be the identity on a closed set A, then we say that $\mathcal{G}$ is an *adjustment of $\mathcal{F}$* $\operatorname{rel} A$.

(c) If additionally $G_i = F_i$ for every i in a given set $J \subset I$, then we say that $\mathcal{G}$ is an *adjustment of $\mathcal{F}$ with fixed J*.

A convenient property of the notion of an adjustment is that for an arbitrary class $\mathcal{C}$ of topological spaces every adjustment of a $\mathcal{C}$-collection is a $\mathcal{C}$-collection.

Lemma 4.1. *For each open subset U of a metric space X there exists an open cover $\mathcal{U}$ of U such that if A is a (closed) subset of X, then every $\mathcal{U}$-approximation of the inclusion $A \cap U \subset U$ by a (closed) embedding of $A \cap U$ into U extends over A, by the* *identity on $A \setminus U$, to a (closed) embedding of A into X. Hence every (closed in U) $\mathcal{U}$-adjustment of a restriction $\mathcal{F}/U$ of a (closed) collection $\mathcal{F}$ of subsets of X extends in a natural way to a (closed) adjustment* $\operatorname{rel} X \setminus U$ *of the entire collection.*

Proof. The first statement follows from lemma 2.38 applied to the inclusion $A \subset X$. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a (closed) collection of subsets of X and let $\mathcal{G} = \{G_i\}_{i\in I}$ be a (closed in U) $\mathcal{U}$-approximation of $\mathcal{F}/U = \{F_i \cap U\}_{i\in I}$. We have to show that the collection $\{G_i \cup (F_i \setminus U)\}_{i\in I}$ is a (closed) $\mathcal{U}$-adjustment $\operatorname{rel} X \setminus U$ of $\mathcal{F}$. Since $\mathcal{G}$ is a $\mathcal{U}$-adjustment of $\mathcal{F}/U$, for each $J \subset I$ there exists a homeomorphism h_J from $F_J \cap U$ onto G_J that is $\mathcal{U}$-close to the inclusion of $F_J \cap U$ into U. By the first statement, the map $h_J \cup (F_J \setminus U)$ is a (closed) embedding. This embedding is $\mathcal{U}$-close to the inclusion and its image is equal to $G_J \cup (F_J \setminus U)$. We are done. □

4.2 Limits of sequences of adjustments

Definition 4.2. For every sequence $\mathcal{F}_k = \{F_i^k\}_{i\in I}$ we let

$$\lim_{k\to\infty} \mathcal{F}_k = \{\bigcap_{j\geq 1}\bigcup_{k\geq j} F_i^k\}_{i\in I}.$$

Proposition 4.3. *Assume that $\mathcal{V} = \{V_k\}_{k\geq 1}$ is a locally finite collection of open subsets of a Polish space X. For every open cover $\mathcal{U}$ of X there exists a sequence $\mathcal{U}_1, \mathcal{U}_2, \ldots$ of open covers of X such that if $\mathcal{F}_k$ is a $\mathcal{U}_k$-adjustment* $\operatorname{rel} X \setminus V_k$ *of $\mathcal{F}_{k-1}$, then $\lim_{k\to\infty} \mathcal{F}_k$ is a $\mathcal{U}$-adjustment of $\mathcal{F}_0$.*

Proof. Without a loss of generality we may assume that a second star of each element of $\mathcal{U}$ intersects only finitely many elements of $\mathcal{V}$. Let $\mathcal{U}_1, \mathcal{U}_2, \ldots$ be a sequence of covers of X obtained via lemma 2.19 applied to $\mathcal{U}$. We will show that it satisfies our claim.

Let $\mathcal{F}_k = \{F_i^k\}_{i\in I}$ and $\mathcal{F}_\infty = \lim_{k\to\infty} \mathcal{F}_k$. Let $F_J^k = \bigcap_{j\in J} F_j^k$. By the assumptions, for each $J \subset I$ there exists a homeomorphism $h_J^k \colon F_J^{k-1} \to F_J^k$ that is $\mathcal{U}_k$-close to the inclusion of F_J^{k-1} into X. By lemma 2.19 and by the choice of $\mathcal{U}_k$'s, the limit $h_J^\infty = \lim_{k\to\infty} h_J^k \circ \cdots \circ h_J^2 \circ h_J^1$ exists, is continuous and $\mathcal{U}$-close to the inclusion of F_J^0 into X.

Observe that if a map is $\mathcal{U}$-close to the identity, then to check that it is a homeomorphism (onto its image) it suffices to check it is a homeomorphism on $\operatorname{st}_{\mathcal{U}} U$ for every U in $\mathcal{U}$. By the choice of $\mathcal{U}$, for each U in $\mathcal{U}$ the second star $\operatorname{st}^2_{\mathcal{U}} U$ intersects only finitely many elements of $\mathcal{V}$. Since each $h_J^k \circ \cdots \circ h_J^2 \circ h_J^1$ is $\mathcal{U}$-close to the identity, the sequence $h_k^J \circ \cdots \circ h_1^J \circ h_0^J$ restricted to $F_J^0 \cap \operatorname{st}_{\mathcal{U}} U$ stabilizes after finitely many steps. By the assumption every h_k^J is a homeomorphism, so h_J^∞ restricted to $\operatorname{st}_{\mathcal{U}} U$ is a homeomorphism. We are done, because the image of h_J^∞ is equal to F_J^∞, by local finiteness of $\mathcal{V}$. □

4.3 Construction of a closed $\mathcal{N}_n$-swelling

Our first application of the technique of adjustments is a construction of a closed interior $\mathcal{N}_n$-cover given some constraints on its size and structure.

Lemma 4.4. *If $\mathcal{G}$ is a closed star-finite cover of an $\mathcal{N}_n$-space Y and $\mathcal{V}$ is an open swelling of $\mathcal{G}$, then for each open cover $\mathcal{U}$ of Y and each Z-set Z in Y there exists a closed $\mathcal{U}$-adjustment $\mathcal{F}$ of $\mathcal{V}$ that is a swelling of $\mathcal{G}$ and such that Z is a $Z(\mathcal{F})$-set.*

Observe that the assumption that $\mathcal{G}$ is star-finite and has an open swelling implies that it is locally finite.

Proof. Let $\mathcal{G} = \{G_i\}_{i\in I}$, $\mathcal{V} = \{V_i\}_{i\in I}$ and $V_J = \bigcap_{j\in J} V_j$ for each non-empty subset J of I. Let $\mathcal{I} = \{J \subset I \colon J \neq \emptyset \text{ and } V_J \neq \emptyset\}$. Order $\mathcal{I}$ into a sequence $\{J_k\}_{k\geq 1}$ non increasing in the order by inclusion. Such ordering exists because $\mathcal{V}$ is star-finite. Let $W_k = \bigcup_{i\in I\setminus J_k} V_i$. Let B_k be a boundary of $V_{J_k} \setminus W_k$ taken in V_{J_k} (it is not necessarily closed in Y). Let

$$U_k = (V_{J_k} \setminus \bigcup_{i\in I\setminus J_k} G_i) \cap \{y \in Y \colon d(y, B_k) < d(y, Y \setminus V_{J_k})\},$$

where $d(y, A)$ is the distance of a point y from a set A (we let $d(y, \emptyset) = \infty$). By the assumptions, $\mathcal{V}$ is a swelling of $\mathcal{G}$, so $\bigcup_{i\in I\setminus J_k} G_i \subset W_k$. Hence U_k is an open neighborhood of B_k. We shall prove that if U_k intersects U_l, then either $J_k \subset J_l$ or $J_l \subset J_k$. Let y be an element of $U_k \cap U_l$. If $B_k \cap V_{J_l} = \emptyset$ and $B_l \cap V_{J_k} = \emptyset$, then $d(y, B_k) < d(y, Y \setminus V_{J_k}) \leq d(y, B_l) < d(y, Y \setminus V_{J_l}) \leq d(y, B_k)$ - a contradiction. Hence either B_k intersects V_{J_l} or B_l intersects V_{J_k}. Assume the former case, i.e. $B_k \cap V_{J_l} \neq \emptyset$. By the definition, B_k is disjoint from W_k. Since $V_{J_l} = \bigcap_{i\in J_l} V_i$ intersects B_k, it is not a subset of $W_k = \bigcup_{i\in I\setminus J_k} V_i$, hence J_l must be disjoint from $I \setminus J_k$, so it must be a subset of J_k. Same argument applies in the latter case, i.e. when $B_l \cap V_{J_k} \neq \emptyset$.

Let $\mathcal{U}_1, \mathcal{U}_2, \ldots$ be a sequence of covers of Y obtained via proposition 4.3 applied to a locally finite collection $\mathcal{V}$ and an open cover $\mathcal{U}$ of Y. If P and Q are open subsets of Y, then the boundary of $P \setminus Q$ taken in P is equal to the boundary of Q taken in $P \cup Q$. We let $P = V_{J_k}$ and $Q = W_k$, and obtain the equality $B_k = \mathrm{Bd}_{V_{J_k}\cup W_k} W_k$. Since U_k is a neighborhood of B_k, $W_k \setminus U_k$ is closed in $W_k \cup V_{J_k}$. By theorem 2.18, $W_k \cup V_{J_k}$ is an $\mathcal{N}_n$-space. Therefore, by proposition 2.39 applied to $A = W_k \setminus U_k$, $V = W_k$ and $Z_Y = Z \cap U_k$, the inclusion $W_k \subset W_k \cup V_{J_k}$ has a $\mathcal{U}_k$-approximation $h_k \colon W_k \to W_k \cup V_{J_k}$ such that the following condition is satisfied.

(h_k) h_k is a closed embedding into $W_k \cup V_{J_k}$, is $\mathcal{U}_k$-close to the inclusion, is equal to the inclusion on $W_k \setminus U_k$ and has image disjoint from $Z \cap U_k$.

We shall recursively construct a sequence $\mathcal{F}_k = \{F_i^k\}_{i\in I}$ of covers of Y, starting with $\mathcal{F}_0 = \mathcal{V}$, such that for $k > 0$ the following conditions are satisfied.

(1_k) $\mathcal{F}_k$ is a $\mathcal{U}_k$-adjustment of $\mathcal{F}_{k-1}$ $\operatorname{rel} Y \setminus U_k$ with fixed J_k.

(2_k) $\mathcal{F}_k$ restricted to $\bigcup_{0<l\leq k} V_{J_l}$ is a closed cover, Z is a $Z(\mathcal{F}_k)$-set and $\mathcal{G}$ is a shrinking of $\mathcal{F}_k$.

Observe that Z is a $Z(\mathcal{F}_0)$-set by remark 3.1 and $\mathcal{G}$ is a shrinking of $\mathcal{F}_0$. Before we construct the sequence, we prove that for each $k \geq 0$ if (1_l) and (2_l) are satisfied for each l greater than 0 and less than or equal to k, then the following conditions hold:

(3_k) $\bigcup_{i\in I\setminus J_{k+1}} F_i^k$ is a subset of W_{k+1},

(4_k) $U_{k+1} \cup (V_{J_{k+1}} \setminus W_{k+1})$ is a subset of $F^k_{J_{k+1}}$.

To prove (3_k) fix $j \in I \setminus J_{k+1}$. We will show by induction on l that $F_j^l \subset W_{k+1}$ for each $0 \leq l \leq k$. It is true for $l = 0$, because $F_j^0 = V_j$. Let $l > 0$. By (1_l), $\mathcal{F}_l$ is equal to $\mathcal{F}_{l-1}$ on the complement of U_l, hence $F_j^l \subset F_j^{l-1} \cup U_l$. By the inductive assumption $F_j^{l-1} \subset W_{k+1}$. If $l \leq k$, then $J_l \setminus J_{k+1}$ is non-empty, by the order of J_k's. Hence $U_l \subset V_{J_l} \subset V_{J_l \setminus J_{k+1}} \subset W_{k+1}$. Hence $F_j^l \subset F_j^{l-1} \cup U_l \subset W_{k+1}$.

We will show by induction on l that $U_{k+1} \cup (V_{J_{k+1}} \setminus W_{k+1})$ is a subset of $F^l_{J_{k+1}}$ for each $0 \leq l \leq k$, thus proving (4_k). From the definition $U_{k+1} \subset V_{J_{k+1}} = F^0_{J_{k+1}}$, so the assertion is true for $l = 0$. Let $0 < l \leq k$. We either have $J_{k+1} \not\subset J_l$ or or $J_{k+1} \subset J_l$. By (1_l), $\mathcal{F}_l$ is an adjustment of $\mathcal{F}_{l-1}$ with fixed J_l so in the latter case, $F^l_{J_{k+1}} = F^{l-1}_{J_{k+1}}$ and the assertion is true. In the former case, U_l is disjoint from U_{k+1} and $U_l \subset V_{J_l} \subset W_{k+1}$. Hence $U_{k+1} \cup (V_{J_{k+1}} \setminus W_{k+1})$ is disjoint from U_l. By (1_l), $\mathcal{F}_l$ is equal to $\mathcal{F}_{l-1}$ on the complement of U_l, so the inductive step is done.

The construction. Fix $k > 0$ and assume that we already constructed $\mathcal{F}_{k-1}$. Let

$$F_i^k = \begin{cases} h_k(F_i^{k-1}) & i \in I \setminus J_k \\ F_i^{k-1} & i \in J_k. \end{cases}$$

Consider $J \subset I$. If $J \subset J_k$, then $F_J^k = F_J^{k-1}$. If $J \not\subset J_k$, then a direct computation using (4_{k-1}) and (h_k) shows that $F_J^k = h_k(F_J^{k-1})$. Hence by (3_{k-1}), the restriction of h_k to F_J^{k-1} is a homeomorphism onto F_J^k. By the definition of a $\mathcal{U}_k$-adjustment, (1_k) is satisfied.

By (h_k), h_k is a closed embedding into $W_k \cup V_{J_k}$. Hence its restriction to $h_k^{-1}(\bigcup_{0<l\leq k} V_{J_l}) = \bigcup_{0<l<k} V_{J_l}$ is a closed embedding into $\bigcup_{0<l\leq k} V_{J_l}$. By ($2_{k-1}$), $\mathcal{F}_{k-1}$ restricted to $\bigcup_{0<l<k} V_{J_l}$ is closed. Hence for each $i \in I \setminus J_k$, $F_i^k \cap \bigcup_{0<l\leq k} V_{J_l} = h_k(F_i^{k-1} \cap \bigcup_{0<l<k} V_{J_l})$ is closed in $\bigcup_{0<l\leq k} V_{J_l}$. If $i \in J_k$, then $F_i^k = F_i^{k-1}$. By (4_{k-1}), $V_{J_k} \setminus \bigcup_{0<l<k} V_{J_l}$ is a subset of F_i^{k-1}. The intersection of F_i^{k-1} with $\bigcup_{0<l<k} V_{J_l}$ is closed in $\bigcup_{0<l<k} V_{J_l}$, so $F_i^{k-1} \cap \bigcup_{0<l\leq k} V_{J_l} = (V_{J_k} \setminus \bigcup_{0<l<k} V_{J_l}) \cup (F_i^{k-1} \cap \bigcup_{0<l<k} V_{J_l})$ is closed in $\bigcup_{0<l\leq k} V_{J_l}$. Hence $\mathcal{F}_k$ restricted to $\bigcup_{0<l\leq k} V_{J_l}$ is a closed cover.

By (2_{k-1}), $G_i \subset F_i^{k-1}$ for each $i \in I$. If $i \in J_k$, then $F_i^k = F_i^{k-1}$, so $G_i \subset F_i^k$. If $i \in I \setminus J_k$, then G_i is disjoint from U_k by the definition. By (1_k), F_i^k is equal to F_i^{k-1} on the complement of U_k, hence $G_i \subset F_i^k$. Therefore $\mathcal{G}$ refines $\mathcal{F}_k$.

By (2_{k-1}), Z is a $Z(\mathcal{F}_{k-1})$-set. Hence for each $J \subset I$, $Z \cap F_J^{k-1}$ is a Z-set in F_J^{k-1}. If $J \subset J_k$, then $F_J^k = F_J^{k-1}$, so $Z \cap F_J^k$ is a Z-set in F_J^k as well. If $J \not\subset J_k$, then $F_J^k = h_k(F_J^{k-1})$. By (h_k), the image of h_k is disjoint from $Z \cap U_k$ and h_k is equal to the inclusion on the complement of U_k. Hence $Z \cap F_J^k = Z \cap (F_J^{k-1} \setminus U_k)$. By (2_{k-1}), the latter set is a Z-set in F_J^{k-1}, hence its image under h_k is a Z-set in $h_k(F_J^{k-1}) = F_J^k$. But h_k is the identity map on this set, hence $Z \cap F_J^k$ is a Z-set in F_J^k. Hence Z is a $Z(\mathcal{F}_k)$-set.

By (1_k), by proposition 4.3 and by the choice of $\mathcal{U}_k$'s, the limit $\mathcal{F} = \lim_{k\to\infty} \mathcal{F}_k$ exists and is a $\mathcal{U}$-adjustment of $\mathcal{F}_0 = \mathcal{V}$. By (3_k), $\mathcal{F}$ is closed. By (2_k), it is a swelling of $\mathcal{G}$. By (2_k) and by proposition 3.4, Z is a $Z(\mathcal{F})$-set. We are done. $\square$

Theorem 4.5. *Every closed star-finite locally finite cover that refines an open cover $\mathcal{U}$ of an $\mathcal{N}_n$-space X has a swelling $\mathcal{F}$ that refines $\mathcal{U}$ and is a closed locally finite interior $\mathcal{N}_n$-cover. Moreover, we may require that a specified Z-set is a $Z(\mathcal{F})$-set.*

Proof. Let $\mathcal{U}$ be an open cover of an $\mathcal{N}_n$-space X and let $\mathcal{G} = \{G_i\}_{i\in I}$ be a closed star-finite locally finite refinement of $\mathcal{U}$. We may assume that $\mathcal{G}$ is an interior cover [18, Exercise 7.1.G]. By the same argument, there exists an open locally finite swelling $\mathcal{V} = \{V_i\}_{i\in I}$ of $\mathcal{G}$ such that the closure of each element of $\mathcal{V}$ lies in an element of $\mathcal{U}$. Let Z be an arbitrary Z-set in X. By lemma 4.4, $\mathcal{V}$ can be adjusted to a closed cover $\mathcal{F}$ of X such that Z is a $Z(\mathcal{F})$-set and $\mathcal{G}$ is a refinement of $\mathcal{F}$. Obviously, if the adjustment is small enough, then $\mathcal{F}$ refines $\mathcal{U}$ and we are done. $\square$

Chapter 5

Carrier and nerve theorems

Almost entire chapter is cited word-for-word after [32] and it is included here for completeness.

5.1 Regular covers

An efficient way to investigate properties of a topological space is to divide it into pieces and examine how they are glued together. We show how to divide a general topological space and endow it with a structure that resembles a triangulation.

In order to divide a space that belongs to a class $\mathcal{C}$ of topological spaces we must decide what a piece is. We want it to resemble a simplex as much as possible. As a simplex is an archetype of an absolute extensor, the choice of absolute extensors for $\mathcal{C}$ as pieces is quite natural.

Definition. A space Y is an *absolute extensor* for a space X if each map from a closed subset of X into Y extends over the entire space X. The class of absolute extensors for all spaces from a class $\mathcal{C}$ is denoted by $AE(\mathcal{C})$. We write $AE(X)$ for $AE(\{X\})$.

The following defines regular covers that endow a space with structures similar to triangulations.

Definition. Let $\mathcal{C}$ be a class of topological spaces. A locally finite locally finite-dimensional closed $AE(\mathcal{C})$-cover is said to be *regular for the class* $\mathcal{C}$. Recall that a cover is *locally finite dimensional* if its nerve is such.

Examples of regular covers include a locally finite and locally finite dimensional cover of an Euclidean space by its closed balls and a cover of a finite simplicial complex by its simplices.

5.2 Carrier theorem

How to extend a partial map from a subcomplex over the entire CW complex? If each map from the boundary of an Euclidean ball into the codomain extends over the ball, then the answer is easy: order cells by inclusion and construct an extension inductively. But the asphericity of the codomain (vanishing of all its homotopy groups) is a rare luxury. The same technique would work though if we were able to restrict ranges of the map on individual cells of the CW complex to aspherical subspaces of the codomain. This idea leads to the notion of a carrier and to the aspherical carrier theorem [30, II §9]. We generalize this notion to arbitrary spaces using a cover to replace the cell structure in the domain.

Definition. A *carrier* is a function $C\colon \mathcal{F} \to \mathcal{G}$ from a cover $\mathcal{F}$ of a space X into a collection $\mathcal{G}$ of subsets of a topological space such that for each $\mathcal{A} \subset \mathcal{F}$ if $\bigcap \mathcal{A} \neq \emptyset$, then $\bigcap_{A \in \mathcal{A}} C(A) \neq \emptyset$. We say that a map f is *carried by* C if it is defined on a closed subset of X and $f(F) \subset C(F)$ for each $F \in \mathcal{F}$.

Carrier theorem. *Assume that $C\colon \mathcal{F} \to \mathcal{G}$ is a carrier such that $\mathcal{F}$ is a closed cover of a space X and $\mathcal{G}$ is an $AE(X)$-cover of another space. If $\mathcal{F}$ is locally finite and locally finite dimensional, then each map carried by C extends to a map of the entire space X, also carried by C.*

Special cases of the carrier theorem follow from Michael's selection theorem, as the multivalued map given by the formula $F(x) = \bigcap_{F \ni x} C(F)$ is lower semi-continuous.

Proof. Let f_0 be a map carried by C and let $A = \operatorname{dom} f_0$. Let $\{\mathcal{E}_\gamma\}_{0<\gamma<\Gamma}$ be a transfinite sequence of all subcollections of $\mathcal{F}$ with non-empty intersections, such that the sequence $\{\delta_\gamma = \bigcap \mathcal{E}_\gamma\}_{0<\gamma<\Gamma}$ is non decreasing in the order by inclusion. Its existence is guaranteed by the assumption of local finite dimensionality of $\mathcal{F}$. Let $C_\gamma = \bigcap_{E \in \mathcal{E}_\gamma} C(E)$ and observe that an arbitrary map f is carried by C if and only if $f(\delta_\gamma) \subset C_\gamma$ for each $0 < \gamma < \Gamma$. Let $\delta_\Gamma = \emptyset$. We shall construct a transfinite sequence of maps $\{f_\gamma\colon A \cup \bigcup_{0<\alpha\leq\gamma} \delta_\alpha \to Y\}_{\gamma \leq \Gamma}$ such that f_α extends f_β for all $0 \leq \beta \leq \alpha \leq \Gamma$ and $f_\gamma(\delta_\gamma) \subset C_\gamma$ for each $0 < \gamma < \Gamma$. The map f_Γ will be an extension that we are looking for, since $\bigcup_{\gamma<\Gamma} \delta_\gamma = X$.

We proceed by transfinite induction. Fix $\gamma \leq \Gamma$ and assume that for each $\alpha < \gamma$ we already constructed f_α. The map $f'_\gamma = \bigcup_{\alpha<\gamma} f_\alpha$ is well defined, continuous and its domain is closed in X because maps f_α agree on intersections of their domains and $\mathcal{F}$ is closed and locally finite. If $\gamma = \Gamma$, then $\delta_\gamma = \emptyset$ and we may put $f_\Gamma = f'_\Gamma$. If $\gamma < \Gamma$, then by the order of δ_γ and by inductive assumptions f'_γ maps δ_γ into C_γ. The set C_γ is non-empty because C is a carrier and is an absolute extensor for δ_γ because $\mathcal{G}$ is an $AE(X)$-cover. So f'_γ extends onto δ_γ to a map f_γ such that $f_\gamma(\delta_\gamma) \subset C_\gamma$ and our construction is finished. ☐

Definition. A *cover is regular for a space* X if it is regular for the class $\{X\}$.

Corollary 5.1. *If a closed cover $\mathcal{G}$ of a space Y is regular for $X \times [0,1]$, then every two $\mathcal{G}$-close maps from X into Y are $\mathcal{G}$-homotopic. Moreover such homotopy exists with the additional property that if endpoints of its path lie in an element of $\mathcal{G}$, then the entire path lies in it.*

Proof. Let f and g be two $\mathcal{G}$-close maps from X into Y. Let $\mathcal{F} = \{F_G\}_{G \in \mathcal{G}}$ be the collection of subsets of $X \times [0,1]$ defined by the formula

$$F_G = (f^{-1}(G) \cap g^{-1}(G)) \times [0,1].$$

It is a cover of $X \times [0,1]$ because f and g are $\mathcal{G}$-close. Define a carrier $C \colon \mathcal{F} \to \mathcal{G}$ by the formula $C(F_G) = G$ and a map $F \colon X \times \{0,1\} \to Y$ by $F(x,0) = f(x)$ and $F(x,1) = g(x)$. By the definition, F is carried by C and by the carrier theorem, it admits an extension over the entire space $X \times [0,1]$, also carried by C. This extension is a $\mathcal{G}$-homotopy that satisfies our claim. □

5.3 Nerve theorem

Nerve theorems give conditions under which the nerve of a cover is equivalent to the underlying space. First examples of such theorems are attributed to K. Borsuk [10, p. 234] (for closed covers) and A. Weil [37, p. 141] (for open covers), both for homotopy equivalences. Since then several generalizations were made. First generalizations by W. Holsztyński [24] and J. N. Haimov [19] relaxed conditions on the cover. Next weak homotopy equivalences were studied in this context by M. McCord [31] and weak n-homotopy equivalences by A. Björner [9].

The nerve theorem that we shall use in the present paper is proved in the next chapter. Here we develop tools used in its proof. As an easy application, we prove a nerve theorem for closed covers (theorem 5.5), which is more general than results previously published in the literature.

Definition. A carrier is *invertible* if it is bijective and its inverse is a carrier.

Lemma 5.2. *If a collection $\mathcal{F} = \{F_i\}_{i \in I}$ is point-finite, then its nerve is well defined and a function $F_i \mapsto \operatorname{bst} v(F_i)$ is an invertible carrier, where $\operatorname{bst} v(F_i)$ is a barycentric star of a vertex of the nerve of $\mathcal{F}$ that corresponds to F_i.*

Proof. This is a reformulation of lemma 2.11. □

Lemma 5.3. *If a simplicial complex is locally finite dimensional and is endowed with either weak or metric topology, then its cover by barycentric stars of its vertices is regular for the class of metric spaces. If a simplicial complex is both locally countable and locally finite dimensional and is endowed with the metric topology, then its cover by barycentric stars of its vertices is regular for the class of normal spaces.*

Proof. Let K be a locally finite dimensional simplicial complex and let $\mathcal{B}_K$ be a cover of K by barycentric stars of its vertices. By lemma 2.11, the nerve of $\mathcal{B}_K$ is isomorphic to K. Hence $\mathcal{B}_K$ is locally finite dimensional. By lemma 2.11, $\mathcal{B}_K$ is locally finite. What is left to prove is that every intersection of elements of $\mathcal{B}_K$ is either empty or is an absolute extensor for the class of metric spaces (or for the class of normal spaces if K is countable and endowed with the metric topology). By theorem 11.7 on page 109 of [25] and by theorem 10.4 on page 105 of [25], every simplicial complex is an absolute neighborhood extensor for the class of metric spaces. By theorem 11.7 on page 109 of [25], a countable simplicial complex with the metric topology is an absolute neighborhood extensor for the class of normal spaces. By theorem 7.1 on page 43 of [25], every contractible absolute neighborhood extensor is an absolute extensor. Hence it suffices to prove that an intersection of a collection of barycentric stars of vertices of a simplicial complex is either empty, or contractible. Lemma 2.12 finishes the proof. □

Remark 5.4. Conditions listed in theorem 11.7 on page 109 of [25] allow to state other variants of lemma 5.3, i.e. if the complex is endowed with the metric topology but is not necessarily countable, then the cover is regular for the class of fully normal spaces, etc.

Putting everything together we obtain a nerve theorem for homotopy equivalences. We state the theorem for closed covers, which generalizes a nerve theorem by J. N. Haimov [19]. An analogous theorem for open covers may also be proved. We do not state it here as it turns out to be equivalent to the nerve theorem by A. Weil [37].

Theorem 5.5. *Assume that a closed cover $\mathcal{F}$ of a normal space X is regular for the class of metric spaces. If $\mathcal{F}$ is star-countable, then X and the nerve of $\mathcal{F}$ are homotopy equivalent.*

The main theorem of [19] states the same conclusion under the additional assumption that $\mathcal{F}$ is star-finite and X is paracompact. The topology on the nerve of $\mathcal{F}$ is either weak or metric; by theorem 6.4 on page 81 of [7], the identity map from a simplicial complex endowed with a weak topology into itself, endowed with a metric topology, is a homotopy equivalence.

Proof. Let $B\colon \mathcal{F} \to \mathcal{B}_{\mathcal{F}}$ be an invertible carrier as defined in lemma 5.2. By the carrier theorem and lemma 5.3 there exist $\kappa\colon X \to N(\mathcal{F})$ and $\lambda\colon N(\mathcal{F}) \to X$ carried by B and B^{-1} respectively. Then $\lambda \circ \kappa$ is carried by $B^{-1} \circ B$ so it is $\mathcal{F}$-close to id_X and by corollary 5.1 λ is a homotopy inverse of κ. Analogously κ is a homotopy inverse of λ so X and $N(\mathcal{F})$ are homotopy equivalent. □

Chapter 6

Anticanonical maps and semiregularity

6.1 A nerve theorem and the notion of semiregularity

The crux of the proof of theorem 5.5 is the proof of the existence of a map λ carried by the carrier B^{-1} defined in lemma 5.2. These maps are very important in the sequel.

Definition. A map from the nerve of a closed cover $\mathcal{F}$ into the underlying space is an *anticanonical map of* $\mathcal{F}$ if it is carried by the carrier B^{-1} defined in lemma 5.2.

Proposition 6.1. *If an $AE(n)$-cover is point-finite and has at most n-dimensional nerve, then it admits an anticanonical map.*

Proof. Let $\mathcal{F}$ be a point-finite $AE(n)$-cover with at most n-dimensional nerve. By lemma 5.3, the cover $\mathcal{B}_{\mathcal{F}}$ of the nerve of $\mathcal{F}$ by barycentric stars of its vertices is locally finite (by the definition of a regular cover). By lemma 5.2, $\mathcal{B}_{\mathcal{F}}$ is isomorphic to $\mathcal{F}$. In particular, it is locally finite dimensional. By the carrier theorem, the empty map extends over the nerve of $\mathcal{F}$ to a map carried by B^{-1}. This map is an anticanonical map of $\mathcal{F}$. ☐

The notion of an anticanonical map is dual to the notion of a canonical map. For closed covers, we define canonical maps as follows.

Definition. A map from a space into the nerve of its closed cover $\mathcal{F}$ is a *canonical map of* $\mathcal{F}$ if it is carried by the carrier B defined in lemma 5.2.

Proposition 6.2. *Every closed locally finite and locally finite dimensional cover of a metric space admits a canonical map.*

Proof. Apply lemma 5.3 and the carrier theorem. ☐

If a cover satisfies assumptions of propositions 6.1 and 6.2, then it is regular for the class of at most n-dimensional metric spaces (the assumption that its nerve is at most n-dimensional is not needed). For brevity, we employ the following definition.

Definition. We say that a closed cover is *n-regular* if it is a locally finite locally finite dimensional $AE(n)$-cover, i.e. if it is regular for the class of at most n-dimensional metric spaces.

Nerve theorem. *The n-dimensional skeleton of the nerve of a closed n-regular cover of an at most n-dimensional metric space X is n-homotopy equivalent to X. In particular, if two at most n-dimensional metric spaces admit isomorphic closed n-regular covers, then they are n-homotopy equivalent.*

Proof. We prove the second assertion first. Let X and Y be at most n-dimensional metric spaces and let $\mathcal{F} = \{F_i\}_{i\in I}$ and $\mathcal{G} = \{G_i\}_{i\in I}$ be isomorphic closed n-regular covers of X and Y respectively. The map $C\colon \mathcal{F} \to \mathcal{G}$ that assigns G_i to F_i for each $i \in I$ is an invertible carrier. By the assumption, $\mathcal{F}$ is a closed locally finite locally finite dimensional cover of X and $\mathcal{G}$ is an $AE(X)$-cover, because X is at most n-dimensional. Hence, by the carrier theorem, there exists a map $\kappa\colon X \to Y$, carried by C. By the symmetry of assumptions, there exists a map $\lambda\colon Y \to X$, carried by C^{-1}. Let $\varPhi$ be a map from an at most $(n-1)$-dimensional simplicial complex into X. The map $\lambda \circ \kappa$ is carried by $C^{-1} \circ C$, hence it is $\mathcal{F}$-close to the identity on X. Therefore, by corollary 5.1, $\lambda \circ \kappa \circ \varPhi$ is homotopic to $\varPhi$, hence λ is an n-homotopy inverse of κ. Analogously, κ is an n-homotopy inverse of λ, so κ and λ are n-homotopy equivalences and X is n-homotopy equivalent to Y.

Let $\mathcal{B}_{\mathcal{F}}$ be the cover of the nerve $N(\mathcal{F})$ of $\mathcal{F}$ by barycentric stars of its vertices. Let $\mathcal{B}_{\mathcal{F}}^{(n)}$ denote the restriction of $\mathcal{B}_{\mathcal{F}}$ to the n-dimensional skeleton of the first barycentric subdivision of the nerve of $\mathcal{F}$. It is isomorphic to $\mathcal{B}_{\mathcal{F}}$, which is isomorphic to $\mathcal{F}$ by lemma 5.2. By lemma 5.3, $\mathcal{B}_{\mathcal{F}}$ is regular for the class of metric spaces. Hence, by the cellular approximation theorem (see [21, Corollary 4.12]), $\mathcal{B}_{\mathcal{F}}^{(n)}$ is n-regular. Therefore, by what we proved in the first paragraph, X is n-homotopy equivalent to the n-dimensional skeleton of the first barycentric subdivision of the nerve of $\mathcal{F}$. By the cellular approximation theorem, inclusions of n-dimensional skeleton of $N(\mathcal{F})$ into $N(\mathcal{F})$ and of n-dimensional skeleton of the first barycentric subdivision of $N(\mathcal{F})$ into $N(\mathcal{F})$ are weak n-homotopy equivalences. Hence, the inclusion of the former skeleton into the latter skeleton is a weak n-homotopy equivalence. By Whitehead's characterization, both skeletons are n-homotopy equivalent. We are done. ☐

Remark 6.3. If a closed n-regular cover of an at most n-dimensional space has an at most n-dimensional nerve, then by proposition 6.2 and by proposition 6.1, it admits both a canonical and an anticanonical map. By an argument used in the proof of the nerve theorem, these maps are n-homotopy equivalences and are n-homotopy inverses of each other.

Definition. We say that a closed cover is *n-semiregular* if it is a locally finite cover with at most n-dimensional nerve, which admits an anticanonical map that is an n-homotopy equivalence.

Lemma 6.4. *Every closed n-regular cover with at most n-dimensional nerve is n-semiregular.*

Proof. Apply proposition 6.1 and remark 6.3. □

Proposition 6.5. *Every $ANE(n)$-space admits an open cover $\mathcal{U}$ such that every anticanonical map of a closed locally finite locally finite dimensional cover that refines $\mathcal{U}$ is an n-homotopy equivalence. In particular, if a closed locally finite cover refines $\mathcal{U}$, has at most n-dimensional nerve and admits an anticanonical map, then it is n-semiregular.*

Proof. Let X be an $ANE(n)$-space. By theorem 2.4, there exists an open cover $\mathcal{U}$ of X such that every two $\mathcal{U}$-close maps into X are homotopic, provided that their domain is an at most $(n-1)$-dimensional metric space. We will show that $\mathcal{U}$ satisfies our claim. Let $\mathcal{F}$ ba a closed locally finite locally finite dimensional cover of X that refines $\mathcal{U}$ and has an anticanonical map $\lambda\colon N(\mathcal{F}) \to X$. By proposition 6.2, there exisits a canonical map $\kappa\colon X \to N(\mathcal{F})$ of $\mathcal{F}$. By lemma 5.3 and corollary 5.1, the composition $\kappa \circ \lambda$ is homotopic to the identity of $N(\mathcal{F})$. By the definitions, the composition $\lambda \circ \kappa$ is $\mathcal{F}$-close to the identity of X. Hence, for each map f from an at most $(n-1)$-dimensional simplicial complex into X, the composition $\lambda \circ \kappa \circ f$ is $\mathcal{U}$-close to f, hence homotopic to f, by the choice of $\mathcal{U}$. So, $\lambda \circ \kappa$ is n-homotopic to the identity on X. Hence, κ and λ are n-homotopy inverses of each other. □

Proposition 6.6. *If $\mathcal{F}$ is a closed n-semiregular cover of an at most n-dimensional $ANE(n)$-space X, then every map from an at most $(n-1)$-dimensional simplicial complex into an element of $\mathcal{F}$ is null-homotopic in X.*

Proof. Let f be a map from an at most $(n-1)$-dimensional simplicial complex into an element of $\mathcal{F}$. By proposition 6.2, there exists a canonical map of $\mathcal{F}$, which we denote by κ. Let λ be an anticanonical map of $\mathcal{F}$ that is an n-homotopy equivalence. By Whitehead's characterization, there exists a map $\mu\colon X \to N(\mathcal{F})$ such that $\lambda \circ \mu$ is n-homotopic to the identity on X. The composition $\kappa \circ \lambda$ is carried by $B \circ B^{-1}$, so it is $\mathcal{B}_{\mathcal{F}}$-close to the identity on the nerve of $\mathcal{F}$ and by corollary 5.1, it is homotopic to it. Hence, μ is homotopic to $\kappa \circ \lambda \circ \mu$, which is n-homotopic to κ. The identity on X is n-homotopic to $\lambda \circ \mu$, which is n-homotopic to $\lambda \circ \kappa$. Hence, by the definition of n-homotopy, f is homotopic to $(\lambda \circ \kappa) \circ f$. But $\kappa \circ f$ is null-homotopic because its image lies entirely in a barycentric star of a vertex of the nerve of $\mathcal{F}$, by the definition of κ and the assumption that the image of f lies entirely in an element of $\mathcal{F}$. Thus f, being homotopic to $\lambda \circ \kappa \circ f$, is null-homotopic. □

6.2 A construction of regular covers

We will now present a theorem (theorem 6.7) that allows us to construct an n-regular cover from an n-semiregular cover. We are still far away from its proof, which will be given in chapter 11.

Definition. We say that a cover $\mathcal{F}$ is *n-contractible in a cover* $\mathcal{G}$ if it refines $\mathcal{G}$ and if every map of an at most $(n-1)$-dimensional simplicial complex into an element of $\mathcal{F}$ is null-homotopic in an element of $\mathcal{G}$.

By a way of example, note that proposition 6.6 states that every closed n-semiregular cover of an at most n-dimensional $ANE(n)$-space X is n-contractible in $\{X\}$.

Theorem 6.7. *There exists a constant N such that if a closed star-finite k-regular n-semiregular ($k < n$) interior $\mathcal{N}_n$-cover $\mathcal{F}$ is n-contractible in a cover $\mathcal{E}$, then there exists a closed $(k+1)$-regular n-semiregular interior $\mathcal{N}_n$-cover isomorphic to $\mathcal{F}$ and refining $\operatorname{st}^N \mathcal{E}$. Moreover, we may require that the constructed cover is equal to $\mathcal{F}$ on a neighborhood of a given $Z(\mathcal{F})$-set.*

The rest of this section is devoted to proofs of three corollaries of theorem 6.7 that will be used in the proof of the main theorem.

Corollary 6.8. *If $\mathcal{F}$ is a closed star-finite n-semiregular interior $\mathcal{N}_n$-cover of a space X and Z is a $Z(\mathcal{F})$-set, then there exists a closed n-regular interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$ and is equal to $\mathcal{F}$ on a neighborhood of Z.*

Proof. By proposition 6.6, every closed n-semiregular cover of X is n-contractible in $\{X\}$. Let $\mathcal{F}_0 = \mathcal{F}$. By theorem 6.7 applied recursively with $\mathcal{E} = \{X\}$, for each $0 < k < n$ there exists a closed star-finite k-regular n-semiregular interior $\mathcal{N}_n$-cover $\mathcal{F}_k$ of X that is isomorphic to $\mathcal{F}_{k-1}$ and is equal to $\mathcal{F}_{k-1}$ on a neighborhood of Z. The cover $\mathcal{F}_n$ is a cover that satisfies our claim. ☐

Corollary 6.9. *For every cover $\mathcal{U}$ of an $\mathcal{N}_n$-space X there exists a cover $\mathcal{V}$ of X such that if $\mathcal{F}$ is a closed star-finite n-semiregular interior $\mathcal{N}_n$-cover that refines $\mathcal{V}$ and Z is a $Z(\mathcal{F})$-set, then there exists a closed n-regular interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$, is equal to $\mathcal{F}$ on an open neighborhood of Z and refines $\mathcal{U}$.*

Proof. Let N be the constant obtained via theorem 6.7. By theorem 2.4, for each open cover $\mathcal{U}_k$ of X there exists an open cover $\mathcal{V}_k$ of X that is n-contractible in $\mathcal{U}_k$. Define recursively a sequence $\mathcal{U}_{n-1}, \mathcal{V}_{n-1}, \ldots, \mathcal{U}_0, \mathcal{V}_0$ of open covers of X such that $\operatorname{st}^N \mathcal{U}_{n-1}$ refines $\mathcal{U}$, $\mathcal{V}_k$ is n-contractible in $\mathcal{U}_k$ and $\operatorname{st}^N \mathcal{U}_{k-1}$ refines $\mathcal{V}_k$. We will show that $\mathcal{V} = \mathcal{V}_0$ satisfies our claim. Let $\mathcal{F}_0 = \mathcal{F}$. By the assumption, $\mathcal{F}_0$ refines $\mathcal{V}_0$, hence it is n-contractible in $\mathcal{U}_0$. For $k = 1, 2, \ldots, n$, we apply theorem 6.7 to a cover $\mathcal{F}_{k-1}$ with $\mathcal{E} = \mathcal{U}_{k-1}$ and obtain a closed k-regular n-semiregular interior $\mathcal{N}_n$-cover $\mathcal{F}_k$ of X that is isomorphic to $\mathcal{F}_{k-1}$, refines $\operatorname{st}^N \mathcal{U}_{k-1}$ (and hence $\mathcal{V}_k$) and is equal to $\mathcal{F}_{k-1}$ on a neighborhood of Z. The cover $\mathcal{F}_n$ is a cover that satisfies our claim. ☐

Definition. We say that a cover is *m-barycentric* if its nerve has a structure of an mth barycentric subdivision of a locally finite simplicial complex K. We say that K is a *primary complex* of the cover.

Corollary 6.10. *For every n and k there is a constant $m_{n,k}$ such that if $\mathcal{E}$ is an n-regular $m_{n,k}$-barycentric cover of an $\mathcal{N}_n$-space X, $\mathcal{F}$ is a closed star-finite n-semiregular interior $\mathcal{N}_n$-cover that refines $\operatorname{st}^k \mathcal{E}$ and Z is a $Z(\mathcal{F})$-set, then there exists a closed n-regular interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$, refines $\operatorname{st}^{m_{n,k}} \mathcal{E}$ and is equal to $\mathcal{F}$ on an open neighborhood of Z.*

To prove corollary 6.10 we need four lemmas stated below. The notion of a join is auxiliary and won't be used later.

Definition. A kth *join* $\operatorname{jn}^k \mathcal{F}$ of an m-barycentric cover $\mathcal{F}$ with primary complex K is a cover by those (2^k-1)th stars of elements of $\mathcal{F}$ that correspond to vertices of the $(m-k)$th barycentric subdivision of K.

Remark 6.11. If a cover is m-barycentric, then its kth join is $(m-k)$-barycentric, when considered with the same primary complex. Also, every m-barycentric cover is k-barycentric for each $k < m$. For each $(k+l)$-barycentric cover $\mathcal{F}$, we have $\operatorname{jn}^k \operatorname{jn}^l \mathcal{F} = \operatorname{jn}^{k+l} \mathcal{F}$.

Lemma 6.12. *If $\mathcal{F}$ is m-barycentric, then $\operatorname{st}^{2^{m-1}-1} \mathcal{F} \prec \operatorname{jn}^m \mathcal{F} \subset \operatorname{st}^{2^m-1} \mathcal{F}$.*

Proof. The second relation follows directly from the definition. To prove the first relation, consider a discrete metric on the set $\mathcal{F}$, with the distance between two elements of $\mathcal{F}$ defined to be the length of a shortest chain containing them, decreased by 1 (a chain is a sequence of sets in which every two consecutive elements intersect). Then elements of $\operatorname{st}^{2^{m-1}-1} \mathcal{F}$ correspond to balls of radius $2^{m-1}-1$ in $\mathcal{F}$ and elements of $\operatorname{jn}^m \mathcal{F}$ correspond to balls of radius 2^m-1 in $\mathcal{F}$, but with centers in elements of $\mathcal{F}$ corresponding to vertices of the primary complex K of $\mathcal{F}$. The nerve of $\mathcal{F}$ is mth barycentric subdivision of K, hence a distance of an element of $\mathcal{F}$ from an element that corresponds to a vertex of K is not greater than 2^{m-1}. We are done. □

Definition. We say that a subcomplex L of a complex K is *full* if it contains every simplex spanned in K by its vertices.

Lemma 6.13. *If $L_1, L_2, \ldots, L_k$ are subcomplexes of some other complex, then*

$$\operatorname{bst}^m L_1 \cap \ldots \cap \operatorname{bst}^m L_k = \operatorname{bst}^{m-1}(\operatorname{bst} L_1 \cap \ldots \cap \operatorname{bst} L_k),$$

for each $m > 1$.

Proof. Assume L and M are subcomplexes of some other complex. We shall prove that

(1) If $L \cup M$ is full, then $\operatorname{bst} L \cap \operatorname{bst} M = \operatorname{bst}(L \cap M)$.

We have to show that $\operatorname{bst} L \cap \operatorname{bst} M \subset \operatorname{bst}(L \cap M)$. Assume that $L \cup M$ is a full subcomplex of a complex K. Observe that for each pair of vertices $v_1, v_2 \in K$ the interesction $\operatorname{bst} v_1 \cap \operatorname{bst} v_2$ is non-empty if and only if v_1, v_2 are connected by an edge. Let v be a vertex in $\operatorname{bst} L \cap \operatorname{bst} M$. Then $v \in \operatorname{bst} v_1$ and $v \in \operatorname{bst} v_2$ for some $v_1 \in L$ and $v_2 \in M$. Then v_1 and v_2 are connected by an edge, which must be contained either in L or in M, because $L \cup M$ is full. Hence either v_1 or v_2 is contained in $L \cap M$. Therefore v is contained in $\operatorname{bst}(L \cap M)$ and we are done.

A barycentric star of a subcomplex is always full, hence (1) yields the equality

$$\operatorname{bst}(\operatorname{bst}^{m-1} L_1) \cap \operatorname{bst}(\operatorname{bst}^{m-1} L_2 \cap \ldots \cap \operatorname{bst}^{m-1} L_k) = \operatorname{bst}(\operatorname{bst}^{m-1} L_1 \cap \ldots \operatorname{bst}^{m-1} L_k).$$

An elementary induction on m and k finishes the proof (observe that $\operatorname{bst}(\operatorname{bst}^{m-1} L) = \operatorname{bst}^{m-1}(\operatorname{bst} L)$). ☐

Lemma 6.14. *If L is a full subcomplex of a locally finite dimensional simplicial complex K, then* $\operatorname{bst} L$ *and L are homotopy equivalent.*

Proof. Let $\mathcal{L}$ denote the collection of barycentric stars of vertices of L, taken in K. Since L is a full subcomplex of K, $\mathcal{L}$ is isomorphic to the cover of L by barycentric stars of its vertices, taken in L. Hence, by lemma 5.2, the nerve of $\mathcal{L}$ is equal to L. By lemma 5.3 and by the definition of $\operatorname{bst} L$, $\mathcal{L}$ is a regular cover of $\operatorname{bst} L$. By theorem 5.5, $\operatorname{bst} L$ is homotopy equivalent to the nerve of $\mathcal{L}$. We are done. ☐

Lemma 6.15. *An mth join a closed n-regular m-barycentric cover of an at most n-dimensional metric space is n-regular.*

Proof. Let $\mathcal{F}$ be a closed n-regular m-barycentric cover of an at most n-dimensional metric space X. Let $\mathcal{B}_{\mathcal{F}}$ be a cover of the nerve of $\mathcal{F}$ by barycentric stars of its vertices. Let $\mathcal{B}_{\mathcal{F}}^{(n)}$ be the restriction of $\mathcal{B}_{\mathcal{F}}$ to the n-dimensional skeleton of the first barycentric subdivision of the nerve of $\mathcal{F}$. We have to show that non-empty intersections of elements of $\operatorname{jn}^m \mathcal{F}$ are absolute extensors in dimension n. Fix any such intersection A. Let B be the intersection of the corresponding elements of $\operatorname{jn}^m \mathcal{B}_{\mathcal{F}}^{(n)}$. As we argued in the proof of the nerve theorem, $\mathcal{B}_{\mathcal{F}}^{(n)}$ is n-regular and isomorphic to $\mathcal{F}$. Clearly, the restrictions of $\mathcal{F}$ to A and of $\mathcal{B}_{\mathcal{F}}^{(n)}$ to B are n-regular and isomorphic. Hence, by the nerve theorem, A is n-homotopy equivalent to B. By theorem 2.6, being an absolute extensor in dimension n is n-homotopy invariant in the class of absolute neighborhood extensors in dimension n. Hence it suffices to show that B is an absolute extensor in dimension n, which in turn would follow from the cellular approximation theorem (see [21, Corollary 4.12]) if the intersections of elements of $\operatorname{jn}^m \mathcal{B}_{\mathcal{F}}$ were contractible (cf. proof of lemma 5.3), which we are about to prove. Let K be a primary complex of $\mathcal{B}_{\mathcal{F}}$ such that the nerve of $\mathcal{B}_{\mathcal{F}}$ is the mth barycentric subdivision of K. Observe that $\operatorname{jn}^m \mathcal{B}_{\mathcal{F}}$ is a collection of mth barycentric stars of vertices of K. By lemma 6.13, if $\{v_i\}$

is a collection of vertices of K, then $\bigcap \operatorname{bst}^m v_i = \operatorname{bst}^{m-1}(\bigcap \operatorname{bst} v_i)$. By lemma 6.14, $\operatorname{bst}^{m-1}(\bigcap \operatorname{bst} v_i)$ is homotopy equivalent to $\bigcap \operatorname{bst} v_i$. The latter intersection is contractible by lemma 5.2. We are done. ☐

Proof of corollary 6.10. By the definitions, if a cover refines an n-regular cover, then it is n-contractible in it. By lemma 6.12 and lemma 6.15, the proof is reduced to an easy recursive application of theorem 6.7 and we omit its details. ☐

The following technical lemma will be used in chapter 8.

Lemma 6.16. *Assume that $\mathcal{G}$ is a 2-barycentric and n-regular cover of a space Y and $\mathcal{F}$ is a closed locally finite locally finite dimensional cover of an at most n-dimensional metric space X. If $\mathcal{F}$ and $\mathcal{G}$ are isomorphic, then every map f defined on a closed subset of X that maps elements of $\mathcal{F}$ into the corresponding elements of $\operatorname{st} \mathcal{G}$ extends over the entire domain to a map that maps elements of $\mathcal{F}$ into the corresponding elements of $\operatorname{st}^7 \mathcal{G}$.*

Proof. Let $\mathcal{F} = \{F_i\}_{i \in I}$ and $\mathcal{G} = \{G_i\}_{i \in I}$. By lemma 6.12 there exists a map q from the indexing set of $\operatorname{st} \mathcal{G}$ into the indexing set of $\operatorname{jn}^2 \mathcal{G}$ such that $\operatorname{st}_{\mathcal{G}} G_i \subset \operatorname{st}^3 G_{q(i)}$. The function $F_i \mapsto \operatorname{st}^3 G_{q(i)}$ is a carrier and f is carried by it. By lemma 6.15, $\operatorname{jn}^2 \mathcal{G}$ is n-regular, hence by the carrier theorem f extends over the entire domain to a map g such that $g(F_i) \subset \operatorname{st}^3 G_{q(i)}$. Since $G_i \subset \operatorname{st}^3 G_{q(i)}$, we have $G_{q(i)} \subset \operatorname{st}^4 G_i$. Hence $\operatorname{st}^3 G_{q(i)} = \operatorname{st}^3_{\mathcal{G}} G_{q(i)} \subset \operatorname{st}^3_{\mathcal{G}} \operatorname{st}^4 G_i = \operatorname{st}^7 G_i$. Hence g maps elements of $\mathcal{F}$ into the corresponding elements of $\operatorname{st}^7 \mathcal{G}$. ☐

6.3 A construction of semiregular covers

First we briefly describe the technique of the construction (cf. figure 6.1). Consider an $\mathcal{N}_n$-space X. By lemma 2.26, there exists a weak n-homotopy equivalence of an at most n-dimensional countable locally finite simplicial complex L and X. By the strong univerality of X, we may approximate it by a closed embedding. By lemma 2.21, we may require (and we do) that the approximation is also a weak n-homotopy equivalence. Let K denote its image. It is a locally finite simplicial complex embedded as a closed subset of X and the inclusion $K \subset X$ is a weak n-homotopy equivalence. By theorem 2.23, there exists a retraction $r\colon X \to K$. Let $\mathcal{B}_K$ be the cover of K by barycentric stars of its vertices. The nerve of $\mathcal{B}_K$ is isomorphic to K. Let $\mathcal{F} = r^{-1}(\mathcal{B}_K)$. It is isomorphic to $\mathcal{B}_K$, hence its nerve is isomorphic to K. The identity on K is an anticanonical map of $\mathcal{F}$. It is a weak n-homotopy equivalence by the construction. By Whitehead's characterization, it is an n-homotopy equivalence, hence $\mathcal{F}$ is n-semiregular.

The construction described above allows us to construct a closed n-semiregular cover of an $\mathcal{N}_n$-space, but it doesn't give us a control over the size of the constructed



Figure 6.1: Construction of a semiregular cover.

cover. It is because theorem 2.23 does not give a way to limit the size of fibers of r. Theorem 6.17 allows us to do just that.

Theorem 6.17. *For every open cover $\mathcal{U}$ of an $\mathcal{N}_n$-space X there exists a locally finite simplicial complex K embedded as a closed subset of X and a retraction of X onto K that is $\mathcal{U}$-close to the identity.*

The proof of theorem 6.17 will be given in section 10.4.

Theorem 6.18. *For each integer m, each $\mathcal{N}_n$-space X and each Z-set Z in X there exist arbitrarily small closed star-finite m-barycentric n-semiregular interior $\mathcal{N}_n$-covers $\mathcal{F}$ of X such that Z is a $Z(\mathcal{F})$-set.*

Proof. Let $\mathcal{U}$ be an open cover of X. By theorem 6.17, there exists a locally finite simplicial complex K embedded as a closed subset of X and a retraction $r\colon X \to K$ that is $\mathcal{U}$-close to the identity. We may assume that the set of simplices of K refines $\mathcal{U}$ and that K has a structure of an mth barycentric subdivision of another complex. Let $\mathcal{F} = r^{-1}(\mathcal{B}_K)$ and observe that $\mathcal{F}$ is a closed locally finite star-finite m-barycentric cover that refines $\operatorname{st}\mathcal{U}$. By theorem 4.5, there is a swelling $\mathcal{G}$ of $\mathcal{F}$ to a closed interior $\mathcal{N}_n$-cover such that Z is a $Z(\mathcal{G})$-set and $\mathcal{G} \prec \operatorname{st}\mathcal{U}$. As K is the nerve of $\mathcal{G}$, the inclusion of K into X is an anticanonical map of $\mathcal{G}$. By proposition 6.5, if $\mathcal{U}$ is small enough, then $\mathcal{G}$ is n-semiregular. We are done. ☐

Chapter 7

Extending homeomorphisms by the use of a "brick partitionings" technique

The classical approach to the problem of constructing a homeomorphism between complete spaces is the brick partitionings technique. Theorem 7.1 enhances this technique and sets the strategy for the proof of the main theorem.

Definition. We write $\mathcal{F}_1 \prec_f \mathcal{F}_2$ if $\mathcal{F}_1$ and $\mathcal{F}_2$ are covers of the same space and f is a map from the indexing set of $\mathcal{F}_1$ into the indexing set of $\mathcal{F}_2$ such that for all i the element of $\mathcal{F}_1$ indexed by i is a subset of the element of $\mathcal{F}_2$ indexed by $f(i)$.

Theorem 7.1 (The homeomorphism extension theorem)**.** *Assume that $\mathcal{F}_k$ and $\mathcal{G}_k$ are sequences of locally finite closed covers of complete spaces X and Y. We do not assume that X or Y is separable or that $\mathcal{F}_k$ or $\mathcal{G}_k$ is countable. Assume that $\mathcal{F}_k$ and $\mathcal{G}_k$ are isomorphic for all k and f_k is a sequence of maps such that for some m and each k*

$$\begin{array}{rcccl} \mathcal{F}_{2k+1} & \prec_{f_{2k+1}} & \operatorname{st}^m \mathcal{F}_{2k} & \prec_{f_{2k}} & \mathcal{F}_{2k-1}, \\ \operatorname{st}^m \mathcal{G}_{2k+1} & \prec_{f_{2k+1}} & \mathcal{G}_{2k} & \prec_{f_{2k}} & \operatorname{st}^m \mathcal{G}_{2k-1}. \end{array}$$

If meshes of $\mathcal{F}_k$ and of $\mathcal{G}_k$ converge to zero, then there exists a unique homeomorphism from X onto Y that maps each element of $\mathcal{F}_k$ into the corresponding element of $\operatorname{st}^{m+1} \mathcal{G}_k$.

The name of *the homeomorphism extension theorem* deserves an explanation as the theorem does not mention explicitly extensions of homeomorphisms. It is justified by the fact that the uniqueness of the homeomorphism implies that it extends every partial homeomorphism that maps elements of $\mathcal{F}_k$'s into the corresponding elements of $\operatorname{st}^{m+1} \mathcal{G}_k$'s.

Proof. The uniqueness of a homeomorphism from X onto Y that maps elements of $\mathcal{F}_k$ into the corresponding elements of $\operatorname{st}^{m+1}\mathcal{G}_k$ is a direct consequence of the assumption that meshes of $\mathcal{G}_k$ converge to zero. Every two maps that satisfy this condition must be $\operatorname{st}^{m+1}\mathcal{G}_k$-close for all k.

The existence is proved in two steps. Let $\mathcal{F}_k = \{F_i^k\}_{i\in I_k}$ and $\mathcal{G}_k = \{G_i^k\}_{i\in I_k}$. In the first step we prove that there exists a function $f\colon X \to Y$ such that $f(F_i^k) \subset \operatorname{st}^{m+1} G_i^k$ for all $k \in \mathbb{N}$ and $i \in I_k$. In the second step we prove that it is a homeomorphism.

By the definition, an mth star of an element of a collection of sets is a union of some of the elements of the collection. We shall call summands of this union *components* of the mth star. Note that the set of components of an mth star may differ from the set of elements of the collection that are subsets of the mth star. If $\mathcal{C}$ is a collection of sets and if C_1, C_2 are elements of $\mathcal{C}$, then C_1 is a component of $\operatorname{st}^m_{\mathcal{C}} C_2$ if and only if C_2 is a component of $\operatorname{st}^m_{\mathcal{C}} C_1$.

For each $x \in X$ let $A_k(x) = \bigcup\{\operatorname{st}^m G_i^k \colon i \in I_k, x \in F_i^k\}$. We first wish to apply the Cantor theorem to prove that $\bigcap_k A_k(x) \neq \emptyset$ for each $x \in X$. By local finiteness of $\mathcal{G}_k$'s, each $A_k(x)$ is closed. The sequence of diameters of $A_k(x)$ converges to zero as k goes to infinity since $A_k(x)$ is an union of mth stars of intersecting elements of $\mathcal{G}_k$. The proof of the inclusion $A_{k+1}(x) \subset A_k(x)$ depends on the parity of k. From the definition of $A_{k+1}(x)$ it suffices to check that for each $i \in I_{k+1}$ such that $x \in F_i^{k+1}$ the inclusion $\operatorname{st}^m G_i^{k+1} \subset A_k(x)$ holds.

Assume that k is even. Then $F_i^{k+1} \subset \operatorname{st}^m F^k_{f_{k+1}(i)}$. Then for some $j \in I_k$ such that F_j^k is a component of $\operatorname{st}^m F^k_{f_{k+1}(i)}$, we have $x \in F_j^k$. Then from the definition $\operatorname{st}^m G_j^k \subset A_k(x)$. But $\mathcal{F}_k$ and $\mathcal{G}_k$ are isomorphic, so $G^k_{f_{k+1}(i)}$ is a component of $\operatorname{st}^m G_j^k$. In particular, $G^k_{f_{k+1}(i)} \subset \operatorname{st}^m G_j^k$. Hence $\operatorname{st}^m G_i^{k+1} \subset G^k_{f_{k+1}(i)} \subset \operatorname{st}^m G_j^k \subset A_k(x)$, the first inclusion from the assumption that k is even.

Assume that k is odd. Let $j \in I_{k+1}$ such that F_j^{k+1} is a component of $\operatorname{st}^m F_i^{k+1}$. Then F_i^{k+1} is a component of $\operatorname{st}^m F_j^{k+1}$. In particular, $F_i^{k+1} \subset \operatorname{st}^m F_j^{k+1}$. Hence $x \in F_i^{k+1} \subset \operatorname{st}^m F_j^{k+1} \subset F^k_{f_{k+1}(j)}$, the last inclusion from the assumption that k is odd. By the definition, $\operatorname{st}^m G^k_{f_{k+1}(j)} \subset A_k(x)$ and by the assumptions $G_j^{k+1} \subset \operatorname{st}^m G^k_{f_{k+1}(j)}$. Therefore $G_j^{k+1} \subset A_k(x)$ so $\operatorname{st}^m G_i^{k+1} \subset A_k(x)$.

Thus Cantor's theorem applies to give a unique function $x \mapsto f(x) \in \bigcap_k A_k(x)$ such that if $x \in F_i^k$ then $f(x) \in A_k(x) \subset \operatorname{st}^{m+1} G_i^k$ for all $k \in \mathbb{N}$ and $i \in I_k$. Let $x \in X$ and an open neighborhood U of $f(x)$ in Y be given. Take k so large that the star of $f(x)$ in $\operatorname{st}^{m+1}\mathcal{G}_k$ lies in U. By local finiteness of $\mathcal{F}_k$, there is open neighborhood V of x in X that lies in a star of x in a cover $\mathcal{F}_k$. Then $f(V) \subset U$, showing that f is continuous. By the symmetry of assumptions (which we obtain after throwing away $\mathcal{F}_1$ and $\mathcal{G}_1$ from the sequences), there exists a continuous map $g\colon Y \to X$ such that $g(G_i^k) \subset \operatorname{st}^{m+1} F_i^k$. By these conditions, $g(f(F_i^k)) \subset \operatorname{st}^{2m+2} F_i^k$ so $g \circ f = id_X$ because it is $\operatorname{st}^{2m+2}\mathcal{F}_k$-close to id_X for all k. Similarly $f \circ g = id_Y$, so f is a homeomorphism from X onto Y. □

Chapter 8

Proof of the main results

In this chapter we prove the main theorem and derive all theorems stated in the introduction. We use theorems 6.7 and 6.17, whose proofs will be given in the third part of the paper.

Definition. We say that a cover is a *closed partition of an* $\mathcal{N}_n$*-space* X if it is a closed star-finite n-regular interior $\mathcal{N}_n$-cover of X with at most n-dimensional nerve.

Lemma 8.1. *For each open cover* $\mathcal{U}$ *of an* $\mathcal{N}_n$*-space* X *there exists a closed partition* $\mathcal{F}$ *of* X *that refines* $\mathcal{U}$*. Moreover, if* m *is an integer and* Z *is a* Z*-set in* X*, then we may require that* $\mathcal{F}$ *is* m*-barycentric and* Z *is a* $Z(\mathcal{F})$*-set.*

Proof. Let $\mathcal{V}$ be an open cover of X obtained via corollary 6.9 applied to $\mathcal{U}$. By theorem 6.18 there exists a closed star-finite m-barycentric n-semiregular interior $\mathcal{N}_n$-cover $\mathcal{E}$ that refines $\mathcal{V}$ and such that Z is a $Z(\mathcal{E})$-set. Let $\mathcal{F}$ be a closed n-regular interior $\mathcal{N}_n$-cover of X that refines $\mathcal{U}$ and that is isomorphic to $\mathcal{E}$ and equal to $\mathcal{E}$ on an open neighborhood of Z. The existence of $\mathcal{F}$ follows again from corollary 6.9, this time applied to $\mathcal{E}$, Z, $\mathcal{V}$ and $\mathcal{U}$. By the construction, $\mathcal{F}$ is m-barycentric and Z is a $Z(\mathcal{F})$-set. We are done. $\square$

Definition. If $\mathcal{F}$ and $\mathcal{G}$ are isomorphic covers of spaces X and Y respectively and h is a homeomorphism from a subset of X onto a subset of Y, then we say that $\mathcal{F}$ *and* $\mathcal{G}$ *are compatible with* h if h maps elements of $\mathcal{F}$ into the corresponding elements of $\operatorname{st}\mathcal{G}$ and h^{-1} maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}\mathcal{F}$.

Lemma 8.2. *If* $\mathcal{F}$ *is a closed partition of an* $\mathcal{N}_n$*-space* X *and* Y *is an* $\mathcal{N}_n$*-space* n*-homotopic to* X*, then there exists a closed parition* $\mathcal{G}$ *of* Y *that is isomorphic to* $\mathcal{F}$*. Moreover, if an* n*-homotopy equivalence* $f\colon X \to Y$ *is a* Z*-embedding on a closed subset* A *of* X*, then we may require that* $\mathcal{F}$ *and* $\mathcal{G}$ *are compatible with* $f_{|A}$ *and that* $f(A)$ *is a* $Z(\mathcal{G})$*-set.*

Proof. By theorem 2.4, if a map into an $ANE(n)$-space is close enough to an n-homotopy equivalence, then it is an n-homotopy equivalence. Hence, by theorem 2.35, we may approximate f rel A by a closed embedding h that is an n-homotopy equivalence. By theorem 2.23 there exists a retraction $r\colon Y \to h(X)$. Let $\mathcal{F} = \{F_i\}_{i\in I}$ and let $\mathcal{R} = \{r^{-1}(h(F_i))\}_{i\in I}$. The collection $\mathcal{R}$ is a closed locally finite cover of Y. It is isomorphic to $\mathcal{F}$, hence it is star-finite and has at most n-dimensional nerve. Covers $\mathcal{F}$ and $\mathcal{R}$ are compatible with $f_{|A}$ by the definition of r and h. Let λ be an anticanonical map of $\mathcal{F}$ and an n-homotopy equivalence of $N(\mathcal{F})$ and X. The map $f \circ \lambda$ is an anticanonical map of $\mathcal{R}$ and an n-homotopy equivalence of $N(\mathcal{R})$ and Y. Hence $\mathcal{R}$ is n-semiregular. By theorem 4.5 there is a closed interior $\mathcal{N}_n$-cover $\mathcal{H}$ that is a swelling of $\mathcal{R}$ and such that $h(A)$ is a $Z(\mathcal{H})$-set. We may assume that the swelling is so small that $\mathcal{F}$ and $\mathcal{H}$ are compatible with $f_{|A}$. The cover $\mathcal{H}$ is isomorphic to $\mathcal{R}$, hence it is star-finite and has at most n-dimensional nerve. As a swelling of an n-semiregular cover, $\mathcal{H}$ admits an anticanonical map that is an n-homotopy equivalence. Hence $\mathcal{H}$ is n-semiregular. We apply corollary 6.8 to $\mathcal{H}$ and to the $Z(\mathcal{H})$-set $h(A)$ to obtain a closed n-regular interior $\mathcal{N}_n$-cover $\mathcal{G}$ isomorphic to $\mathcal{H}$ and equal to $\mathcal{H}$ on an neighborhood of $h(A)$. This cover satisfies all prescribed conditions. ☐

Lemma 8.3. *For each n there exists an integer m such that if $\mathcal{F}$ and $\mathcal{G}$ are isomorphic closed m-barycentric partitions of $\mathcal{N}_n$-spaces X and Y respectively and $\mathcal{P}$ is a closed partition of X such that $\mathcal{P} \prec_p \mathcal{F}$ (where p is a map from the indexing set of $\mathcal{P}$ into the indexing set of $\mathcal{F}$), then there exists a closed partition $\mathcal{Q}$ of Y that is isomorphic to $\mathcal{F}$ and such that $\mathcal{Q} \prec_p \operatorname{st}^m \mathcal{G}$. Moreover, if h is a $Z(\mathcal{G})$-embedding of a closed subset A of X into Y such that $\mathcal{F}$ and $\mathcal{G}$ are compatible with h, then we may require that $\mathcal{P}$ and $\mathcal{Q}$ are compatible with h as well.*

Proof. We let m to be equal to $m_{n,9}+1$ or 2, whichever is greater, with $m_{n,9}$ obtained via corollary 6.10. Let $\mathcal{F} = \{F_i\}_{i\in I}$ and $\mathcal{G} = \{G_i\}_{i\in I}$.

By the carrier theorem, there is a map $f\colon X \to Y$ that maps elements of $\mathcal{F}$ into the corresponding elements of $\mathcal{G}$. The composition of f with a canonical map of $\mathcal{G}$ is a canonical map of $\mathcal{F}$. By the nerve theorem, canonical maps of $\mathcal{F}$ and of $\mathcal{G}$ are n-homotopy equivalences. Hence f is an n-homotopy equivalence as well. By lemma 2.21 and by Whitehead's characterization, every approximation of f that is sufficiently close is an n-homotopy equivalence. By the definition, every approximation of f within $\mathcal{G}$ maps elements of $\mathcal{F}$ into the corresponding elements of $\mathcal{G}$. Hence by corollary 3.7 we may assume that f is a closed embedding that has image disjoint from $h(A)$. The function $h^{-1} \cup f^{-1}$ is well defined and continuous and maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}\mathcal{F}$. By lemma 6.16, $h^{-1} \cup f^{-1}$ extends over Y to a map g that maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}^7 \mathcal{F}$. Note that $g^{-1}(F_i) \subset \operatorname{st}^8 G_i$.

Let $\mathcal{P} = \{P_j\}_{j\in J}$. Let $\mathcal{R} = \{g^{-1}(P_j)\}_{j\in J}$. We have $\mathcal{R} \prec_p \operatorname{st}^8 \mathcal{G}$. The collection $\mathcal{R}$ is a closed locally finite cover of Y. It is isomorphic to $\mathcal{P}$, hence it is star-finite and has at most n-dimensional nerve. Covers $\mathcal{P}$ and $\mathcal{R}$ are compatible with h by the definition of g. Let λ be an anticanonical map of $\mathcal{P}$ and an n-homotopy equivalence of $N(\mathcal{P})$ and X. The map $f \circ \lambda$ is an anticanonical map of $\mathcal{R}$ and an n-homotopy equivalence of $N(\mathcal{R})$ and Y. Hence $\mathcal{R}$ is n-semiregular. By 4.5 there is a closed, locally finite interior $\mathcal{N}_n$-cover $\mathcal{S}$ that is a swelling of the cover $\mathcal{R}$ such that $h(A)$ is a $Z(\mathcal{S})$-set. If the swelling is small enough, then $\mathcal{S} \prec \operatorname{st}^9 \mathcal{G}$ and covers $\mathcal{P}$ and $\mathcal{S}$ are compatible with h. Let Z be a $Z(\mathcal{S})$-set that contains $h(A)$ and meets every non-empty intersection of elements of $\mathcal{S}$ (we may let Z to be the union of $h(A)$ and $\operatorname{im} f \circ \lambda$, by remark 3.5 and by proposition 3.3 this set is a $Z(\mathcal{S})$-set). We apply corollary 6.10 to $\mathcal{S}$, $Z(\mathcal{S})$-set Z and $\mathcal{E} = \mathcal{G}$ to obtain a closed n-regular $\mathcal{N}_n$-cover $\mathcal{Q}$ that refines $\operatorname{st}^{m_{n,9}}$, is isomorphic to $\mathcal{S}$ and is equal to $\mathcal{S}$ on an neighborhood of Z. We have $\mathcal{Q} \prec \operatorname{st}^{m_{n,9}} \mathcal{G}$. By the assumption that Z meets every non-empty intersection of elements of $\mathcal{S}$, we have $\mathcal{Q} \prec_p \operatorname{st}^{m_{n,9}+1} \mathcal{G}$. The cover $\mathcal{Q}$ is a closed partition of Y. Since $\mathcal{Q}$ is equal to $\mathcal{S}$ on $h(A)$, partitions $\mathcal{P}$ and $\mathcal{Q}$ are compatible with h. We are done. ☐

Lemma 8.4. *For each n there exists a constant m such that if $\mathcal{F}$ is an m-barycentric closed partition of an n-dimensional Nöbeling manifold X and $\mathcal{G}$ is an isomorphic closed partition of an n-dimensional Nöbeling manifold Y, then there exists a homeomorphism from X onto Y that maps elements of $\mathcal{F}$ into the corresponding elements of $\operatorname{st}^m \mathcal{G}$. Moreover, if $\mathcal{F}$ and $\mathcal{G}$ are compatible with a homeomorphism h of a Z-set in X onto a $Z(\mathcal{G})$-set in Y, then we may require that the ambient homeomorphism extends h.*

Proof. Let m be a constant obtained via lemma 8.3 applied to n. We will construct a sequence of covers and functions $\mathcal{G}_1, \mathcal{F}_1, \mathcal{F}_2, f_2, \mathcal{G}_2, \mathcal{G}_3, f_3, \mathcal{F}_3, \dots$ (in this particular order), where f_k is a map from the indexing set of $\mathcal{F}_k$ (and of $\mathcal{G}_k$) into the indexing set of $\mathcal{F}_{k-1}$ (and of $\mathcal{G}_{k-1}$), such that the following conditions are satisfied for each k.

(1) $\mathcal{F}_k$ and $\mathcal{G}_k$ are isomorphic closed m-barycentric partitions of X and Y respectively.

(2) Z_1 is a $Z(\mathcal{F}_{2k})$-set and Z_2 is a $Z(\mathcal{G}_{2k-1})$-set.

(3) $\operatorname{mesh} \mathcal{F}_{2k} < \frac{1}{k}$ and $\operatorname{mesh} \mathcal{G}_{2k+1} < \frac{1}{k+1}$.

(4) $$\begin{aligned} \operatorname{st}^m \mathcal{F}_{2k} &\prec_{f_{2k}} \mathcal{F}_{2k-1}, & \mathcal{F}_{2k+1} &\prec_{f_{2k+1}} \operatorname{st}^m \mathcal{F}_{2k} \\ \mathcal{G}_{2k} &\prec_{f_{2k}} \operatorname{st}^m \mathcal{G}_{2k-1}, & \operatorname{st}^m \mathcal{G}_{2k+1} &\prec_{f_{2k+1}} \mathcal{G}_{2k}. \end{aligned}$$

(5) $\mathcal{F}_k$ and $\mathcal{G}_k$ are compatible with h.

By corollary 2.36, there exists a closed embedding $h_1 \colon X \to Y$ and a closed embedding $h_2 \colon Y \to X$ such that $h_{1|Z_1} = h$, $h_{2|Z_2} = h^{-1}$ and both h_1 and h_2 are n-homotopy equivalences.

We let $\mathcal{G}_1 = \mathcal{G}$ and $\mathcal{F}_1 = \mathcal{F}$. Let $k \geq 1$ and assume that we already constructed $\mathcal{G}_{2k-1}$ and $\mathcal{F}_{2k-1}$. Let $\mathcal{U}$ be an open cover of X with mesh less than $1/k$ and such that $\operatorname{st}^m \mathcal{U}$ refines $\mathcal{F}_{2k-1}$. By lemma 8.1, there exists a closed m-barycentric partition $\mathcal{F}_{2k}$ of X that refines $\mathcal{U}$ and such that Z_1 is a $Z(\mathcal{F}_{2k})$-set. By the choice of $\mathcal{U}$, the mesh of $\mathcal{F}_{2k}$ is less than $1/k$ and there exists a function f_{2k} such that $\mathcal{F}_{2k} \prec_{f_{2k}} \mathcal{F}_{2k-1}$. We let f_{2k} to be any such function. By the choice of m and by lemma 8.3 applied with $\mathcal{F} = \mathcal{F}_{2k-1}$, $\mathcal{G} = \mathcal{G}_{2k-1}$, $\mathcal{P} = \mathcal{F}_{2k}$ and $p = f_{2k}$, there exists a closed partition $\mathcal{G}_{2k}$ of Y that is isomorphic to $\mathcal{F}_{2k}$, satisfies the identity $\operatorname{st}^m \mathcal{G}_{2k} \prec_{f_{2k}} \mathcal{G}_{2k-1}$ and such that $\mathcal{F}_{2k}$ and $\mathcal{G}_{2k}$ are compatible with h.

The construction of $\mathcal{G}_{2k+1}$ and f_{2k+1} is similar to the construction of $\mathcal{F}_{2k}$ and f_{2k} and the construction of $\mathcal{F}_{2k+1}$ is similar to the construction of $\mathcal{G}_{2k}$.

We constructed a sequence $\mathcal{G}_1, \mathcal{F}_1, \mathcal{F}_2, f_2, \ldots$ that satisfies conditions (1) - (5) stated above. Hence by theorem 7.1, there exists a homeomorphism from X onto Y that maps elements of $\mathcal{F}_k$ into the corresponding elements of $\operatorname{st}^{m+1} \mathcal{G}_k$. By (5), the restriction of this homeomorphism to Z_1 is $\operatorname{st}^{m+1} \mathcal{G}_k$-close to h for each k. By (3), meshes of $\operatorname{st}^{m+1} \mathcal{G}_k$ converge to 0, so this restriction must be equal to h, hence the constructed homeomorphism extends h. We are done. □

Now we are set to prove the main theorem. Let us recall the statement.

Main theorem. *A homeomorphism $h\colon Z_1 \to Z_2$ of a Z-set Z_1 in an $\mathcal{N}_n$-space X onto a Z-set Z_2 in an $\mathcal{N}_n$-space Y extends to an ambient homeomorphism if and only if it extends to an n-homotopy equivalence of X and Y.*

Proof. Only the “if” part requires a proof. Let m be a constant obtained via lemma 8.4 applied to n. Let $h_2\colon X \to Y$ be an extension of h^{-1} to an n-homotopy equivalence. It exists by corollary 2.36. By lemma 8.1, there exists a closed m-barycentric partition $\mathcal{G}$ of Y such that Z_2 is a $Z(\mathcal{G})$-set. By lemma 8.2 applied to $\mathcal{G}$ and an n-homotopy equivalence h_2, there exists a closed partition $\mathcal{F}$ of X that is isomorphic to $\mathcal{G}$ and such that $Z_1 = h_2(Z_2)$ is a $Z(\mathcal{F})$-set and $\mathcal{F}$ and $\mathcal{G}$ are compatible with h. An application of lemma 8.4 finishes the proof. □

The proof of the local Z-set unknotting is similar. Let us recall the statement.

Local Z-set unknotting theorem. *For every open cover $\mathcal{U}$ of a Nöbeling manifold X there exists an open cover $\mathcal{V}$ such that every homeomorphism $h\colon Z_1 \to Z_2$ between Z-subsets of the manifold that is $\mathcal{V}$-close to the inclusion $Z_1 \subset X$ extends to a homeomorphism of the entire manifold that is $\mathcal{U}$-close to the identity.*

Proof. Let $\mathcal{U}$ be an open cover of an n-dimensional Nöbeling manifold X. Let m be a constant obtained via lemma 8.4 applied to n. Let $\mathcal{W}$ be an open cover of X whose $(2m+1)$-th star refines $\mathcal{U}$. By lemma 8.1, there exists a closed m-barycentric partition $\mathcal{G}$ of X that refines $\mathcal{W}$. We let $\mathcal{V}$ be an open cover whose star refines $\mathcal{G}$. We will show that it satisfies the assertion of the theorem.

Let $h\colon Z_1 \to Z_2$ be a homeomorphism between Z-subsets of X that is $\mathcal{V}$-close to the inclusion of Z_1 into X. Observe that neither Z_1 nor Z_2 has to be a $Z(\mathcal{G})$-set. By theorem 3.6, there exists a closed $Z(\mathcal{G})$-embedding $i\colon Z_1 \to X$ that is $\mathcal{V}$-close to the inclusion of Z_1 into X. Let $\mathcal{F} = \mathcal{G}$. Since $\operatorname{st} \mathcal{V}$ refines $\mathcal{G}$, both i and $i \circ h^{-1}$

map elements of $\mathcal{G}$ into their stars. Hence partitions $\mathcal{F}$ and $\mathcal{G}$ are compatible with i and with $i \circ h^{-1}$. By lemma 8.4, there exists a homeomorphism h_1 of X onto X that extends i and maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}^m \mathcal{G}$ and a homeomorphism h_2 of X onto X that extends $i \circ h^{-1}$ and maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}^m \mathcal{G}$. Observe that h_2^{-1} maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}^{m+1} \mathcal{G}$, hence $h_2^{-1} \circ h_1$ maps elements of $\mathcal{G}$ into the corresponding elements of $\operatorname{st}^{2m+1} \mathcal{G}$. By the construction, $h_2^{-1} \circ h_1$ extends h and is $\mathcal{U}$-close to the identity. We are done. ☐

Z-Set unknotting theorem. *A homeomorphism between Z-sets in n-dimensional Nöbeling manifolds extends to an ambient homeomorphism if and only if it extends to an n-homotopy equivalence.*

Proof. By theorem 2.15, the n-dimensional Nöbeling space is strongly universal in dimension n. By corollary 2.28, it is absolute extensor in dimension n. Therefore theorems 2.4 and 2.16 imply that Nöbeling manifolds are $\mathcal{N}_n$-spaces. Therefore the Z-set unknotting theorem is just a special case of the main theorem, which we just proved. ☐

Open embedding theorem. *Every n-dimensional Nöbeling manifold is homeomorphic to an open subset of the n-dimensional Nöbeling space.*

Proof. By proposition 2.29, every $\mathcal{N}_n$-space is n-homotopy equivalent to an open subset of ν^n. Hence, by the main theorem, it is homeomorphic to an open subset of ν^n. ☐

Characterization theorem. *An n-dimensional Polish space is a Nöbeling manifold if and only if it is an $ANE(n)$-space and is strongly universal in dimension n.*

Proof. By the definition, every open subset of ν^n is an n-dimensional Nöbeling manifold. See the proof of the open embedding theorem. ☐

Sum theorem. *If a space X is an union of two closed n-dimensional Nöbeling manifolds whose intersection is also an n-dimensional Nöbeling manifold, then X is an n-dimensional Nöbeling manifold.*

Proof. By the characterization theorem, every n-dimensional Nöbeling manifold is an $\mathcal{N}_n$-space. By corollary 3.8, an union of two closed $\mathcal{N}_n$-spaces that meet in an $\mathcal{N}_n$-space, is an $\mathcal{N}_n$-space. By the characterization theorem, every $\mathcal{N}_n$-space is an n-dimensional Nöbeling manifold. We are done. ☐

Part III

Constructing n-semiregular and n-regular $\mathcal{N}_n$-covers

Chapter 9

Basic constructions in $\mathcal{N}_n$-spaces

9.1 Adjustment to a Z-collection

See page 30 for the definition of a Z-collection.

Lemma 9.1. *Every star-finite locally finite closed $\mathcal{N}_n$-cover $\mathcal{F}$ admits arbitrarily small adjustment to a Z-collection. Moreover, given arbitrary $Z(\mathcal{F})$-set and a closed shrinking of of $\mathcal{F}$, we may assume that the Z-collection is equal to the shrinking on the $Z(\mathcal{F})$-set.*

Proof. Let $\mathcal{F} = \{F_i\}_{i \in I}$ denote a star-finite locally finite closed $\mathcal{N}_n$-cover of a space X. Let $\mathcal{I} = \{J \subset I \colon J \neq \emptyset, F_J \neq \emptyset\}$. Order $\mathcal{I}$ into a sequence $J_1, J_2, \ldots$ non increasing in the order by inclusion. Such ordering exists because $\mathcal{F}$ is star-finite. Let Z be a $Z(\mathcal{F})$-set and let $\mathcal{G} = \{G_i\}_{i \in I}$ be a closed shrinking of $\mathcal{F}$. Let $\mathcal{U}$ be an open cover of X. We construct a sequence of maps $\{\varphi_k \colon F_{J_k} \to X\}_{k \in \mathbb{N}}$ such that for each k the following conditions are satisfied.

- (1_k) φ_k is a $Z(\mathcal{F})$-embedding that is $\mathcal{U}$-close to the inclusion of F_{J_k} into X,
- (2_k) $\operatorname{im} \varphi_k \subset F_{J_k}$ and $\operatorname{im} \varphi_k \cap Z = G_{J_k} \cap Z$,
- (3_k) if $J_k \cup J_l = J_m$ and $l < k$, then $\operatorname{im} \varphi_k \cap \operatorname{im} \varphi_l = \operatorname{im} \varphi_m$,
- (4_k) if $J_k \subsetneq J_l$, then $\varphi_k \circ \varphi_l = \varphi_l$.

Fix $k \in \mathbb{N}$ and assume that for each $l < k$ we constructed map φ_l that satisfies conditions (1_l), (2_l) and (3_l). Let $Z_1 = F_{J_k} \cap (Z \cup \bigcup_{l<k} \operatorname{im} \varphi_l)$. By (1), φ_l is a $Z(\mathcal{F})$-set for each $l < k$, so Z_1 is a $Z(\mathcal{F})$-set. Let $Z_2 = (G_{J_k} \cap Z) \cup \bigcup_{l<k \colon J_l \supset J_k} \operatorname{im} \varphi_l$ and observe that $Z_2 \subset Z_1$ and $Z_2 \cap Z = G_{J_k} \cap Z$. Let $A = F_{J_k} \sqcup_{Z_2} Z_1$ denote the disjoint sum of F_{J_k} and Z_1 glued along Z_2. We treat F_{J_k} and Z_2 as subspaces of A. Let $f \colon A \to X$ denote the map induced on A by identities on F_{J_k} and Z_1. By the definition, f restricted to Z_1 is a $Z(\mathcal{F})$-embedding. By theorem 3.6, there is a $\mathcal{U}$-approximation g of f, $\operatorname{rel} Z_1$ and within $\mathcal{F}$, by a $Z(\mathcal{F})$-embedding. We let φ_k to be

the restriction of g to F_{J_k}. By the definition of g, φ_k is a $Z(\mathcal{F})$-embedding that is $\mathcal{U}$-close to the inclusion of F_{J_k} into X and its image lies in F_{J_k}. By the definition of g, $\operatorname{im}\varphi_k \cap Z_1 = Z_2$, hence $\operatorname{im}\varphi_k \cap Z = \operatorname{im}\varphi_k \cap Z_1 \cap Z = Z_2 \cap Z = G_{J_k} \cap Z$. Hence ($1_k$) and ($2_k$) are satisfied. To verify (3_k) observe that if $J_k \cup J_l = J_m$ and $l < k$, then $m < l$ and by (3_l), $\operatorname{im}\varphi_m \subset \operatorname{im}\varphi_l$. By the definition, $\operatorname{im}\varphi_m \subset Z_2$, so $\operatorname{im}\varphi_m \subset \operatorname{im}\varphi_k$. Hence $\operatorname{im}\varphi_l \cap \operatorname{im}\varphi_k \supset \operatorname{im}\varphi_m$. We have $\operatorname{im}\varphi_l \cap \operatorname{im}\varphi_k = \operatorname{im}\varphi_l \cap \operatorname{im}\varphi_k \cap Z_1$, because $\operatorname{im}\varphi_l \subset Z_1$ as $l < k$. Hence $\operatorname{im}\varphi_l \cap \operatorname{im}\varphi_k = \operatorname{im}\varphi_l \cap Z_2$, because as we noted above $\operatorname{im}\varphi_k \cap Z_1 = Z_2$. Recall that $Z_2 = (G_{J_k} \cap Z) \cup \bigcup_{l'<k\colon J_{l'} \supset J_k} \operatorname{im}\varphi_{l'}$. By ($2_l$) and ($2_m$), $\operatorname{im}\varphi_l \cap G_{J_k} \cap Z = G_{J_l} \cap G_{J_k} \cap Z = G_{J_m} \cap Z \subset \operatorname{im}\varphi_m$. Let $l' < k$ such that $J_{l'} \supset J_k$ and observe that by (3_l) and ($3_{l'}$), $\operatorname{im}\varphi_l \cap \operatorname{im}\varphi_{l'} \subset \operatorname{im}\varphi_m$ as $J_m \subset J_l \cup J_{l'}$. Hence $\operatorname{im}\varphi_l \cap \operatorname{im}\varphi_k = \operatorname{im}\varphi_l \cap Z_2 \subset \operatorname{im}\varphi_m$, by the definition of Z_2. Hence (3_k) is satisfied. To verify (4_k) it is sufficient to see that if $J_k \subsetneq J_l$, then the restriction of φ_k to $\operatorname{im}\varphi_l$ is equal to the inclusion of $\operatorname{im}\varphi_l$ into X. But by the definition the restriction of φ_k to Z_2 is equal to the inclusion and $\operatorname{im}\varphi_l \subset Z_2$ by the definition of Z_2.

For every $i \in I$ let $k(i)$ denote an integer such that $J_{k(i)} = \{i\}$. Claim: the collection $\mathcal{F}' = \{F'_i = \operatorname{im}\varphi_{k(i)}\}_{i\in I}$ satisfies the assertion. Observe that $F'_{J_k} = \operatorname{im}\varphi_k$, by (3), therefore $\mathcal{F}'$ is a $\mathcal{U}$-adjustment of $\mathcal{F}$, as by (1) every φ_k is $\mathcal{U}$-close to the identity. Let $J_k \subsetneq J_{k_1}$. Since φ_{k_1} is a $Z(\mathcal{F})$-embedding, $F'_{J_{k_1}} = \operatorname{im}\varphi_{k_1}$ is a Z-set in F_{J_k}. Therefore $\varphi_k(F'_{J_{k_1}})$ is a Z-set in $F'_{J_k} = \operatorname{im}\varphi_k$, but by (4) $\varphi_k(F'_{J_{k_1}}) = F'_{J_{k_1}}$, therefore $\mathcal{F}'$ is a Z-collection. By (2) $\mathcal{F}'$ is equal to $\mathcal{G}$ on Z, so $\mathcal{F}'$ satisfies the assertion. □

9.2 Fitting closed $\mathcal{N}_n$-neighborhoods

Definition. Let $\mathcal{F} = \{F_i\}_{i\in I}$ denote a collection of subsets of a space X. We say that a set $A \subset X$ *fits* $\mathcal{F}$ if it is an $AE(n)$-space and if for each $J \subset I$ the intersection $A \cap F_J$ is an $AE(n)$-space.

Lemma 9.2. *Assume that a subset B of a space X fits a collection $\mathcal{F} = \{F_i\}_{i\in I}$ and $\mathcal{F}$ restricted to B is both a Z-collection and an $\mathcal{N}_n$-collection. Assume that $f\colon B \to X$ is an embedding such that $f^{-1}(F_i) = F_i \cap B$. Then $f(B)$ fits $\mathcal{F}$ and $\mathcal{F}$ restricted to $f(B)$ is both a Z-collection and an $\mathcal{N}_n$-collection.*

Proof. Let $\mathcal{F} = \{F_i\}_{i\in I}$ and let $F_J = \bigcap_{i\in J} F_i$ for each non-empty subset J of I. We have $F_J \cap f(B) = F_J \cap \operatorname{im} f = f(f^{-1}(F_J)) = f(F_J \cap B)$, the last equality by the assumption that $f^{-1}(F_i) = F_i \cap B$ for each $i \in I$. By the assumption B fits $\mathcal{F}$, so $F_J \cap B$ is $AE(n)$ for each $J \subset I$ and so is $F_J \cap f(B)$. Therefore $\mathcal{F}$ fits $f(B)$. The homeomorphism $f\colon B \to f(B)$ maps collection $\{F_i \cap B\}_{i\in I}$ onto a collection $\{F_i \cap f(B)\}_{i\in I}$. Since the first collection is a Z-collection and an $\mathcal{N}_n$-collection in B, the second collection is a Z-collection and an $\mathcal{N}_n$-collection in $f(B)$. □

Lemma 9.3. *Assume that* $\mathcal{G} = \{G_i\}_{i\in I}$ *is an* $\mathcal{N}_n$*-collection and a* Z*-collection in an* $\mathcal{N}_n$*-space* Y*,* V *is an open subset of* Y *and* f *is a map from an at most* n*-dimensional Polish space* X *into* Y*. If* f *is a closed embedding into* V *on* $f^{-1}(V)$ *and* A *is a closed (in* Y*) subset of* V*, then* f *is approximable* rel $f^{-1}(A)$ *by closed embeddings* g *such that* $g^{-1}(G_i) = f^{-1}(G_i)$ *for each* $i \in I$*. Moreover, if* Z *is a subset of* X *and* $f_{|Z}$ *is a* $Z(\mathcal{G})$*-embedding, then such approximations exist* rel $f^{-1}(A) \cup Z$*.*

Proof. The proof follows the proof of proposition 2.39. Let $\mathcal{V}$ be an open cover of Y. Let $U = Y \setminus A$ and let $\mathcal{U}$ be an open cover of U obtained via lemma 2.38 applied to U and Y. Without a loss of generality, we may assume that $\mathcal{U}$ refines $\mathcal{V}$. By proposition 3.4, the restriction $\mathcal{G}/U$ of $\mathcal{G}$ to U is a Z-collection in U and the restriction of f to $Z \cap f^{-1}(U)$ is a $Z(\mathcal{G}/U)$-embedding into U. By theorem 2.4 and by theorem 2.16, U is an $\mathcal{N}_n$-space and $\mathcal{G}/U$ is an $\mathcal{N}_n$-collection, hence by theorem 3.9, there exists a $\mathcal{U}$-approximation g of $f_{|f^{-1}(U)}$ rel $Z \cap f^{-1}(U)$ by a closed embedding into U such that $g^{-1}(G_i) = f^{-1}(G_i)$ for each $i \in I$. By lemma 2.38 and by the choice of $\mathcal{U}$, the map $g \cup f_{|f^{-1}(U)}$ is a closed embedding into U and is $\mathcal{V}$-close to f. ☐

Lemma 9.4. *Let a finite collection* $\mathcal{F}$ *of subsets of an* $\mathcal{N}_n$*-space* X *be both an* $\mathcal{N}_n$*-collection and a* Z*-collection. Then every* $Z(\mathcal{F})$*-set* $A \subset X$ *that fits* $\mathcal{F}$ *has arbitrarily small closed* $\mathcal{N}_n$*-neighborhoods* B *that fit* $\mathcal{F}$*. Moreover, we may require that* $\mathcal{F}$ *restricted to* B *is both a* Z*-collection and an* $\mathcal{N}_n$*-collection of subsets of* B*.*

Observe that we do not assume above that A is an $\mathcal{N}_n$-space.

Proof. The proof is by induction on the cardinality of $\mathcal{F}$. Assume that $\mathcal{F}$ is empty. We have to prove that if A is an absolute extensor in dimension n and a Z-set in X, then it has arbitrarily small closed $\mathcal{N}_n$-neighborhoods that are absolute neighborhood extensors in dimension n. Fix an open neighborhood U of A and let V be another open neighborhood of A whose closure lies in U. By lemma 2.9 and by remark 2.40, there is a closed $\mathcal{N}_n$-neighborhood B of A with a property that every partial map from an n-dimensional Polish space into B extends over the entire domain to a map into V. By theorem 2.15, there exists a closed embedding $f\colon B \to \nu^n$. By the choice of B, there is an extension $g\colon \nu^n \to V$ of f^{-1}. By corollary 3.7, there is an approximation h of g within an $\mathcal{N}_n$-cover $\{X, B\}$ by a closed embedding with image disjoint from A. By Whitehead's characterization and by lemma 2.21, if the approximation is close enough (which we assume), then the inclusion of $h(f(B))$ into B is an n-homotopy equivalence. Consider a space $C = B \sqcup_{h(f(B))} h(\nu^n)$, a disjoint sum of B and $h(\nu^n)$ glued along $h(f(B))$. By corollary 3.8, it is an $\mathcal{N}_n$-space. By lemma 2.24, it is an $AE(n)$-space. Let $i\colon C \to B$ denote a map induced by inclusions $B \subset X$ and $h(\nu^n) \subset X$. It is an embedding on some neighborhood of A. Hence, by proposition 2.39, it can be approximated by a closed embedding rel some neighborhood of A. The image of this embedding is a closed $\mathcal{N}_n$-neighborhood of A that is an $AE(n)$-space. We may assume that both approximations done above are close to the identity, so this image lies in U. The basis step of induction is done.

Let $\mathcal{F} = \{F_i\}_{i\in I}$, fix $j \in I$, let $J = I \setminus \{j\}$ and assume that the theorem is proved for collections of cardinality J. Fix a pair of open neighborhoods U, V of A as above. Let $\mathcal{F}' = \{F_i\}_{i\in J}$. By the inductive assumption, there is a closed $\mathcal{N}_n$-neighborhood B' of A in X that fits $\mathcal{F}'$ and lies in V. Let $\mathcal{F}'' = \{F_i \cap F_j\}_{i\in J}$. Since $\mathcal{F}$ is both a Z-collection and an $\mathcal{N}_n$-collection, $\mathcal{F}''$ is a Z-collection and an $\mathcal{N}_n$-collection in F_j. Since A is a $Z(\mathcal{F})$-set, $A\cap F_j$ is a $Z(\mathcal{F}'')$-set. By the inductive assumption, there is a closed $\mathcal{N}_n$-neighborhood B'' of $A \cap F_j$ in F_j that fits $\mathcal{F}''$ and lies in $F_j \cap \operatorname{Int}_X B'$. Let $B''' = B' \sqcup_{B''} F_j$ be a disjoint union of B' and F_j glued along B''. Let $f\colon B''' \to X$ be a map induced by inclusions on B' and F_j. By the assumption that $\mathcal{F}$ is a Z-collection, F_j is a $Z(\mathcal{F}')$-set. By the construction, f is an embedding on an open neighborhood of A. By lemma 9.3 applied to a Z-collection $\mathcal{F}'$, map f and a set $F_j \subset B'''$ (f restricted to F_j is a $Z(\mathcal{F}')$-embedding), there exists an approximation of f by a closed embedding g that is equal to f on an union of F_j with a neighborhood of A and that satisfies equality $g^{-1}(F_i) = f^{-1}(F_i)$ for each $i \in J$. Let $B = g(B')$. It is a closed $\mathcal{N}_n$-neighborhood of A. We will show that it fits $\mathcal{F}$ and that the restriction of $\mathcal{F}$ to B is both a Z-collection and an $\mathcal{N}_n$-collection in B, thus ending the inductive step.

By the construction, B' fits $\mathcal{F}'$ and the restriction of $\mathcal{F}'$ to B' is both an $\mathcal{N}_n$-collection and a Z-collection. Hence by lemma 9.2 applied to an embedding $g_{|B'}$, B fits $\mathcal{F}'$ and the restriction of $\mathcal{F}'$ to B is both an $\mathcal{N}_n$-collection and a Z-collection. Let J be a subset of I such that $j \in J$. We have $F_J \cap B = (F_j \cap B) \cap F_J = B'' \cap F_J$. Since by the construction B'' fits $\mathcal{F}''$, the intersection $B'' \cap F_J$ is an $AE(n)$-space and so is $B \cap F_J$. Hence B fits the entire collection $\mathcal{F}$. Again, by the construction $\mathcal{F}''$ restricted to B'' is an $\mathcal{N}_n$-collection, hence the entire collection $\mathcal{F}$ restricted to B is an $\mathcal{N}_n$-collection. To see that this restriction is a Z-collection in B observe that B'' is a Z-set in B (because F_j is a Z-set in X and B'' is a closed subset of F_j that lies in the interior of B), $\mathcal{F}''$ is a Z-collection in B'' (by the construction) and $\mathcal{F}'$ is a Z-collection in B (by lemma 9.2). $\square$

9.3 Patching of holes

Lemma 9.5. *Assume that B is a closed $\mathcal{N}_n$-subspace and a Z-set in an $\mathcal{N}_n$-space X. If Δ is a k-dimensional simplex embedded in X such that $\partial\Delta = \Delta \cap B$ and N is an open neighborhood of Δ, then there exists a closed (in X) $\mathcal{N}_n$-subset A of N such that $A \cap B$ is an $\mathcal{N}_n$-space, $\Delta \subset A$ and the inclusions $\partial\Delta \subset A \cap B$ and $\Delta \subset A$ are n-homotopy equivalences.*

Proof. Let $\delta \subset \nu^n$ be a standard k-dimensional simplex isometrically embedded into a k-dimensional hyperplane that lies in ν^n. Fix an $\varepsilon > 0$ and let $U = \{x \in \nu^n \colon d(x, \delta) < \varepsilon\}$ and $V = \{x \in \nu^n \colon d(x, \partial\delta) < \varepsilon\}$. Assume that ε is so small that the inclusion $\partial\delta \subset V$ is an n-homotopy equivalence (cf. proposition 2.27). The inclusion $\delta \subset U$ is also an n-homotopy equivalence. By corollary 2.37 applied with $D = \delta \setminus \partial\delta$, the inclusion $V \subset U$ can be approximated $\operatorname{rel} \partial\delta$ by a closed embedding $g\colon V \to U$ with image disjoint from $\delta \setminus \partial\delta$. Let $C = g(V)$. Since the inclusion $\partial\delta \subset V$ is an n-homotopy equivalence, the inclusion $g(\partial\delta) = \partial\delta \subset C = g(V)$ is an n-homotopy equivalence. Therefore, by theorem 2.23, there is a retraction $C \to \partial\delta$. Since δ is an absolute extensor and $\delta \cap C = \partial\delta$, this retraction can be extended to a retraction $r\colon U \to \delta$. Note that by the construction, $r(C) \subset \partial\delta$. Let $H\colon \delta \to \Delta$ be any homeomorphism. By theorem 3.6, $(H \circ r)\colon U \to \Delta$ can be approximated by a closed embedding $h\colon U \to X$ $\operatorname{rel} \delta$ such that $h(C) \subset B$. By the assumption that B is a Z-set, and by corollary 2.37 applied with $D = B \setminus h(C)$ and $\operatorname{rel} F \cup \delta$, we may assume that $h(C) = h(U) \cap B$. Observe that $\Delta \subset h(U)$, $\partial\Delta \subset h(C) = h(U) \cap B$ and these inclusions are n-homotopy equivalences. Hence letting $A = h(C)$ finishes the proof. □

Remark 9.6. Assume that Δ, A and B are closed subsets of $A \cup B$ such that Δ, A, B and their intersections are at most n-dimensional $ANE(n)$-spaces. If inclusions $\partial\Delta \subset A \cap B$, $\Delta \subset A$ hold and are n-homotopy equivalences, then the inclusion $\Delta \cup B \subset A \cup B$ is an n-homotopy equivalence.

Proof. By corollary 2.25 applied with $X = A$, $A_1 = \Delta$ and $A_2 = A \cap B$, the inclusion $\Delta \cup (A \cap B) \subset A$ is an n-homotopy equivalence. Hence, by lemma 2.24 applied with $A_1 = \Delta \cup B$ and $A_2 = A$, the inclusion $\Delta \cup B \subset A \cup B$ is an n-homotopy equivalence. □

Chapter 10

Core of a cover

Our goal is a proof of theorem 6.7 and we will attain it by "patching of holes" in a cover. In this chapter we introduce the notion of an n-core of a cover that aids the "patching" process and allows us to control it.

Definition. Assume that $\mathcal{F} = \{F_i\}_{i \in I}$ is a cover of a space X. A pair $\langle K, \mathcal{K} \rangle$ consisting of an at most n-dimensional locally finite simplicial complex K embedded as a closed subset of X and of an interior cover $\mathcal{K} = \{K_i\}_{i \in I}$ of K by subcomplexes is an *n-core of* $\mathcal{F}$ if for every $J \subset I$ the inclusion $K_J \subset F_J$ holds and induces an n-homotopy equivalence. If additionally $\mathcal{K}$ is equal to $\mathcal{F}$ on K, then we say that the core is *exact*. We shall say that K is an exact n-core of $\mathcal{F}$ if $\langle K, \{F_i \cap K\}_{i \in I} \rangle$ is an n-core of $\mathcal{F}$.

Observe that the conditions of the above definition imply that collections $\mathcal{F}$ and $\mathcal{K}$ must be isomorphic, because the empty space is not n-homotopy equivalent to an non-empty space.

Definition. Let $\mathcal{F}$ be a cover of a space X, let $\langle K, \mathcal{K} \rangle$ be an n-core of $\mathcal{F}$ and let $\mathcal{U}$ be an open cover of X. We say that a cover $\mathcal{G}$ of X is a *$\mathcal{U}$-deformation of $\mathcal{F}$ with fixed* $\langle K, \mathcal{K} \rangle$ if $\langle K, \mathcal{K} \rangle$ is an n-core of $\mathcal{G}$ and if G_i is a subset of $\operatorname{st}_{\mathcal{U}} F_i$ for each pair of corresponding elements $F_i \in \mathcal{F}$ and $G_i \in \mathcal{G}$ (note that $\mathcal{F}$ and $\mathcal{G}$ must be isomorphic because they have the same n-core). Sometimes we shall write that $\mathcal{G}$ is a $\mathcal{U}$-deformation of $\mathcal{F}$ or that $\mathcal{G}$ is a deformation of $\mathcal{F}$, if $\mathcal{U}$ and $\langle K, \mathcal{K} \rangle$ are known from the context.

Theorem 10.1 is the main theorem of this chapter.

Theorem 10.1. *Every closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{F}$ with an n-core $\langle K, \mathcal{K} \rangle$ admits arbitrarily small deformation to a closed interior $\mathcal{N}_n$-cover of which $\langle K, \mathcal{K} \rangle$ is an exact n-core. Moreover, if L is a subcomplex of K and $\mathcal{F}$ is equal to $\mathcal{K}$ on $K \setminus L$, then we may additionally require that the deformed cover is equal to $\mathcal{F}$ on a complement of arbitrarily small open neighborhood of L.*



Figure 10.1: Sample application of theorem 10.1.

We prove it in section 10.3. Here we illustrate its typical usage. Consider a cover $\mathcal{F}$ drawn on figure 10.1(a). It contains three sets: F_1 (dark one), F_2 (the large transparent box - it contains F_1) and F_3 (light one). Note that $\mathcal{F}$ is not 1-regular: F_3 is disconnected. We want to "patch" F_3 by attaching arc $\overline{ab}$ to it. But $\overline{ab}$ crosses F_1 and the cover $\{F_1, F_2, F_3 \cup \overline{ab}\}$, although it is 1-regular, is not isomorphic to $\mathcal{F}$. Theorem 10.1 gives a solution to this problem, as follows.

Consider a graph $K = \overline{ab} \cup \overline{cd}$ and its cover $\mathcal{K} = \{\{c\}, K, \{a, b\}\}$. A pair $\langle K, \mathcal{K} \rangle$ is a 1-core of $\mathcal{F}$. Assume that $\mathcal{F}$ is a closed interior $\mathcal{N}_1$-cover and apply theorem 10.1 to it. We get a cover $\{F'_1, F'_2, F'_3\}$ drawn on picture 10.1(b) that has $\langle K, \mathcal{K} \rangle$ as an exact 1-core. Now the construction described above works: if we attach $\overline{ab}$ to F'_3 we get a 1-regular cover that is isomorphic to $\mathcal{F}$.

10.1 The existence of an n-core

First, to show that theorem 10.1 has a point, we prove that every closed star-finite locally finite $\mathcal{N}_n$-cover has an n-core. To this end we introduce an auxiliary notion of an outer n-core.

Definition 10.2. Assume that $\mathcal{F} = \{F_i\}_{i \in I}$ is a cover of a space X. A triple $\langle K, \mathcal{K}, k \rangle$ is an *outer n-core of $\mathcal{F}$* if

(1) K is an at most n-dimensional locally finite simplicial complex,

(2) $\mathcal{K} = \{K_i\}_{i \in I}$ is a cover of K by subcomplexes,

(3) $k \colon K \to X$ is a map such that $k(K_i) \subset F_i$ and for every $J \subset I$ the restriction of k to K_J is an n-homotopy equivalence of K_J and F_J.

For example, if $\mathcal{F}$ is an n-regular cover and λ is its anticanonical map, then the triple $\langle N(\mathcal{F}), \mathcal{B}_{\mathcal{F}}, \lambda\rangle$ is an outer n-core of $\mathcal{F}$.

Lemma 10.3. *Assume that $\mathcal{F} = \{F_i\}_{i\in I}$ is a star-finite $ANE(n)$-cover of a space X and $\langle L, \mathcal{L}, l\rangle$ is a triple that satisfies all conditions of definition 10.2 with the exception that the restriction of l to L_J need not to be an n-homotopy equivalence of L_J and F_J. Then there is an outer n-core $\langle K, \mathcal{K}, k\rangle$ of $\mathcal{F}$ such that L is a subcomplex of K, $\mathcal{K}$ restricted to L is equal to $\mathcal{L}$ and k is an extension of l. Moreover, if $\mathcal{L}$ is an interior cover, then we may additionally require that $\mathcal{K}$ is also an interior cover.*

Proof. Let $\mathcal{I} = \{J \subset I \colon F_J \neq \emptyset\}$. Order $\mathcal{I}$ into a sequence $\{J_k\}$ non increasing in the order by inclusion. Such ordering exists because $\mathcal{F}$ is star-finite. By a recursive application of lemma 2.26 we construct a sequence of triples $\langle W_k, \mathcal{W}_k, w_k\rangle$, starting with $\langle W_0, \mathcal{W}_0, w_0\rangle = \langle L, \mathcal{L}, l\rangle$, such that

(1) W_{k-1} is a subcomplex of an at most n-dimensional locally finite complex W_k,

(2) $\mathcal{W}_k = \{W_i^k\}_{i\in I}$ is a cover of W_k by subcomplexes and

$$W_i^k = \begin{cases} W_i^{k-1} & i \notin J_k \\ W_i^{k-1} \cup (W_k \setminus W_{k-1}) & i \in J_k, \end{cases}$$

(3) w_k is an extension of w_{k-1} onto W_k such that w_k restricted to $W_{J_k}^k$ is an n-homotopy equivalence of $W_{J_k}^k$ and F_{J_k}.

We take $K = \bigcup W_k$ (with the direct limit topology), $k = \bigcup w_k$ and $\mathcal{K} = \{K_i = \bigcup W_i^k\}_{i\in I}$. Obviously the triple $\langle K, \mathcal{K}, k\rangle$ has all required properties; the complex K is locally finite because of the assumption that $\mathcal{F}$ is star-finite.

Now assume that $\mathcal{L}$ is an interior cover. A simple modification of the construction described above assures that the constructed cover $\mathcal{K}$ is also an interior cover. Let $W_{k-1/2}$ denote a complex $W_{k-1} \cup C_k$, where $C_k = W_{J_k}^{k-1} \times \{0,1\} \cup W_{J_k}^{k-1(n-1)} \times [0,1]$ - a cylinder taken over $W_{J_k}^{k-1}$ with removed interiors of n-dimensional simplices. Let $p \colon W_{k-1/2} \to W_{k-1}$ be a natural projection. Let $\mathcal{W}_{k-1/2} = p^{-1}(\mathcal{W}_{k-1})$ and $w_{k-1/2} = w_{k-1} \circ p$. Construct $\langle W_k, \mathcal{W}_k, w_k\rangle$ as above, but over $\langle W_{k-1/2}, \mathcal{W}_{k-1/2}, w_{k-1/2}\rangle$ instead of $\langle W_{k-1}, \mathcal{W}_{k-1}, w_{k-1}\rangle$ and in such a way, that the closure of $W_k \setminus W_{k-1}$ intersects only $W_{J_k}^{k-1} \times \{1\}$. It is easy to verify that with such construction if $\mathcal{L}$ is an interior cover, then $\mathcal{K}$ is also an interior cover. □

By proving theorem 10.5 we will show that every closed star-finite locally finite $\mathcal{N}_n$-cover has an n-core.

Lemma 10.4. *For each closed locally finite $ANE(n)$-cover $\mathcal{F} = \{F_i\}_{i\in I}$ of a space X there exists an open cover $\mathcal{U}$ of X satisfying the following condition.*

> *For each $J \subset I$ and each map φ from at most $(n-1)$-dimensional space into $F_J = \bigcap i \in JF_i$, every map into F_J that is $\mathcal{U}$-close to φ is homotopic to it (in F_J).*

Proof. By theorem 2.4(3) for every $J \subset I$ there is an open cover $\mathcal{U}_J$ of F_J such that every two $\mathcal{U}_J$-close maps into F_J from at most $(n-1)$-dimensional Polish space are homotopic. Since $\mathcal{F}$ is closed and locally finite there is an open cover $\mathcal{U}$ of X such that for every $J \subset I$ the cover $\mathcal{U}$ restricted to F_J refines $\mathcal{U}_J$. This cover satisfies our claim. □

Theorem 10.5. *Let $\mathcal{F} = \{F_i\}_{i \in I}$ be a closed star-finite locally finite $\mathcal{N}_n$-cover of a space X. Let L be a locally finite simplicial complex embedded as a closed subset of X. Let $\mathcal{L} = \{L_i\}_{i \in I}$ be an interior cover of L by subcomplexes such that $L_i \subset F_i$ for each $i \in I$. Then there exists an n-core $\langle K, \mathcal{K} \rangle$ of $\mathcal{F}$ such that L is a subcomplex of K and $\mathcal{K}$ is equal to $\mathcal{L}$ on L. Moreover, if Z is a $Z(\mathcal{F})$-set, then we may additionaly require that $K \setminus L$ is disjoint from Z.*

Proof. Let $\mathcal{F} = \{F_i\}_{i \in I}$ be a closed star-finite locally finite $\mathcal{N}_n$-cover of a space X. By lemma 10.4, there exist an open cover $\mathcal{U}$ of X that satisfies the following condition.

> (*) *For each $J \subset I$ and each map φ from at most $(n-1)$-dimensional space into F_J, every map into F_J that is $\mathcal{U}$-close to φ is homotopic to it (in F_J).*

Let l denote the inclusion of L into X. By lemma 10.3 the triple $\langle L, \mathcal{L}, l \rangle$ can be enlarged to an outer n-core $\langle W, \mathcal{W}, w \rangle$ of $\mathcal{F}$. By theorem 3.6 and by remark 3.5 there exists a $\mathcal{U}$-approximation of w within $\mathcal{F}$ rel L by a closed embedding v with image disjoint from $Z \setminus L$. Then $v_{|W_J}$ is $\mathcal{U}$-close to $w_{|W_J}$ so by (*), the inclusion of $v(W_J)$ into F_J is an n-homotopy equivalence (see the proof of Whitehead's characterization). Therefore $\langle v(W), v(\mathcal{W}) \rangle$ is an n-core of $\mathcal{F}$. Since $v_{|L}$ is equal to l, L is a subcomplex of $v(W)$ and $v(\mathcal{W})$ is equal to $\mathcal{L}$ on L. □

Remark 10.6. Let $\mathcal{F}$ be a closed star-finite locally finite $\mathcal{N}_n$-cover of a space X. Let $\mathcal{U}$ be a cover satisfying the condition stated in lemma 10.4. Assume that $\langle K, \mathcal{K} \rangle$ is an n-core of $\mathcal{F}$, where $\mathcal{K} = \{K_i\}_{i \in I}$. If $\mathcal{G} = \{G_i\}_{i \in I}$ is a $\mathcal{U}$-adjustment of $\mathcal{F}$ such that $K_i \subset G_i$ for each $i \in I$, then $\langle K, \mathcal{K} \rangle$ is an n-core of $\mathcal{G}$. To prove it fix $J \subset I$ and let $h \colon G_J \to F_J$ be a homeomorphism that is $\mathcal{U}$-close to the identity. The inclusion of $h(K_J)$ into F_J is $\mathcal{U}$-close to the identity on K_J, hence it is n-homotopic to it. Therefore the inclusion of $K_J = h^{-1}(h(K_J))$ into $G_J = h^{-1}(F_J)$ is a weak n-homotopy equivalence.

10.2 An n-core of a limit of a sequence of deformations

The purpose of this section is to prove an analogue of proposition 4.3 for sequences of deformations. Observe that, even if deformations are done on a discrete family of open subsets, the limit of the sequence doesn't have to be a deformation of the original cover. Figure 10.2 gives an appropriate example: the singleton K is weak 1-homotopy equivalent to the set F_n obtained from F_0 by disconnecting n "bridges". But it is not weak 1-homotopy equivalent to the set that we get in the limit of this construction, because $F_\infty = \bigcap F_n$ is disconnected.



Figure 10.2: K is not a 1-core of $F_\infty = \bigcap F_n$.

Lemma 10.7. *Let* $\{U_m\}_{m\geq 1}$ *be a discrete family of open subsets of a space* X *and for each* m *let* V_m *be an open subset of* U_m. *Let* $\mathcal{F}_m$ $(m \geq 0)$ *be a closed star-finite locally finite* $\mathcal{N}_n$*-cover of* $X \setminus \bigcup_{k>m} V_k$ *and assume that* $\mathcal{F}_m$*'s are pairwise isomorphic. Assume that* $\mathcal{F}_{m-1}$ *is equal to* $\mathcal{F}_m$ *on* $X \setminus U_m$ *and that it is a shrinking of* $\mathcal{F}_m$ *on* $U_m \setminus V_m$. *If* $\langle K, \mathcal{K}\rangle$ *is an* n*-core of* $\mathcal{F}_m$ *for each* m, *then it is an* n*-core of* $\lim_{m\to\infty} \mathcal{F}_m$.

Lemma 10.7 is a direct consequence of the following lemma.

Lemma 10.8. *Let* $\{U_m\}_{m\geq 1}$ *be a discrete family of open subsets of a space* X *and for each* m *let* V_m *be an open subset of* U_m. *Let* $\mathcal{F}_m$ $(m \geq 0)$ *be a closed star-finite locally finite* $\mathcal{N}_n$*-cover of* $X \setminus \bigcup_{k>m} V_k$ *and assume that* $\mathcal{F}_m$*'s are pairwise isomorphic. If* $\mathcal{F}_m$ *is equal to* $\mathcal{F}_{m-1}$ *on the complement of* U_m, *then* $\mathcal{F} = \lim_{m\to\infty} \mathcal{F}_m$ *is a closed star-finite locally finite* $\mathcal{N}_n$*-cover of* X. *Assume that* $\mathcal{F}_m$ *is a swelling of* $\mathcal{F}_{m-1}$ *on* $U_m \setminus V_m$ *and let* $\langle K_m, \mathcal{K}_m\rangle$ *be an* n*-core of* $\mathcal{F}_m$, *where* $\mathcal{K}_m = \{K_i^m\}_{i\in I}$. *If* K_{m-1} *is a subcomplex of* K_m, $\mathcal{K}_{m-1}$ *is a shrinking of* $\mathcal{K}_m$ *and* $K = \bigcup K_m$ *is a locally finite complex embedded as a closed subset of* X, *then* $\langle K, \mathcal{K}\rangle$ *is an* n*-core of* $\mathcal{F}$, *where* $\mathcal{K} = \{K_i\}_{i\in I}$ *and* $K_i = \bigcup_{m\geq 1} K_i^m$.

Proof. To prove that $\mathcal{F}$ is a closed locally finite $\mathcal{N}_n$-cover of X it is sufficient to observe that the sequence $\mathcal{F}_m$ stabilizes locally and consists of closed locally finite $\mathcal{N}_n$-covers. Also $\mathcal{F}$ must be star-finite because $\mathcal{F}_m$'s are pairwise isomorphic. What is left is to check that $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}$. Let $\mathcal{F}_m = \{F_i^m\}_{i\in I}$ and $\mathcal{F} = \{F_i\}_{i\in I}$

where $F_i = \bigcap_j \bigcup_{k\geq j} F_i^k$. Fix $J \subset I$. By the definition it is sufficient to prove that the inclusion of K_J into F_J is a weak n-homotopy equivalence.

Fix $k < n$ and a map $\varphi\colon S^k \to F_J$. Since S^k is compact $\operatorname{im}\varphi$ intersects only finitely many U_m's so there is M such that $\operatorname{im}\varphi \cap U_m = \emptyset$ for every $m \geq M$. Therefore φ is homotopic with a map into $K_M \cap F_J^M$ in $F_J^M \setminus \bigcup_{m>M} V_m$. But $F_J^M \setminus \bigcup_{m>M} V_m \subset F_J$ and $K_M \cap F_J^M \subset K \cap F_J$ so the inclusion induces epimorphisms on homotopy groups of dimensions less than n.

Fix a map $\varphi\colon S^k \to K \cap F_J$ and assume that it has an extension $\Phi\colon D^{k+1} \to F_J$. By compactness and local finiteness of K there is M such that $\operatorname{im}\varphi \subset K_M$ and $\operatorname{im}\Phi \subset F_J \backslash \bigcup_{m>M} U_m \subset F_J^M$. But K_M is a core of $\mathcal{F}_M$ therefore φ must be contractible in $K_M \cap F_J^M$. But $K_M \cap F_J^M \subset K \cap F_J$ so the inclusion induces monomorphisms on homotopy groups of dimensions less than n and we are done. □

10.3 Proof of theorem 10.1

Let K be a locally finite simplicial complex embedded as a closed subset of an $\mathcal{N}_n$-space X and let L be a subcomplex of K. Let U_L be an open neighborhood of L. Let $\mathcal{K} = \{K_i\}_{i\in I}$ be an interior cover of K by subcomplexes. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a star-finite interior $\mathcal{N}_n$-cover of X such that $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}$. We fix a triangulation τ of K such that for each simplex δ in τ there exist i and j in I such that $\delta \subset \operatorname{Int}_X F_i$ (this is possible because $\mathcal{F}$ is an interior cover) and $\delta \subset \operatorname{Int}_K K_j$ (if a triangulation does not have this property, it suffices to take its first barycentric subdivision, because $\mathcal{K}$ is an interior cover). Let τ_L be a subset of τ that contains simplices of L.

For each $l \in \{0, 1, \ldots, n+1\}$ and $m \in \{0, 1, \ldots, \infty\}$ we let $\mathcal{F}_{l,m} = \{F_i^{l,m}\}_{i\in I}$ denote a collection (not yet defined) of subsets of X. Let $(1_{l,m})$, $(2_{l,m})$, $(3_{l,m})$ and $(4^{\delta,i}_{l,m})$ denote the following statements, with $0 \leq l \leq n+1$, $0 \leq m \leq \infty$, $\delta \in \tau$ and $i \in I$.

$(1_{l,m})$ $\mathcal{F}_{l,m}$ is a closed star-finite interior $\mathcal{N}_n$-cover of X.

$(2_{l,m})$ $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}_{l,m}$.

$(3_{l,m})$ For each $\delta \in \tau$ there is $i \in I$ such that $\delta \subset \operatorname{Int}_X F_i^{l,m}$.

$(4^{\delta,i}_{l,m})$ $\delta \subset \operatorname{Int}_K K_i \Rightarrow \delta \subset \operatorname{Int}_X F_i^{l,m}$.

On a very general level, the proof is organized as follows. We let $\mathcal{F}_{0,0} = \mathcal{F}$ and construct $\mathcal{F}_{l,m}$ recursively. In (l,m)th step of the construction, with $0 \leq l \leq n+1$ and $0 \leq m < \infty$, we construct a collection $\mathcal{F}_{l,m+1}$. If $l < n+1$, then $\mathcal{F}_{l,m+1}$ is a deformation of $\mathcal{F}_{l,m}$ with fixed n-core $\langle K, \mathcal{K}\rangle$. If $l = n+1$, then $\mathcal{F}_{l,m+1}$ is an adjustment of $\mathcal{F}_{l,m}$. In lth limit step of the construction, with $0 \leq l \leq n+1$, we let $\mathcal{F}_{l,\infty} = \lim_{m\to\infty} \mathcal{F}_{l,m}$. We also let $\mathcal{F}_{l+1,0}$ denote $\mathcal{F}_{l,\infty}$ for each $0 \leq l < n+1$. The collection $\mathcal{F}_{n+1,\infty}$ shall satisfy the assertion of the theorem.

A general constraint on the construction is that conditions $(1_{l,m})$, $(2_{l,m})$ and $(3_{l,m})$ are satisfied for all l and m. Observe that $(1_{0,0})$, $(2_{0,0})$ and $(3_{0,0})$ are satisfied by the assumptions of the theorem (we did let $\mathcal{F}_{0,0} = \mathcal{F}$).

The first part of the proof includes the first $n+1$ limit steps (i.e. the construction of $\mathcal{F}_{0,\infty}, \mathcal{F}_{1,\infty}$ up to $\mathcal{F}_{n,\infty}$). In lth limit step of the construction, with $0 \leq l \leq n$, we construct a cover $\mathcal{F}_{l,\infty}$ that satisfies $(4^{\delta,i}_{l,\infty})$ for each at most l-dimensional simplex $\delta \in \tau$ and each $i \in I$. A general constraint in the first part of the construction is that in its (l,m)th step, with $0 \leq l \leq n$ and $0 \leq m < \infty$, *the fourth condition is preserved*, i.e. the implication $(4^{\delta,i}_{l,m}) \Rightarrow (4^{\delta,i}_{l,m+1})$ is true for each at most l-dimensional simplex δ of triangulation τ and each $i \in I$. The first part of the proof concludes with a construction of $\mathcal{F}_{n,\infty}$ that satisfies $(1_{n,\infty})$, $(2_{n,\infty})$, $(3_{n,\infty})$ and, for each $\delta \in \tau$ and each $i \in I$, $(4^{\delta,i}_{n,\infty})$ (no other property of $\mathcal{F}_{n,\infty}$ is important in the second part of the proof).

The second part of the proof includes the last limit step. It starts with a cover $\mathcal{F}_{n+1,0} = \mathcal{F}_{n,\infty}$ satisfying $(1_{n+1,0})$, $(2_{n+1,0})$, $(3_{n+1,0})$ and $(4^{\delta,i}_{n+1,0})$ for each $\delta \in \tau$ and $i \in I$. A cover $\{V_m\}_{m\geq 0}$ of K, consisting of open subsets of X, is picked and in $(n+1,m)$th step of the construction an adjustment $\operatorname{rel} X \setminus V_m$ of $\mathcal{F}_{n+1,m}$ is done in such a way that $\mathcal{F}_{n+1,m+1}$ is equal to $\mathcal{K}$ on V_m. Under some additional constraints, the limit cover $\mathcal{F}_{n+1,\infty} = \lim_{m\to\infty} \mathcal{F}_{n+1,m}$ satisfies the assertion of the theorem.

First part.

In lth limit step, with $0 \leq l \leq n$, we shall take $\mathcal{F}_{l,\infty} = \lim_{m\to\infty} \mathcal{F}_{l,m}$ and argue that $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}_{l,\infty}$. To this end we shall apply lemma 10.7. To be able to do that we must construct an auxiliary n-core of $\mathcal{F}_{0,0}$ that is disjoint from K. We start with its construction.

Let $K' = K \times \{0,1\} \cup K^{(n-1)} \times [0,1]$ and let $p\colon K' \to K$ be a projection along $[0,1]$. By theorem 3.6, there exists an approximation $\tilde{p}\colon K' \to X$ of p within $\mathcal{F}_{0,0}$ $\operatorname{rel} K \times \{0\}$ by a closed embedding. Let $L = \tilde{p}(K \times \{1\})$ and $M = \tilde{p}(K')$. Note that $K = \tilde{p}(K \times \{0\}) \subset M$. Let $\mathcal{L} = \{L_i\}_{i\in I}$ and $\mathcal{M} = \{M_i\}_{i\in I}$, where $M_i = \tilde{p}(p^{-1}(K_i))$ and $L_i = M_i \cap L$. Note that $\langle L, \mathcal{L}\rangle$ and $\langle M, \mathcal{M}\rangle$ are n-cores of $\mathcal{F}_{0,0}$ and L is disjoint from some open neighborhood of K. Also note that if $M_i \subset F^{l,k}_i$ for some $0 \leq l \leq n$ and $0 \leq k \leq \infty$, and each $i \in I$, then $\langle L, \mathcal{L}\rangle$ is an n-core of $\mathcal{F}_{l,k}$ if and only if $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}_{l,k}$.

Let $\delta_0, \delta_1, \ldots$ be a sequence of all 0-dimensional simplices of τ_L. Let $J_k = \{i \in I\colon \delta_k \subset \operatorname{Int}_X F^{0,0}_i\}$. Fix an open neighborhood U_k of δ_k in X whose closure lies in the interior of $F^{0,0}_{J_k}$ and meets only those simplices of K that contain δ_k. Without a loss of generality we may assume that $\{U_k\}$ is a discrete family and each U_k is disjoint from L. Let $h_k\colon U_k \to U_k$ be an approximation of id_{U_k} within $\mathcal{F}_{0,0}$ by a closed embedding with image disjoint from δ_k. We assume that the approximation is such that $h_k \cup id_{X \setminus U_k}$ is a closed embedding (see lemma 4.1). We let $V_k = U_k \setminus h_k(U_k)$. Note that the choice of sets V_k will be important only when passing to the limit and for now the reader may think that $V_k = U_k$.

We are going to construct a sequence $\mathcal{F}_{0,1}, \mathcal{F}_{0,2}, \ldots$ in such a way that the following conditions will be satisifed for each $k > 0$.

(5_k) $\mathcal{F}_{0,k}$ is a deformation of $\mathcal{F}_{0,k-1}$ $\operatorname{rel} X \setminus V_m$, for m such that $2m < k \leq 2m+2$,

(6_k) M_i is a subset of $F_i^{0,k}$ for each $i \in I$.

(7_{2k}) the condition ($4_{0,2k}^{\delta_{k-1},i}$) is satisfied,

Also, as we stated in the introductory part of the proof, for each m conditions ($1_{0,m}$), ($2_{0,m}$) and ($3_{0,m}$) will be satisfied and the fourth condition will be preserved in each step of the construction.

Assume that we already constructed $\mathcal{F}_{0,1}, \mathcal{F}_{0,2}, \ldots, \mathcal{F}_{0,2k}$. Let $F_i = F_i^{0,2k} \cap V_k$ and $J'_k = \{i \in I \setminus J_k \colon F_i^{0,2k} \cap V_k \neq \emptyset\}$. By star-finiteness of $\mathcal{F}_{0,2k}$, the set J'_k is finite. By lemma 9.1, there exists an adjustement of a collection $\{F_i\}_{i \in J'_k}$ to a Z-collection $\{F'_i\}_{i \in J'_k}$ that is equal to $\{M_i\}_{i \in J'_k}$ on $M \cap V_k$. Let

$$F_i^{0,2k+1} = \begin{cases} F'_i \cup (F_i^{0,2k} \setminus V_k) & i \in J'_k \\ F_i^{0,2k} & i \in I \setminus J'_k. \end{cases}$$

By lemma 4.1, if the adjustement was small enough (which we assume), then the collection $\mathcal{F}_{0,2k+1}$ is an adjustment of $\mathcal{F}_{0,2k}$ $\operatorname{rel} X \setminus V_k$. Therefore $\mathcal{F}_{0,2k+1}$ is a closed star-finite $\mathcal{N}_n$-collection. It is an interior cover because the adjustment was done with fixed J_k $\operatorname{rel} X \setminus V_k$ and the closure of V_k lies in the interior of $F_{J_k}^{0,2k}$ by the definition of U_k. By remark 10.6, if the adjustement was small enough (which we assume), then the condition ($2_{0,2k+1}$) holds. Condition ($3_{0,2k+1}$) is satisfied and the fourth condition is preserved by the construction, because the closure of V_k is disjoint from every simplex not containing δ_k and if $\delta_l \supset \delta_k$, then $J_l \subset J_k$ and the adjustement is done with fixed J_k. The condition (5_{2k+1}) is satisfied because $\mathcal{F}_{0,2k+1}$ is an adjustement $\operatorname{rel} X \setminus V_k$ of $\mathcal{F}_{0,2k}$ and $\langle K, \mathcal{K} \rangle$ is its n-core by ($2_{0,2k+1}$). The condition 6_{2k+1} follows from the construction of $\{F'_i\}_{i \in J'_k}$. Note that there is no condition (7_{2k+1}) to check.

By lemma 9.4, there is a closed (in X) $\mathcal{N}_n$-neighborhood A_k of δ_k that lies in V_k and such that for each $J''_k \subset J'_k$ (hence for each $J''_k \subset I$) the intersection $A_k \cap F_{J''_k}^{0,2k+1}$ is an $(AE(n) \cap \mathcal{N}_n)$-space. Let

$$F_i^{0,2k+2} = \begin{cases} F_i^{0,2k+1} \cup A_k & i \in J'_k \\ F_i^{0,2k+1} & i \in I \setminus J'_k. \end{cases}$$

The collection $\mathcal{F}_{0,2k+2}$ is $\mathcal{N}_n$-collection by corollary 3.8, hence ($1_{0,2k+2}$) is satisfied. By lemma 2.24, the pair $\langle K, \mathcal{K} \rangle$ is its n-core (we assume that A_k is so small that $\mathcal{F}_{0,2k+2}$

is isomorphic to $\mathcal{F}_{0,2k+1}$). Hence $(2_{0,2k+2})$ is satisfied. Conditions (3_{2k+2}) and (6_{2k+2}) are satisfied and the fourth condition is preserved, because $F_i^{2k+1} \subset F_i^{2k+2}$ for each $i \in I$. The condition (5_{2k+2}) is satisfied by $(2_{0,2k+2})$ and because A_k is a subset of V_k. By the choice of A_k, $(4_{0,2k+2}^{\delta_k,i})$ is satisfied for each $i \in I$, hence (7_{2k+2}) is satisfied.

Let $\mathcal{F}_{0,\infty} = \lim_{k\to\infty} \mathcal{F}_{0,k}$. We are going to show that $(1_{0,\infty})$, $(2_{0,\infty})$, $(3_{0,\infty})$ and, for each 0-dimensional simplex $\delta \in \tau$ and each $i \in I$, $(4_{0,\infty}^{\delta,i})$ are satisfied. Since the sequence $\mathcal{F}_{0,0}, \mathcal{F}_{0,1}, \dots$ stabilizes locally and $(1_{0,k})$, $(3_{0,k})$ and (7_{2k}) hold for each k, only verification of $(2_{0,\infty})$ needs an argument. What we have to show is that $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}_{0,\infty}$. Let $H_k \colon X \to X$ be a closed embedding defined as the identity on $X \setminus \bigcup_{l\geq k} U_l$ and by the formula $H_k(x) = h_l(x)$ for each $x \in U_l$ with $l \geq k$. Let $\mathcal{G}_k = \{G_i^k\}_{i\in I}$ and let $G_i^k = H_k(F_i^{0,2k})$. Observe that $\mathcal{G}_k$ is a closed interior $\mathcal{N}_n$-cover of $X \setminus \bigcup_{l\geq k} V_l$ and $\langle L, \mathcal{L}\rangle$ is its n-core. Also, $\mathcal{G}_k$ is equal to $\mathcal{G}_{k-1}$ on $X \setminus U_k$ and $\mathcal{G}_{k-1}$ is a shrinking of $\mathcal{G}_k$ on $U_k \setminus V_k$. By lemma 10.7, $\langle L, \mathcal{L}\rangle$ is an n-core of $\lim_{k\to\infty} \mathcal{G}_k$. But $\lim_{k\to\infty} \mathcal{G}_k = \lim_{k\to\infty} \mathcal{F}_{0,k}$, so $\langle L, \mathcal{L}\rangle$, and also $\langle K, \mathcal{K}\rangle$, is an n-core of $\mathcal{F}_{0,\infty}$.

Now we are going to describe a similar construction for simplices of dimension $l > 0$. It presents additional difficulty because simplices of positive dimensions have non-empty boundaries.

Let $0 < l \leq n$ and assume that we already constructed $\mathcal{F}_{l,0}$ and $(1_{l,0})$, $(2_{l,0})$, $(3_{l,0})$ and $(4_{l,0}^{\delta,i})$ are satisfied for each at most $(l-1)$-dimensional simplex $\delta \in \tau$ and each $i \in I$. Let $\delta_0, \delta_1, \dots$ be a sequence of all l-dimensional simplices in τ_L. For each k let δ'_k be a shrinking of δ_k, by a homothety with a ratio close to 1 and with a center equal to the barycenter of δ_k, such that the following conditions are satisfied with $m = 0$.

(8) $\{\delta'_k\}$ is a discrete collection in X,

(9_m) if $\delta_k \subset \operatorname{Int}_K K_i$, then $\operatorname{Cl}_X(\delta_k \setminus \delta'_k) \subset \operatorname{Int}_X F_i^{l,m}$.

The last condition can be satisfied because by the assumption every proper face δ of δ_k satisfies $(4_{l,0}^{\delta,i})$ for each $i \in I$.

Let $J_k = \{i \in I \colon \delta_k \subset \operatorname{Int}_X F_i^{l,0}\}$. For each k fix an open neighborhood U_k of δ'_k whose closure lies in the interior of $F_{J_k}^{l,0}$ and is disjoint from every simplex of τ that is disjoint from δ'_k. Without a loss of generality we may assume that $\{U_k\}$ is a discrete family in X and each U_k is disjoint from L. Define V_k in the same manner as in the 0-dimensional case. Let $J'_k = \{i \in I \setminus J_k \colon F_i^{l,0} \cap V_k \neq \emptyset\}$. Let $\widehat{J}_k = \{i \in J'_k \colon \delta_k \subset \operatorname{Int}_K K_i\}$. Let $m_0 = 0$ and $m_{k+1} = m_k + 4|\widehat{J}_k|$, where $|\widehat{J}_k|$ denotes the number of elements of $\widehat{J}_k$ (the set J_k is finite because $\mathcal{F}_{l,0}$ is locally finite). For each k order $\widehat{J}_k$ into a sequence $j_k^0, j_k^1, \dots, j_k^{|\widehat{J}_k|-1}$.

We are going to construct a sequence $\mathcal{F}_{l,1}, \mathcal{F}_{l,2}, \dots$ in such a way that the following conditions will be satisfied for each $k > 0$.

(10_k) $\mathcal{F}_{l,k}$ is a deformation of $\mathcal{F}_{l,k-1}$ $\operatorname{rel} X \setminus V_m$, for m such that $m_m < k \leq m_{m+1}$,

(11_k) M_i is a subset of $F_i^{l,k}$ for each $i \in I$,

($12_{k,m}$) the condition ($4^{\delta_k, j_k^m}_{l,m_k+4m+4}$) is satisfied, where $0 \leq m < |\widehat{J}_k|$.

Also, for each m, conditions ($1_{l,m}$), ($2_{l,m}$), ($3_{l,m}$) and (9_m) will be satisfied and the fourth condition will be preserved in each step of the construction.

Fix $0 \leq k < \infty$ and $0 \leq \hat{m} < |\widehat{J}_k|$ and let $m = m_k + 4\hat{m}$. Assume that we already constructed $\mathcal{F}_{l,m}$. Let j denote $j_k^{\hat{m}}$. We shall construct covers $\mathcal{F}_{0,m+1}, \ldots, \mathcal{F}_{0,m+4}$. Our goal is to meet condition ($12_{k,\hat{m}}$) and verify that other conditions specified above are satisfied. Let $\hat{\delta}_k$ be a shrinking of δ'_k by a homothety with a ratio less than 1 and a center equal to the barycenter of δ'_k, such that (9_m) is satisfied with $\hat{\delta}_k$ substituted for δ'_k. Let $\hat{J} = \{i \in J'_k \colon i \neq j_k^q, q \leq \hat{m}\}$ and let $\hat{V}_k$ be an open neighborhood of $\hat{\delta}_k$, whose closure lies in the intersection of $(V_k \setminus \delta_k) \cup (\delta'_k \setminus \partial\delta'_k)$ with $\operatorname{Int}_X F^{l,m}_{J_k \cup \{j_k^q \colon q < \hat{m}\}}$. Let m' be an integer greater than or equal to m and less than $m+4$. We shall construct $\mathcal{F}_{l,m'+1}$ as a collection equal to $\mathcal{F}_{l,m'}$ on $\operatorname{rel} X \setminus \hat{V}_k$ and such that $F_i^{l,m'+1} = F_i^{l,m'}$ for each $i \in J_k \cup \{j_k^q \colon q < \hat{m}\}$. This way, the condition ($9_{m'+1}$) will be satisfied and the condition ($10_{m'+1}$) will follow from ($2_{l,m'+1}$).

Let $P = \hat{V}_k \cap \operatorname{Int}_X F_j^{l,m}$. Consider a collection $\{F_i\}_{i \in \hat{J}}$, where $F_i = F_i^{l,m} \cap P$. By lemma 9.1, it can be adjusted to a Z-collection $\{F'_i\}_{i \in \hat{J}}$ that is equal to $\{M_i\}_{i \in \hat{J}}$ on $M \cap P$. Let

$$F_i^{l,m+1} = \begin{cases} F'_i \cup (F_i^{l,m} \setminus P) & i \in \hat{J} \\ F_i^{l,m} & i \in I \setminus \hat{J}. \end{cases}$$

As before, if we assume that the adjustement is small enough, then conditions ($1_{l,m+1}$), ($2_{l,m+1}$), ($3_{l,m+1}$) and (11_{m+1}) are satisfied and the fourth condition is preserved. By our earlier remark, (9_{m+1}) and (10_{m+1}) are satisfied as well.

Let V'_k be an open subset of $\hat{V}_k$ such that $\operatorname{Cl}_X V'_k \cap \operatorname{Cl}_X(\delta_k \setminus \hat{\delta}_k) = \emptyset$ and $V'_k \cup P = \hat{V}_k$. Consider a collection $\{F_i\}_{i \in \hat{J} \cup \{j\}}$ such that $F_i = F^{l,m+1} \cap V'_k$. By lemma 9.1, it can be adjusted to a Z-collection $\{F'_i\}_{i \in \hat{J} \cup \{j\}}$ that is equal to $\{M_i\}_{i \in \hat{J} \cup \{j\}}$ on $M \cap V'_k$. Let

$$F_i^{l,m+2} = \begin{cases} F'_i \cup (F_i^{l,m+1} \setminus V'_k) & i \in \hat{J} \cup \{j\} \\ F_i^{l,m+1} & i \in I \setminus \hat{J}, i \neq j. \end{cases}$$

Again, if the adjustement is small enough, then the collection $\mathcal{F}_{l,m+2}$ satisfies all prescribed conditions. We claim that the following conditions are satisfied as well.

(13) $\{F_i^{l,m+2} \cap \hat{V}_k\}_{i \in \hat{J}}$ is an $\mathcal{N}_n$-collection and a Z-collection in $\hat{V}_k$,

(14) $\{F_i^{l,m+2} \cap \hat{V}_k \cap F_j^{l,m+2}\}_{i \in \hat{J}}$ is an $\mathcal{N}_n$-collection and a Z-collection in $\hat{V}_k \cap F_j^{l,m+2}$,

(15) $\operatorname{Cl}_X(\delta_k \setminus \hat{\delta}_k) \subset \operatorname{Int}_X F_j^{l,m+2}$.

By the construction, $\{F_i^{l,m+2}\}_{i \in \hat{J}}$ is equal to $\mathcal{M}$ on $\hat{V}_k \cap M$, so $\hat{\delta}_k$ fits $\{F_i^{l,m+2}\}_{i \in \hat{J}}$ and by (13) $\{F_i^{l,m+2} \cap \hat{V}_k\}_{i \in \hat{J}}$ is a Z- and $\mathcal{N}_n$-collection in $\hat{V}_k$. Hence, by lemma 9.4, there is a closed (in X) $\mathcal{N}_n$-subset A_k of $\hat{V}_k$ that fits $\{F_i^{l,n+2} \cap \hat{V}_k\}_{i \in \hat{J}}$ and such that $\hat{\delta}_k \subset \operatorname{Int}_X A_k$ and $\{F_i^{l,m+2}\}_{i \in \hat{J}}$ restricted to A_k is both a Z- and an $\mathcal{N}_n$-collection.

Observe that by the choice of $\hat{V}_k$ and $\hat{J}$, A_k fits $\{F_i^{l,m+2} \cap \hat{V}_k\}_{i \in I, i \neq j}$. By lemma 9.2, we may adjust A_k to a closed (in X) $\mathcal{N}_n$-subset A'_k of $\hat{V}_k$ such that $A'_k \cap \delta_k = \hat{\delta}_k$, $\hat{\delta}_k \setminus \partial\hat{\delta}_k \subset \operatorname{Int}_X A'_k$ and $\{F_i^{l,m+2}\}_{i \in \hat{J}}$ restricted to A'_k is both a Z- and an $\mathcal{N}_n$-collection. Let $W_k = (F_j^{l,m+2} \cap \operatorname{Int}_X A'_k) \cup (A'_k \cap \operatorname{Int}_X F_j^{l,m+2})$. It is an open neighborhood of $\hat{\delta}_k$ in $A'_k \cap F_j^{l,m+2}$ and it is an $\mathcal{N}_n$-space. By (14) and by lemma 9.4, there is a closed $\mathcal{N}_n$-subset B_k of W_k that fits $\{F_i^{l,m+2} \cap W_k\}_{i \in I}$ and such that $\hat{\delta}_k \subset \operatorname{Int}_{W_k} B_k$. If it is small enough, then it is closed in X. Let W'_k be an open subset of V_k such that $W'_k \cap (A'_k \cap F_j^{l,m+2}) = (A'_k \cap F_j^{l,m+2}) \setminus B_k$ and $\operatorname{Cl}_X W'_k \cap \delta_k = \emptyset$. By lemma 9.2, we may adjust A'_k and $F_j^{l,m+2}$ rel $X \setminus W'_k$ to obtain sets A''_k and $\widetilde{F}_j^{l,m+2}$ such that $\mathcal{F}_{l,m+3}$ defined in the following way

$$F_i^{l,m+3} = \begin{cases} \widetilde{F}_j^{l,m+2} & i = j \\ F_i^{l,m+2} & i \in I \text{ and } i \neq j. \end{cases}$$

is an adjustment of $\mathcal{F}_{l,m+2}$. Observe that A''_k fits $\mathcal{F}_{l,m+3}$, because its intersection with $\widetilde{F}_j^{l,m+2}$ is equal to B_k. Observe that $\hat{\delta}_k \setminus \partial\hat{\delta}_k \subset \operatorname{Int}_X A''_k$ and the restriction of $\mathcal{F}_{l,m+3}$ to A''_k is an $\mathcal{N}_n$-cover. By the construction, $\operatorname{Cl}_X \delta_k \setminus \hat{\delta}_k \subset \operatorname{Int} A''_k$.

Let

$$F_i^{l,m+4} = \begin{cases} F_i^{l,m+3} \cup A''_k & i \in J'_k \\ F_i^{l,m+3} & i \in I \setminus J'_k. \end{cases}$$

Since $\delta_k \subset \operatorname{Int}_X F_j^{l,m+4} \cup \operatorname{Int}_X A''_k$, $(12_{k,\hat{m}})$ is satisfied. Verification of other conditions is similar to the verification done in the previous steps.

The limit argument is similar to the 0-dimensional case. Let $\mathcal{F}_{l,\infty} = \lim_{k \to \infty} \mathcal{F}_{l,k}$. We are going to show that $(1_{l,\infty})$, $(2_{l,\infty})$, $(3_{l,\infty})$ and, for each at most l-dimensional simplex $\delta \in \tau$ and each $i \in I$, $(4_{l,\infty}^{\delta,i})$ are satisfied. Since the sequence $\mathcal{F}_{l,0}, \mathcal{F}_{l,1}, \ldots$ stabilizes locally and $(1_{l,k})$, $(2_{l,k})$ and $(12_{k,m})$ are satsified for all k and m, all we have to show is that $\langle K, \mathcal{K} \rangle$ is an n-core of $\mathcal{F}_{l,\infty}$. Let $H_k \colon X \to X$ be a closed embedding defined as the identity on $X \setminus \bigcup_{l \geq k} U_l$ and by the formula $H_k(x) = h_m(x)$ for each $x \in U_m$ with $m \geq k$. Let $\mathcal{G}_k = \{G_i^k\}_{i \in I}$ and let $G_i^k = H_k(F_i^{l,m_{k+1}})$. Observe that $\mathcal{G}_k$ is a closed interior $\mathcal{N}_n$-cover of $X \setminus \bigcup_{m \geq k} V_m$ and $\langle L, \mathcal{L} \rangle$ is its n-core. Also, $\mathcal{G}_k$ is equal to $\mathcal{G}_{k-1}$ on $X \setminus U_k$ and $\mathcal{G}_{k-1}$ is a shrinking of $\mathcal{G}_k$ on $U_k \setminus V_k$. By lemma 10.7, $\langle L, \mathcal{L} \rangle$ is an n-core of $\lim_{k \to \infty} \mathcal{G}_k$. But $\lim_{k \to \infty} \mathcal{G}_k = \lim_{k \to \infty} \mathcal{F}_{l,k}$, so $\langle L, \mathcal{L} \rangle$, and also $\langle K, \mathcal{K} \rangle$, is an n-core of $\mathcal{F}_{l,\infty}$.

Second part.

In the first part we constructed a closed interior $\mathcal{N}_n$-cover $\mathcal{F}_{n,\infty}$ of X that satisfies conditions $(1_{n,\infty})$, $(2_{n,\infty})$, $(3_{n,\infty})$ and $(4_{n,\infty}^{\delta,i})$ for each $\delta \in \tau$ and $i \in I$. By the construction, $\mathcal{F}_{n,\infty}$ is equal to $\mathcal{F}_{0,0}$ on the complement of an open neighborhood U_L of L. Let $\mathcal{U}$ be an open cover of X satisfying conditions of remark 10.6. If $\mathcal{F}_{n+1,\infty}$ is a $\mathcal{U}$-adjustment rel $X \setminus U_L$ of $\mathcal{F}_{n,\infty}$ to a closed interior $\mathcal{N}_n$-cover such that $K_i = K \cap F_i^{n+1,\infty}$

for each $i \in I$, then, by the choice of $\mathcal{U}$, $\langle K, \mathcal{K} \rangle$ is an exact n-core of $\mathcal{F}_{n+1,\infty}$. We shall construct such $\mathcal{F}_{n+1,\infty}$, finishing the proof.

Order τ_L into a sequence $\{\delta_k\}_{k\geq 1}$ non decreasing in the order by inclusion. Let $J_k = \{i \in I \colon \delta_k \subset \operatorname{Int}_K K_i\}$. By the choice of τ made at the beginning of the proof, J_k is non-empty for each $k \geq 1$. Since $\mathcal{F}_{n,\infty}$ satisfies $(4^{\delta_k;i}_{n,\infty})$ for each $k \geq 1$ and each $i \in I$, the inclusion $\delta_k \subset \operatorname{Int}_X F^{n,\infty}_{J_k}$ holds for each k (this is the crucial property obtained in the first part of the proof). Therefore there exists a locally finite collection $\{V_k\}_{k\geq 1}$ of open subsets of U_L such that $L \subset \bigcup_{k\geq 1} V_k$ and the following conditions are satisfied for each k.

(16) $K \cap V_k \subset K_{J_k}$,

(17) if $\operatorname{Cl}_X V_l$ intersects $\operatorname{Cl}_X V_k$, then either δ_l is a face of δ_k or δ_k is a face of δ_l,

(18) $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n,\infty}_{J_k}$.

Let $\mathcal{F}_{n+1,0} = \mathcal{F}_{n,\infty}$. Let $\mathcal{U}_k$ be a sequence of open covers of X obtained via lemma 4.3 applied to a locally finite collection $\{V_k\}_{k\geq 1}$ and an open cover $\mathcal{U}$. We recursively define a sequence $\mathcal{F}_{n+1,k}$ of closed interior $\mathcal{N}_n$-covers of X such that the following conditions are satisfied for each $k \geq 1$.

(19) $\mathcal{F}_{n+1,k}$ is a $\mathcal{U}_k$-adjustment of $\mathcal{F}_{n+1,k-1}$ $\operatorname{rel} X \setminus V_k$ with fixed J_k,

(20) $\mathcal{F}_{n+1,k}$ is equal to $\mathcal{K}$ on $K \cap \bigcup_{l\leq k} V_l$.

Fix $k \geq 1$ and assume that we already constructed $\mathcal{F}_{n+1,k-1}$. First, we claim that the inclusion $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,l}_{J_k}$ holds for each $0 \leq l \leq k-1$. The proof is by induction on l. The basis step of induction ($l = 0$) follows directly from (18) and the identity $\mathcal{F}_{n+1,0} = \mathcal{F}_{n,\infty}$. Fix $l > 0$ and assume that we already proved that $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,l-1}_{J_k}$. By (19), $\mathcal{F}_{n+1,l}$ is an adjustment of $\mathcal{F}_{n,l-1}$ $\operatorname{rel} X \setminus V_l$, hence if V_l is disjoint from $Cl_X V_k$, then $\operatorname{Cl}_X V_k \cap \operatorname{Int}_X F^{n+1,l-1}_{J_k} = \operatorname{Cl}_X V_k \cap \operatorname{Int}_X F^{n+1,l}_{J_k}$, hence $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,l}_{J_k}$. If $\operatorname{Cl}_X V_l$ intersects V_k, then by (17) and by the assumption that the sequence $\{\delta_k\}$ is non-decreasing in the order by inclusion, δ_l is a face of δ_k. In this case, J_k is a subset of J_l by the definition. By (19), $\mathcal{F}_{n+1,l}$ is an adjustment of $\mathcal{F}_{n,l}$ with fixed J_l, hence $\operatorname{Int}_X F^{n+1,l}_{J_k} = \operatorname{Int}_X F^{n+1,l-1}_{J_k}$ and the inclusion $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,l}_{J_k}$ follows from the inductive assumption. The inductive step is done. In particular, we have proven that $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,k-1}_{J_k}$.

By lemma 9.1, there exists an adjustment of a collection $\{F^{n+1,k-1}_i \cap V_k\}_{i\in I\setminus J_k}$ of subsets of V_k to a Z-collection that is equal to $\{K_i \cap V_k\}_{i\in I\setminus J_k}$ on $K \cap V_k$. By lemma 4.1, we may require (and we do) that the adjustment is so small, that it extends to an adjustment $\operatorname{rel} X \setminus V_k$ of $\{F^{n+1,k-1}_i\}_{i\in I\setminus J_k}$ to a closed $\mathcal{N}_n$-collection of subsets of X. We let $\mathcal{F}_{n+1,k}$ be a union of this collection with the collection $\{F^{n+1,k-1}\}_{i\in J_k}$. Observe that $\mathcal{F}_{n+1,k}$ is a $\mathcal{U}_k$-adjustment of $\mathcal{F}_{n+1,k-1}$ $\operatorname{rel} X \setminus V_k$ with fixed J_k directly from the construction. It is a closed interior $\mathcal{N}_n$-cover of X because $\operatorname{Cl}_X V_k \subset \operatorname{Int}_X F^{n+1,k}_{J_k}$.

Hence (19) is satisfied. To see that $\mathcal{F}_{n+1,k}$ satisfies condition (20) recall that by (16), $K \cap V_k \subset K_{J_k}$, hence $F_i^{n+1,k} \cap K \cap V_k = K_i \cap V_k$ for each $i \in J_k$. If $i \in I \setminus J_k$, then $F_i^{n+1,k} \cap K \cap V_k = K_i \cap V_k$ directly from the construction.

Let $\mathcal{F}_{n+1,\infty} = \lim_{k\to\infty} \mathcal{F}_{n+1,k}$. By the choice of $\mathcal{U}_k$'s, it is a $\mathcal{U}$-adjustment of $\mathcal{F}_{n+1,0}$. Since the collection $\{V_k\}$ is locally finite, $\mathcal{F}_{n+1,\infty}$ is a closed interior $\mathcal{N}_n$-cover. By the choice of $\mathcal{U}$, $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}_{n+1,\infty}$. By (20), by the assumption that $L \subset \bigcup_{k\geq 1} V_k$ and by the assumption that $\mathcal{F}_{n+1,0}$ is equal to $\mathcal{K}$ on $K \setminus L$, $\langle K, \mathcal{K}\rangle$ is an exact n-core of $\mathcal{F}_{n+1,\infty}$. The cover $\mathcal{F}_{n+1,\infty}$ is equal to $\mathcal{F}_{0,0}$ on the complement of U_L directly from the construction. We are done.

10.4 Retraction onto a core and a proof of theorem 6.17

We shall now obtain theorem 6.17 as an easy corollary of the results obtained earlier in this chapter.

Lemma 10.9. *If K is an exact n-core of a closed star-finite locally finite $ANE(n)$-cover $\mathcal{F}$ of an at most n-dimensional space X, then there exists a retraction of X onto K that is $\mathcal{F}$-close to the identity.*

Proof. Let $\mathcal{F} = \{F_i\}_{i\in I}$ and for each non-empty $J \subset I$ let F_J denote, as usual, the intersection $\bigcap_{i\in J} F_i$. Let J_1, J_2 be subsets of I. Assume that F_{J_2} is non-empty. We will show that the inclusion of $F_{J_2} \cap (K \cup \bigcup_{i\in J_1} F_i)$ into F_{J_2} is a weak n-homotopy equivalence. Without a loss of generality, we may assume that J_1 is disjoint from J_2 and that all elements indexed by J_1 intersect F_{J_2}. By the definition of an exact n-core, for each $i \in I$ the intersection of F_i with K is a subcomplex of K. By the assumption that $\mathcal{F}$ is an $ANE(n)$-cover, for each non-empty $J \subset I$ the intersection F_J is an $ANE(n)$-space. By the assumption, $\mathcal{F}$ is closed and locally finite, hence theorem 2.7 implies that $F_{J_2} \cap (K \cup \bigcup_{i\in J_1} F_i)$ is an $ANE(n)$-space for each J_1 and J_2. Fix J_2 such that F_{J_2} is non-empty and assume that the assertion is already proven for all $J \subset I$ such that $J_2 \subsetneq J$ and $F_J \neq \emptyset$ (there are only finitely many of such J's as $\mathcal{F}$ is star-finite). Let J_1 be a subset of I that is disjoint from J_2 and is such that elements indexed by J_1 intersect F_{J_2}. Again, there are only finitely many of those. We shall prove the assertion by an induction over the cardinality of J_1. By the assumption that K is an exact n-core of $\mathcal{F}$, the inclusion of $K \cap F_{J_2}$ into F_{J_2} is an n-homotopy equivalence. By Whitehead's characterization, this inclusion is a weak n-homotopy equivalence. Hence the assertion is verified for $J_1 = \emptyset$. Let $J_1 \neq \emptyset$. Fix $j \in J_1$. We assumed that J_1 is disjoint from J_2, so $j \notin J_2$. Let $A_1 = F_{J_2} \cap (K \cup \bigcup_{i\in J_1, i\neq j} F_i)$ and $A_2 = F_{J_2} \cap F_j$. As we argued earlier, both A_1 and A_2 are $ANE(n)$-spaces. By the assumption, X is at most n-dimensional, and so are A_1 and A_2. By the inductive assumptions, the inclusions $A_1 \cap A_2 \subset A_2$ and

$A_1 \subset F_{J_2}$ are weak n-homotopy equivalences. Hence by corollary 2.25, the inclusion $A_1 \cup A_2 \subset F_{J_2}$ is a weak n-homotopy equivalence. This is the assertion that we wanted to prove.

By theorem 2.23, for each $J \subset I$ such that F_J is non-empty there exists a retraction $r_J \colon F_J \to F_J \cap (K \cup \bigcup_{i \in I \setminus J} F_i)$. Let $\mathcal{I} = \{J \subset I \colon F_J \neq \emptyset\}$. Order $\mathcal{I}$ into a sequence $\{J_k\}_{k \geq 1}$ non decreasing in the order by inclusion. Such ordering exists because $\mathcal{F}$ is point finite. Let r_0 be the identity on X. Let $r_k \colon \bigcup_{i \in I \setminus J_k} F_i \cup K \cup F_{J_k} \to \bigcup_{i \in I \setminus J_k} F_i \cup K$ be a map defined by the following equation.

$$r_k = \begin{cases} r_{J_k}(x) & x \in F_J \\ x & x \in K \cup \bigcup_{i \in I \setminus J_k} F_i. \end{cases}$$

Obviously each r_k is a contiuous retraction and $r_k(F_i) \subset F_i$ for each $i \in I$. By the order of J_k's for each x in X the sequence of compositions $r_1, r_2 \circ r_1, \ldots$ is well defined. By star-finiteness of $\mathcal{F}$ it stabilizes for each x after finitely many steps. Hence the limit

$$r(x) = \lim_{k \to \infty} r_k \circ \cdots \circ r_2 \circ r_1(x).$$

is well defined for each x in X. It is a retraction that we were looking for. □

Proof of theorem 6.17. Let $\mathcal{U}$ be an open cover of an $\mathcal{N}_n$-space X. By [18, theorem 5.3.10] and the fact that every separable metric space is strongly paracompact, there exists a closed star-finite locally finite cover of X that refines $\mathcal{U}$. By theorem 4.5, it has a closed star-finite swelling to an interior $\mathcal{N}_n$-cover $\mathcal{F}$ refining $\mathcal{U}$. By theorem 10.5, $\mathcal{F}$ has an n-core $\langle K, \mathcal{K} \rangle$. By theorem 10.1, $\mathcal{F}$ can be deformed to a closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{G}$, which refines $\mathcal{U}$ and has an exact n-core K. By lemma 10.9, there is a retraction $r \colon X \to K$ that is $\mathcal{G}$-close to the identity. It satisfies assertion of the theorem because $\mathcal{G}$ refines $\mathcal{U}$. □

Chapter 11

Proof of theorem 6.7

The entire chapter is devoted to the proof of theorem 6.7. Let us recall the statement.

Theorem 6.7. *There exists a constant N such that if a k-regular n-semiregular ($k < n$) closed countable star-finite interior $\mathcal{N}_n$-cover $\mathcal{F}$ is n-contractible in a closed cover $\mathcal{E}$, then there exists a $(k+1)$-regular n-semiregular interior $\mathcal{N}_n$-cover isomorphic to $\mathcal{F}$ and refining $\operatorname{st}^N \mathcal{E}$. Moreover, we may additionally require that the constructed cover is equal to $\mathcal{F}$ on some open neighborhood of a given $Z(\mathcal{F})$-set Z.*

We shall prove that the assertion is true with $N = 472$, but the actual value is of no importance and no effort was made to minimize it.

11.1 Patching of small holes

We fix n and $k < n$. Let $\mathcal{F} = \{F_i\}_{i \in I}$ be a cover of a space X. We say that a map φ from a k-dimensional sphere S^k into X is *a k-hole in $\mathcal{F}$* if the image of φ lies in $F_J = \bigcap_{i \in J} F_i$ for some non-empty subset J of I. When we will want to specify J, we will say that φ is *a k-hole in F_J*. If φ is a k-hole in F_J and is null-homotopic in F_J, then we say that it is a *trivial k-hole*. Observe that if $\mathcal{F}$ is k-regular and all k-holes in $\mathcal{F}$ are trivial, then it is $(k+1)$-regular. We shall describe a construction that enables us to *patch* holes. That is, to make them trivial in a newly constructed cover. We begin with a special case. If φ is a k-hole in F_J and φ is contractible in F_j for some $j \in J$, then we say that φ is a *small k-hole*. We describe a construction of patching of a small k-hole.

Definition 11.1. Let $\langle K, \mathcal{K} \rangle$ be a pair consisting of a locally finite countable simplicial complex K and a cover $\mathcal{K} = \{K_i\}_{i \in I}$ of K by subcomplexes. Let $k < n$ and let B be a $(k+1)$-dimensional ball that is embedded as a subcomplex of K. We will say that a pair $\langle L, \mathcal{L} \rangle$ is *obtained from $\langle K, \mathcal{K} \rangle$ by patching of B*, if the following conditions are satisfied.

(1) (a) L is a locally finite countable simplicial complex and K is a subcomplex of L,

   (b) $\mathcal{L} = \{L_i\}_{i \in I}$ is a cover of L by subcomplexes and $\mathcal{L}$ is equal to $\mathcal{K}$ on K.

(2) There exists a complex

$$C = \begin{cases} B \times [0,1], & \text{if } k < n-1 \\ B \times \{0,1\} \cup \partial B \times [0,1], & \text{if } k = n-1 \end{cases}$$

such that $L = K \cup C$ and the intersection of K with C is equal to B (in K) and to $B \times \{0\}$ (in C). We identify B with $B \times \{0\}$.

(3) Recall that $\mathcal{K} = \{K_i\}_{i \in I}$ and $\mathcal{L} = \{L_i\}_{i \in I}$. For each $i \in I$, L_i is the smallest subset of L for which the following conditions are satisfied.

   (a) if $B \subset K_i$, then $C \subset L_i$.

   (b) if $\partial B \subset K_i$, then $\partial B \times [0,1] \cup B \times \{1\} \subset L_i$.

   (c) $\begin{array}{l} \text{if } k < n-1, \text{then } (B \cap K_i) \times [0,1/2] \subset L_i \\ \text{if } k = n-1, \text{then } (\partial B \cap K_i) \times [0,1/2] \subset L_i \end{array}$.

Note that these conditions uniquely determine L_i's.



Figure 11.1: A picture discussed in example 11.2

Example 11.2. Let K be a square shown on figure 11.1(a) and let B be the white disc embedded as a subcomplex of K. We let $K_1 = B$, K_2 to be the dark disc and K_3 to be the complement of the interior of K_1. Let $\mathcal{K} = \{K_1, K_2, K_3\}$. Figure 11.1(b) shows a pair $\langle L, \mathcal{L} \rangle$ obtained from $\langle K, \mathcal{K} \rangle$ by patching of B. The cylinder $C = B \times [0,1]$ is attached to K along its lower base (in the example we assume that $k < n-1$) to form $L = K \cup C$. The set L_1 is equal to C. The set L_2 is an union of K_2 with a cylinder of height 1/2 and base $K_1 \cap K_2$. The set L_3 is an union of K_3 with $\partial B \times [0,1] \cup B \times \{1\}$.

The cover $\mathcal{K}$ of K is 1-regular, but not 2-regular as both K_3 and $K_1 \cap K_3$ are not simply connected. Observe that $\partial B \subset K_1 \cap K_3$ is a small hole in $K_1 \cap K_3$, as it is null-homotopic in K_1. Observe that $\partial B \subset L_1 \cap L_3$ is null-homotopic in $L_1 \cap L_3$. The small hole $\partial B \subset K_1 \cap K_3$ in $\mathcal{K}$ is "patched" in $\mathcal{L}$ and the cover $\mathcal{L}$ is 2-regular.

Remark 11.3. Assume the notation of definition 11.1. By (1b), $\mathcal{L}$ is a cover of L and by (3), $\mathcal{L}$ covers L if and only if $B \subset K_i$ for some $i \in I$. It follows from (3c) that if $\mathcal{K}$ is an interior cover, then $\mathcal{L}$ is an interior cover as well. This is the only reason for which (3c) is required. By the conditions enforced on $\mathcal{L}$, $\mathcal{L}$ is always isomorphic to $\mathcal{K}$.

Definition 11.4. Let $\mathcal{F} = \{F_i\}_{i \in I}$ be a closed star-finite interior $\mathcal{N}_n$-cover of a space X. Let K be an exact n-core of $\mathcal{F}$. Let B be a subcomplex of K that is homeomorphic to a $(k+1)$-dimensional ball and such that $B \subset F_i$ for some $i \in I$. Let U be an open neighborhood of B in X. We say that a closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{G} = \{G_i\}_{i \in I}$ of X is *obtained from $\mathcal{F}$ by patching of B in U*, if the conditions (0)-(3) stated below are satisfied.

(0) (a) $\mathcal{G}$ is isomorphic to $\mathcal{F}$ and is equal to $\mathcal{F}$ on $K \cup (X \setminus U)$,
  (b) there exists an exact n-core L of $\mathcal{G}$ such that $L \setminus K \subset U$.

Recall that $\mathcal{F}/K = \{F_i \cap K\}_{i \in I}$ and $\mathcal{G}/L = \{G_i \cap L\}_{i \in I}$ (definition 2.1). We require that the pair $\langle L, \mathcal{G}/L \rangle$ is obtained from $\langle K, \mathcal{F}/K \rangle$ by patching of B, i.e. that

(1) (a) L is a locally finite countable simplicial complex and K is a subcomplex of L,
  (b) $\mathcal{G}/L$ is a cover of L by subcomplexes and $\mathcal{F}$ is equal to $\mathcal{G}$ on K.

(2) There exists a complex

$$C = \begin{cases} B \times [0,1], & \text{if } k < n-1 \\ B \times \{0,1\} \cup \partial B \times [0,1], & \text{if } k = n-1 \end{cases}$$

such that $L = K \cup C$ and the intersection of K with C is equal to B (in K) and to $B \times \{0\}$ (in C). We identify B and $B \times \{0\}$.

(3) For each $i \in I$, $G_i \cap L$ is the smallest subset of L for which the following conditions are satisfied.

  (a) if $B \subset F_i$, then $C \subset G_i$.
  (b) if $\partial B \subset F_i$, then $\partial B \times [0,1] \cup B \times \{1\} \subset G_i$.
  (c) $\begin{array}{l} \text{if } k < n-1, \text{then } (B \cap F_i) \times [0,1/2] \subset G_i \\ \text{if } k = n-1, \text{then } (\partial B \cap F_i) \times [0,1/2] \subset G_i \end{array}$.

Note that these conditions uniquely determine $L \cap G_i$'s.

Remark 11.5. Assume the notation of definition 11.4. Let $J \subset I$ and assume that $B \subset F_i$ for some $i \in J$. If $\partial B \subset F_J$ is a small k-hole in F_J, then $\partial B \subset G_J$ is a trivial k-hole in G_J. There are no "new" non-trivial k-holes in $\mathcal{G}$ (cf. example 11.2; $\langle L, \mathcal{G}/L\rangle$ is an exact n-core of $\mathcal{G}$).

Lemma 11.6. *If $\mathcal{F}$ is a closed star-finite interior $\mathcal{N}_n$-cover of a space X, K is an exact n-core of $\mathcal{F}$, B is a $(k+1)$-dimensional ball embedded as a subcomplex of K, B is contained in some element of $\mathcal{F}$ and U is a neighborhood of B, then there exists a closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{G}$ of X that is obtained from $\mathcal{F}$ by patching of B in U.*

Proof. Let $\mathcal{F} = \{F_i\}_{i\in I}$. The construction of $\mathcal{G}$ proceeds in two stages.
Cleaning up. In the first stage we construct a closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{H} = \{H_i\}_{i\in I}$ of X and its exact n-core M that satisfy conditions of definition 11.4 (with H_i substituted for G_i and M substituted for L), with the exception that condition (3b) is replaced by condition

(3b') if $\partial B \subset F_i$, then $\partial B \times [0,1] \subset H_i$.

Let
$$C_0 = \begin{cases} B \times [0,1], & \text{if } k < n-1, \\ B \times \{0,1\} \cup \partial B \times [0,1], & \text{if } k = n-1 \end{cases}.$$

Identify $B \times \{0\}$ with B and let p be the projection of C_0 onto B along $[0,1]$. Let $\imath$ be the inclusion of K into X. The map $p \cup \imath$ is a well-defined map into X, and its domain is an most n-dimensional countable simplicial complex. By theorem 3.6 and by remark 3.5, we can approximate $p \cup \imath \operatorname{rel} K$ within $\mathcal{F}$ by a closed embedding f. Take an approximation close enough, such that the image $f(C_0)$ lies in U and let $C = f(C_0)$. Observe that $C \cap K = B$. Let $\mathcal{K} = \{K_i\}_{i\in I}$ be a cover of $K \cup C$ by subcomplexes, such that $\mathcal{K}$ is equal to $\mathcal{F}$ on K and such that for each $i \in I$, K_i is a smallest subset of K for which the following conditions are satisfied.

(a) if $B \subset F_i$, then $C \subset K_i$.

(b') if $\partial B \subset F_i$, then $\partial B \times [0,1] \subset K_i$.

(c) $\begin{array}{l} \text{if } k < n-1, \text{then } (B \cap F_i) \times [0,1/2] \subset K_i \\ \text{if } k = n-1, \text{then } (\partial B \cap F_i) \times [0,1/2] \subset K_i \end{array}$.

Observe that $\mathcal{K}$ refines $\mathcal{F}$ because f was an approximation within $\mathcal{F}$. By the assumption K is an exact n-core of $\mathcal{F}$, so $\langle K \cup C, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}$. By theorem 10.1 applied to $\langle K \cup C, \mathcal{K}\rangle$ and $\mathcal{F}$ there is a deformation of $\mathcal{F} \operatorname{rel} X \setminus U$ to an interior $\mathcal{N}_n$-cover $\mathcal{H}$ with an exact n-core $\langle K \cup C, \mathcal{K}\rangle$. By theorem 10.1, we may require (and we do), that if $B \times \{1\}$ is a subset of K_i, then it is a subset of $\operatorname{Int}_X H_i$. The cover $\mathcal{H}$

satisfies conditions of definition 11.4 (with H_i substituted for G_i and M substituted for L), with the exception that condition (3b) is replaced by condition (3b').

Patching. Let $\Delta = B \times \{1\}$ be the upper base of C. Let $J = \{i \in I \colon B \subset F_i\}$ and $J_\partial = \{i \in I \colon \partial B \subset F_i\}$. By the construction, Δ is a subset of the interior of H_J and is disjoint from H_i for each $i \in I \setminus J_\partial$. Hence there exists an open neighborhood V of Δ that is disjoint from $\bigcup_{i \in I \setminus J_\partial} H_i$ and whose closure lies in the intersection of U with the interior of H_J. By theorem 2.35 and by lemma 4.1 there exists an approximation $\operatorname{rel} C$ of the inclusion $V \cap \bigcup_{i \in J_\partial \setminus J} H_i \subset V$ by a Z-embedding h into V such that $id_{X \setminus V} \cup h$ is continuous. Let $\mathcal{H}' = \{H'_i\}_{i \in I}$, where

$$H'_i = \begin{cases} H_i & \text{if } i \in J \\ (H_i \setminus V) \cup h(V \cap H_i) & \text{if } i \in I \setminus J. \end{cases}$$

Observe that $\mathcal{H}'$ satisfies conditions (0) - (3ab'c) with H'_i substituted for G_i and $V \cap \bigcup_{i \in I \setminus J} H'_i$ is a Z-set in V.

By lemma 9.5 applied to Δ, a Z-set $\tilde{B} = H'_{J_\partial} \cap V$ and a space V, there exists a closed $\mathcal{N}_n$-subset A of V such that $A \cap H'_{J_\partial}$ is an $\mathcal{N}_n$-space, $\Delta \subset A$ and the inclusions $\partial\Delta \subset A \cap \tilde{B}$, $\Delta \subset A$ are n-homotopy equivalences. We may assume that A is so small that it is closed in X. By corollary 2.37, the inclusion $A \cup H'_{J_\partial} \subset V$ can be approximated $\operatorname{rel} \Delta \cup H'_{J_\partial}$ by a closed embedding with image disjoint from $D = (\bigcup_{i \in I \setminus J} H'_i \cup C) \setminus (\Delta \cup H'_{J_\partial})$. Let A' denote the image of A under this embedding.

Let $\mathcal{G} = \{G_i\}_{i \in I}$, with $G_i = H'_i$ for $i \in I \setminus J_\partial$ and $G_i = H'_i \cup A'$ for $i \in J_\partial$. By the choice of V, $A' \subset H'_i$ for each $i \in J$ and by the construction $H'_i \cap A' = H'_J \cap A$ for each $i \in J_\partial \setminus J$ and the latter intersection is an $\mathcal{N}_n$-space, so $\mathcal{G}$ is a closed interior $\mathcal{N}_n$-cover by corollary 3.8. By the choice of V condition (0a) is satisfied. Let $L = K \cup C$. Since $A' \cap L = \Delta$, conditions (1) - (3abc) are satisfied. What is left is to show that L is an exact n-core of $\mathcal{G}$, i.e. that (0b) holds.

Consider a set $J' \subset I$. We have to show that $L \cap G_{J'} \subset G_{J'}$ is an n-homotopy equivalence. If $J' \not\subset J_\partial$, then $G_{J'} = H'_{J'}$ and $L \cap G_{J'} = L \cap H'_{J'}$. By the construction L is an exact n-core of $\mathcal{H}'$, so we are done. If $J' \subset J_\partial$ then $L \cap G_{J'} = (L \cap H'_{J'}) \cup \Delta$ and $G_{J'} = H'_{J'} \cup A'$. We shall show that inclusions $(L \cap H'_{J'}) \cup \Delta \subset H'_{J'} \cup \Delta$ and $H'_{J'} \cup \Delta \subset H'_{J'} \cup A'$ are n-homotopy equivalences. The latter inclusion is an n-homotopy equivalence by remark 9.6. The former inclusion is an n-homotopy equivalence by lemma 2.24 applied with $A_1 = (L \cap H'_{J'}) \cup \Delta$ and $A_2 = H'_{J'}$. We are done. □

11.2 ◎-Contractibility

Throughout the section we fix a positive integer n and a nonnegative integer $k < n$. We let $B^{k+1} \subset \mathbb{R}^{k+1}$ be the $(k+1)$-dimensional unit ball.

Definition. For each positive integer l and each positive $m \le l$ we let A^m_l denote the annulus $\{x \in B^{k+1} \colon \frac{m-1}{l} \le |x| \le \frac{m}{l}\}$.

Definition 11.7. Let $\mathcal{F}$ and $\mathcal{G}$ be covers of a space X. Let $\varphi\colon S^k \to X$ be a k-hole in $\mathcal{F}$. We say that φ is *$\circledcirc_{\mathcal{F}}$-contractible in $\mathcal{G}$* (pronounced "o"-$\mathcal{F}$-contractible), if it extends to a map $\varPhi$ from B^{k+1} into an element of $\mathcal{G}$ such that for some positive integer l the collection of annuli $\{A_l^m\}_{1\le m\le l}$ refines $\{\varPhi^{-1}(F)\}_{F\in\mathcal{F}}$. We shall call $\varPhi$ an *$\circledcirc_{\mathcal{F}}$-contraction of φ*.

Remark 11.8. If $\varPhi$ is an $\circledcirc_{\mathcal{F}}$-contraction of a map $\varphi\colon S^k \to X$, then for some positive integer l the collection $\{\bigcup_{j\le l,\varPhi(A_l^j)\subset F_i} A_l^j\}_{i\in I}$ is a cover of B^{k+1} by subcomplexes. We can always replace $\varPhi$ by another $\circledcirc_{\mathcal{F}}$-contraction such that this collection (possibly with different l) is an interior cover of B^{k+1}. The new $\circledcirc_{\mathcal{F}}$-contraction can be obtained by a reparametrization of $\varPhi$. In particular, we may assume that its image is equal to the image of $\varPhi$.

Remark 11.9. In definition 11.7, the sole role of a cover $\mathcal{G}$ is to limit the size of the image of $\varPhi$. The role of $\mathcal{F}$ is more interesting and may be interpreted in terms of small holes as follows. Let $\mathcal{F}$ be a cover of a space X and let φ be a k-hole in $\mathcal{F}$. Assume that φ is $\circledcirc_{\mathcal{F}}$-contractible and let $\varPhi$ be a map satisfying conditions of definition 11.7 for a positive integer l. Let φ_m be a map from S^k into X defined by the formula $\varphi_m(x) = \varPhi(mx/l)$. The hole φ_1 is small. The hole φ_2 might not be small, but if we patch φ_1 (a precise meaning of it will be given in the proof of lemma 11.10), then it would become small. Hence we may think that φ_2 is a "small hole of second order". Pushing this argument further, we see that φ_l, which is equal to φ, is a "small hole of lth order" and that it can be patched after recursively patching of $\varphi_1, \varphi_2, \dots$ up to φ_{l-1}.

Definition. We say that a cover $\mathcal{F} = \{F_i\}_{i\in I}$ is *$\circledcirc_k$-contractible in a cover $\mathcal{G}$* if it refines $\mathcal{G}$ and if for every non-empty subset J of I such that $F_J = \bigcap_{i\in J} F_i$ is non-empty, the kth homotopy group $\pi_k(F_J)$ has a set of generators that are $\circledcirc_{\mathcal{F}}$-contractible in $\mathcal{G}$, for some choice of base-points of F_J's.

Lemma 11.10. *If $\mathcal{F}$ is a k-regular ($k < n$) closed star-finite interior $\mathcal{N}_n$-cover that is $\circledcirc_k$-contractible in a closed cover $\mathcal{G}$, then there exists a closed $(k+1)$-regular interior $\mathcal{N}_n$-cover isomorphic to $\mathcal{F}$ and refining $\operatorname{st}_{\operatorname{st}^3_{\mathcal{F}}\mathcal{G}}\mathcal{F}$. Moreover, given a $Z(\mathcal{F})$-set Z, we may require that there exists a neighborhood of Z on which the constructed cover is equal to $\mathcal{F}$.*

Proof. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a k-regular ($k < n$) closed interior $\mathcal{N}_n$-cover of a space X that is $\circledcirc_k$-contractible in a closed cover $\mathcal{G}$. Let $\mathcal{J} = \{J \subset I \colon J \neq \emptyset \text{ and } F_J \neq \emptyset\}$. By proposition 2.5, for every $J \in \mathcal{J}$ the k-dimensional homotopy group $\pi_k(F_J)$ is countable. Therefore for every $J \in \mathcal{J}$ there exists a collection of maps $\{\psi_m^J\colon S^k \to F_J\}_{m\in\mathbb{N}}$ that generates $\pi_k(F_J)$. By the assumption that $\mathcal{F}$ is $\circledcirc_k$-contractible, we may additionaly require that ψ_m^J's are $\circledcirc_{\mathcal{F}}$-contractible in $\mathcal{G}$, i.e. that for every $J \in \mathcal{J}$ and every $m \in \mathbb{N}$ there exists an element G_m^J of $\mathcal{G}$ and a $\circledcirc_{\mathcal{F}}$-contraction $\Psi_m^J\colon B^{k+1} \to G_m^J$ of ψ_m^J. By remark 11.8, we may assume that the following condition is satisfied.

(1) There exists an integer l^J_m such that the collection $\{\bigcup\{A^j_{l^J_m}\colon 1 \le j \le l^J_m, \Psi^J_m(A^j_{l^J_m}) \subset F_i\}\}_{i\in I}$ is an interior cover of B^{k+1}.

By the assumptions, $\mathcal{F}$ is a countable star-finite interior cover, so it is locally finite. Hence every element of $\mathcal{J}$ is a finite subset of I. In particular, $\mathcal{J}$ is countable. We endow $\mathcal{J}$ with the discrete topology. Let $\Psi\colon B^{k+1}\times\mathcal{J}\times\mathbb{N}\to X$ be a map defined by the formula $\Psi(x,J,m)=\Psi^J_m(x)$, i.e. let Ψ be a disjoint union of Ψ^J_m's. The domain of Ψ is a k-dimensional locally finite countable simplicial complex. Let $\mathcal{U}$ be an open cover of X obtained via lemma 10.4 applied to $\mathcal{F}$. By theorem 3.6, there exists a $\mathcal{U}$-approximation of Ψ within $\mathcal{F}$ by a closed embedding $\varPhi$ with image disjoint from Z. Let $\varPhi^J_m\colon B^{k+1}\to X$ be a map defined by the formula $\varPhi^J_m(x)=\varPhi(x,J,m)$, i.e. an approximation of Ψ^J_m obtained as a restriction of Φ to an appropriate subset of its domain. Observe that $\varPhi^J_m$'s satisfy the following conditions.

(2) $\varphi^J_m=\varPhi^J_{m|S^k}$ maps S^k into F_J and $\{\varphi^J_m\}_{m\in\mathbb{N}}$ generates $\pi_k(F_J)$.

(3) There exists an integer l^J_m such that the collection $\{\bigcup\{A^j_{l^J_m}\colon 1 \le j \le l^J_m, \varPhi^J_m(A^j_{l^J_m}) \subset F_i\}\}_{i\in I}$ is an interior cover of B^{k+1}.

(4) Each $\varPhi^J_m$ is an embedding, the collection $\{\operatorname{im}\varPhi^J_m\}_{J\in\mathcal{J},m\in\mathbb{N}}$ is a discrete collection in X and $\operatorname{im}\varPhi^J_m\subset\operatorname{st}_{\mathcal{F}}G^J_m$.

Condition (2) is satisfied because φ^J_m is a $\mathcal{U}$-approximation within $\mathcal{F}$ of ψ^J_m; by the choice of $\mathcal{U}$, every two $\mathcal{U}$-close maps into F_J are homotopic and by the choice of ψ^J_m's, $\pi_k(F_J)$ is generated by $\{\psi^J_m\}_{m\in\mathbb{N}}$. Condition (3) is satisfied because φ^J_m is an approximation within $\mathcal{F}$ of ψ^J_m, so the collection defined in (1) refines the collection defined in (3). Condition (4) is satisfied because $\varPhi$ is a closed embedding into X and is $\mathcal{F}$-close to Ψ.

Let $L=\bigcup_{J\in\mathcal{J},m\in\mathbb{N}}\operatorname{im}\varPhi^J_m$. It is a locally finite simplicial complex embedded as a closed subset of X. By the construction it is disjoint from Z. For each $i\in I$, let

$$L_i=\bigcup\{\varPhi^J_m(A^j_{l^J_m})\colon J\in\mathcal{J}, m\in\mathbb{N}, 1\le j\le l^J_m, \varPhi^J_m(A^j_{l^J_m})\subset F_i\}.$$

By (3), $\mathcal{L}=\{L_i\}_{i\in I}$ is an interior cover of L by subcomplexes. By theorem 10.5, we may enlarge the pair $\langle L,\mathcal{L}\rangle$ to an n-core $\langle K_0,\mathcal{K}_0\rangle$ of $\mathcal{F}$, with $\mathcal{K}_0=\{K^0_i\}_{i\in I}$ and K_0 disjoint from Z. For each $J\in\mathcal{J}$ and $m\in\mathbb{N}$ fix a triangulation of a copy of B^{k+1} such that $A^j_{l^J_m}$ is a subcomplex of B^{k+1} for each $1\le j\le l^J_m$. To avoid undue proliferation of notation, each copy is denoted by the same symbol B^{k+1}. Let $p^J_m\colon C(B^{k+1})\to B^{k+1}$ be a projection along $[0,1]$. By theorem 3.6, we may approximate each $\widetilde{\varPhi}^J_m$ within $\mathcal{F}$ rel $B^{k+1}\times\{0\}$ by a closed embedding $\widehat{\varPhi}^J_m$ with image disjoint from Z. We may require that images of $\widehat{\varPhi}^J_m$'s form a discrete collection in X (by simultaneously approximating the disjoint union of $\varPhi^J_m$'s, like we did when approximating Ψ^J_m's). Since the approximation is within $\mathcal{F}$, $\operatorname{im}\widehat{\Phi}^J_m \subset \operatorname{st}_{\mathcal{F}} G^J_m$ and $\widehat{\Phi}^J_m({\Phi^J_m}^{-1}(K^0_i)) \subset F_i$ for each $i \in I$.

Let $K = K_0 \cup \bigcup_{J\in\mathcal{J}, m\in\mathbb{N}} \operatorname{im}\widehat{\Phi}^J_m$ and $\mathcal{K} = \{K_i\}_{i\in I}$, with $K_i = K^0_i \cup \bigcup \widehat{\Phi}^J_m({\Phi^J_m}^{-1}(K^0_i))$. By the construction, $\langle K, \mathcal{K}\rangle$ is an n-core of $\mathcal{F}$. By theorem 10.1, there exists a deformation of $\mathcal{F}$ to a closed star-finite interior $\mathcal{N}_n$-cover $\mathcal{F}_0$ of X such that $\langle K, \mathcal{K}\rangle$ is an exact n-core of $\mathcal{F}_0$. Since Z is disjoint from K, by letting $L = K$ when applying theorem 10.1, we may require that $\mathcal{F}_0$ is equal to $\mathcal{F}$ on a neighborhood of Z. We may also require that the deformation is so small, that $\mathcal{F}_0 \prec \operatorname{st}_{\mathcal{G}} \mathcal{F}$. Observe that $\langle K_0, \mathcal{K}_0\rangle$ is an exact n-core of $\mathcal{F}_0$ as well.

Order $\mathcal{J} \times \mathbb{N}$ into a sequence $\{(J_q, m_q)\}_{q\in\mathbb{N}}$. For each $q \in \mathbb{N}$ let $C_q = \operatorname{im}\widehat{\Phi}^{J_q}_{m_q}$ and let $D_q = \widehat{\Phi}^{J_q}_{m_q}(B^{k+1} \times \{1\})$ be the upper base of C_q. By the construction of $\widehat{\Phi}^J_m$'s, there exists a discrete collection $\{U_q\}_{q\in\mathbb{N}}$ of open subsets of X such that $D_q \subset U_q$ and $U_q \cap (K_0 \cup \bigcup_{r\neq q} C_r) = \emptyset$ for each $q \in \mathbb{N}$. By the construction, $D_q \subset \operatorname{st}_{\mathcal{F}} G^{J_q}_{m_q}$, hence we may require (and we do) that $U_q \subset \operatorname{st}^2_{\mathcal{F}} G^{J_q}_{m_q}$ (we use the assumption that $\mathcal{F}$ is an interior cover that refines $\mathcal{G}$). By proposition 3.2 and by corollary 3.7, there exists an approximation of the identity on U_q within $\mathcal{F}_0$ by a closed embedding $h_q\colon U_q \to U_q$ with image disjoint from D_q. By lemma 4.1 we may require (and we do) that $id_{X\setminus U_q} \cup h_q$ is a closed embedding into X. Let $V_q = U_q \setminus h_q(U_q)$. By the construction, V_q is an open neighborhood of D_q.

We shall recursively construct a sequence of closed interior $\mathcal{N}_n$-covers $\mathcal{F}_q = \{F^q_i\}_{i\in I}$ of X, $q \geq 1$, that are isomorphic to $\mathcal{F}_0$ and satisfy the following conditions.

(5) There exists an exact n-core K_q of $\mathcal{F}_q$ such that $K_{q-1} \cup C_q$ is a subcomplex of K_q and $K_q \setminus (K_{q-1} \cup C_q) \subset V_q$.

(6) $\mathcal{F}_q$ is equal to $\mathcal{F}_{q-1}$ on $C_q \cup (X \setminus V_q)$.

(7) The inclusion $K_{q-1} \cap F^{q-1}_J \subset K_q \cap F^q_J$ induces epimorphisms on homotopy groups of dimensions less than or equal to k.

(8) $\varphi^{J_q}_{m_q}$ is a trivial hole in $\mathcal{F}_q$.

Before we will do the construction, we will show that under these conditions the limit $\mathcal{F}_\infty = \lim_{q\to\infty} \mathcal{F}_q$ (see definition 4.2) is a cover that satisfies assertion of the lemma. By the construction, $\mathcal{F}$ is isomorphic to $\mathcal{F}_0$ and we required that $\mathcal{F}_0$ is isomorphic to $\mathcal{F}_q$ for each $q \geq 1$. By (6) and by the assumption that the collection $\{V_q\}$ is discrete, the sequence $\mathcal{F}_q$ stabilizes locally, so if elements of $\mathcal{F}_\infty$ intersect, then the corresponding elements of $\mathcal{F}_q$ intersect for large enough q. By (5), $\mathcal{F}_\infty$ is equal to $\mathcal{F}_0$ on K_0 and by the construction, K_0 is an exact core of $\mathcal{F}_0$, hence for each $J \in \mathcal{J}$ the intersection $K_0 \cap F^0_J = K_0 \cap F^\infty_J$ is non-empty. Therefore if elements of $\mathcal{F}_0$ intersect, then the corresponding elements of $\mathcal{F}_\infty$ intersect. Hence $\mathcal{F}_\infty$ is isomorphic to $\mathcal{F}$. As we noted, the sequence $\mathcal{F}_q$ stabilizes locally, hence by theorem 2.18, $\mathcal{F}_\infty$ is a closed interior $\mathcal{N}_n$-cover of X. Let $\mathcal{V} = \{V_q\}_{q\geq 1}$. By the construction, $\mathcal{F}_0 \prec \operatorname{st}_{\mathcal{F}} \mathcal{F}$

and $\mathcal{F}_\infty \prec \operatorname{st}_{\mathcal{V}} \mathcal{F}_0$. We assumed that $\mathcal{V} \prec \operatorname{st}^2_{\mathcal{F}} \mathcal{G}$, hence $\mathcal{F}_\infty \prec \operatorname{st}_{\operatorname{st}^2_{\mathcal{F}} \mathcal{G}} \operatorname{st}_{\mathcal{F}} \mathcal{F}$. By a direct computation, $\mathcal{F}_\infty \prec \operatorname{st}_{\operatorname{st}^3_{\mathcal{F}} \mathcal{G}} \mathcal{F}$. What is left to do is to show that $\mathcal{F}_\infty$ is a $(k+1)$-regular cover.

Let $K_\infty = \bigcup_{q \geq 1} K_q$. Let $f_q \colon X \to X$ be a closed embedding defined as the identity on $X \setminus \bigcup_{r>q} U_r$ and by the formula $f_q(x) = h_r(x)$ for each $x \in U_r$ with $r > q$. Let $\mathcal{H}_q = \{H^q_i\}_{i \in I}$ and let $H^q_i = f_q(F^q_i)$ for each $i \in I$. Observe that $\mathcal{H}_q$ is a closed interior $\mathcal{N}_n$-cover of $X \setminus \bigcup_{r>q} V_r$ and K_q is its exact n-core. Also, $\mathcal{H}_k$ is equal to $\mathcal{H}_{k-1}$ on $X \setminus U_k$ and $\mathcal{H}_{k-1}$ is a shrinking of $\mathcal{H}_k$ on $U_k \setminus V_k$. By lemma 10.8, K_∞ is an exact n-core of $\lim_{q\to\infty} \mathcal{H}_k$. Observe that $\lim_{q\to\infty} \mathcal{H}_q = \mathcal{F}_\infty$, so K_∞ is also an exact n-core of $\mathcal{F}_\infty$. Observe that if $J \in \mathcal{J}$, then $K_\infty \cap F^\infty_J = \bigcup_{q \geq 1} K_q \cap F^q_J$. By the assumptions, $K_0 \cap F^0_J$ has trivial homotopy groups up to dimension k, so by (8), $K_q \cap F^q_J$ has trivial homotopy groups up to dimension k. Hence $K_\infty \cap F^\infty_J$ has trivial homotopy groups up to dimension k, so $\mathcal{F}_\infty$ is k-regular. To see that it is $(k+1)$-regular consider a map $\varphi \colon S^k \to F^\infty_J$. Since K_∞ is an exact n-core of $\mathcal{F}_\infty$, φ is homotopic in F^∞_J with a map $\varphi' \colon S^k \to F^\infty_J \cap K_\infty$. Since the image of φ' is compact, it lies in $K_q \cap F^\infty_J$ for some $q \geq 1$. It follows from (6) and from the assumption that U_q is disjoint from $K_0 \cup \bigcup_{r \neq q} C_q$ that $K_q \cap F^\infty_J = K_q \cap F^q_J$. By (7), the inclusion $K_0 \cap F^0_J \subset K_q \cap F^q_J$ induces an epimorphism on k-dimensional homotopy groups. Hence φ' is homotopic, in $K_q \cap F^q_J$, with a finite product of φ^J_m's, for some $m \in \mathbb{N}$. By (8), φ^J_m is contractible in $K_r \cap F^r_J \subset K_\infty \cap F^\infty_J$ for r such that $J = J_r$ and $m = m_r$. Hence $\mathcal{F}_\infty$ is $(k+1)$-regular.



Figure 11.2: After patching of ξ_1, ξ_2 becomes a small hole in the middle set.

The construction of $\mathcal{F}_q$. Assume that we already constructed $\mathcal{F}_{q-1}$. Let $\mathcal{F}^0_{q-1} = \mathcal{F}_{q-1}$ and let $K^0_{q-1} = K_{q-1} \cup C_q$. By the construction, K^0_{q-1} is an exact n-core of $\mathcal{F}^0_{q-1}$. Let $l = l^{J_{q-1}}_{m_{q-1}}$. We shall construct a finite sequence $\mathcal{F}^1_{q-1}, \mathcal{F}^2_{q-1}, \ldots, \mathcal{F}^l_{q-1}$ of closed interior $\mathcal{N}_n$-covers of X that are isomorphic to $\mathcal{F}_{q-1}$ and satisfy the the following conditions for each $0 < m \leq l$. Below we let $\mathcal{F}^m_{q-1} = \{F^{m,q-1}_i\}_{i \in I}$, $F^{m,q-1}_J = \bigcap_{i \in J} F^{m,q-1}_i$ for each $J \in \mathcal{J}$ and $\xi_m = \Phi^{J_q}_{m_q}(xm/l)$.

(9) There exists an exact n-core K^m_{q-1} of $\mathcal{F}^m_{q-1}$ such that K^{m-1}_{q-1} is a subcomplex of K^m_{q-1} and $K^m_{q-1} \setminus K^{m-1}_{q-1} \subset V_q$.

(10) $\mathcal{F}^m_{q-1}$ is equal to $\mathcal{F}^{m-1}_{q-1}$ on $K^{m-1}_{q-1} \cup (X \setminus V_q)$.

(11) The inclusion $K^{m-1}_{q-1} \cap F^{m-1,q-1}_J \subset K^m_{q-1} \cap F^{m,q-1}_J$ induces epimorphisms on homotopy groups of dimensions less than or equal to k.

(12) ξ_m is a trivial hole in $\mathcal{F}^m_{q-1}$.

Obviously, $\mathcal{F}^l_{q-1}$ is a cover $\mathcal{F}_q$ that we are looking for. Assume that we already contructed $\mathcal{F}^{m-1}_{q-1}$ and K^{m-1}_{q-1}. By the construction, the collection $\{A^j_l\}_{1\leq j\leq l}$ of subsets of D_q refines $\mathcal{F}^{m-1}_{q-1}$. Hence for some $J \in \mathcal{J}$, ξ_m is homotopic with ξ_{m-1} in $F^{m-1,q-1}_J$. By (12), ξ_{m-1} is a trivial hole in $\mathcal{F}^{m-1}_{q-1}$, hence ξ_m is a small hole in $\mathcal{F}^{m-1}_{q-1}$. Let B_m be the "obvious" contraction of ξ_m (cf. figure 11.2). By lemma 11.6, there exists a closed interior $\mathcal{N}_n$-cover $\mathcal{F}^m_{q-1}$ obtained from $\mathcal{F}^{m-1}_{q-1}$ by patching of B_m in V_q. We let K^m_{q-1} to be an exact n-core of $\mathcal{F}^m_{q-1}$ that satisfies conditions of the definition of a cover obtained by patching. A direct verification shows that conditions (9)-(12) are satisfied. We are done. ☐

Remark 11.11. If $\mathcal{G} = \operatorname{st}^q \mathcal{E}$, then the cover constructed in lemma 11.10 refines $\operatorname{st}^{21q+10} \mathcal{E}$. It is beause $\mathcal{F}$ refines $\mathcal{G}$, the constructed cover refines $\operatorname{st}_{\operatorname{st}^3_{\mathcal{F}} \mathcal{G}} \mathcal{F}$ and

$$\operatorname{st}_{\operatorname{st}^3_{\mathcal{F}} \mathcal{G}} \mathcal{F} \prec \operatorname{st}_{\operatorname{st}^3_{\mathcal{G}} \mathcal{G}} \operatorname{st}^3_{\mathcal{G}} \mathcal{G} = \operatorname{st} \operatorname{st}^3 \mathcal{G} = \operatorname{st}^{10} \mathcal{G},$$

the last equality by lemma 2.2. By the same lemma, $\operatorname{st}^{10} \operatorname{st}^q \mathcal{E} = \operatorname{st}^{21q+10} \mathcal{E}$.

As an application we show that every n-semiregular cover is $\circledcirc_0$-contractible, proving theorem 6.7 for $k = 0$.

11.3 Proof of theorem 6.7 for $k = 0$

Let $\mathcal{F} = \{F_i\}_{i\in I}$ be an n-semiregular ($n > 0$) closed star-finite interior $\mathcal{N}_n$-cover n-contractible in a closed cover $\mathcal{E}$. By the assumption that $\mathcal{F}$ is n-semiregular, there exists an anticanonical map λ of $\mathcal{F}$. By the definition, every approximation within $\mathcal{F}$ of λ is an anticanonical map of $\mathcal{F}$. Hence by theorem 3.6, we may assume that λ is a closed embedding. Fix a non-empty subset J of I. Let σ_J be a barycenter of a simplex of the nerve of $\mathcal{F}$ that is spanned by vertices corresponding to elements of $\mathcal{F}$ that are indexed by J. We let $c_0 = \lambda(\sigma_J)$ be the base point of F_J. By the assumption, $\mathcal{F}$ is an $\mathcal{N}_n$-cover and $n > 0$, so F_J is separable and locally connected, hence there are only countably many connected components of F_J. Let $c_1, c_2, \dots$ be a sequence of points of F_J such that every connected component of F_J has an element in the sequence. A set of maps $\varphi_k \colon S^0 \to F_J$ that map 1 to c_0 and -1 to c_k generates $\pi_0(F_J)$. We will show that for each k, φ_k is $\circledcirc_{\mathcal{F}}$-contractible in an element of $\operatorname{st} \mathcal{E}$.

By the assumption that $\mathcal{F}$ is n-contractible in $\mathcal{E}$, there exists a path $\gamma \colon [0,1] \to X$ that connects c_k with c_0 inside an element of $\mathcal{E}$, i.e. such that $\gamma(0) = c_k$, $\gamma(1) = c_0$

and $\operatorname{im}\gamma$ is a subset of some $E \in \mathcal{E}$. Since $\mathcal{F}$ is an interior cover, there exists a positive integer l such that $\{\gamma([(m-1)/l, m/l])\}_{0<m\leq l}$ refines $\mathcal{F}$. For each $0 < m \leq l$, let j_m be an element of I such that $\gamma([(m-1)/l, m/l]) \subset F_{j_m}$. Without a loss of generality we may assume that γ is constant on small neighborhoods of 0 and 1, so we may assume that j_1 and j_l are elements of J. Consider a function $[(m-1)/l, m/l] \mapsto \lambda(\operatorname{bst} v(F_{j_m}))$ defined on a collection $\{[(m-1)/l, m/l]\}_{0<m\leq l}$. It is a carrier and a constant map $\beta\colon \{0,1\} \to X$ that maps 0 and 1 to c_0 is carried by it. By lemma 5.3, the cover of the nerve of $\mathcal{F}$ by barycentric stars of its vertices is regular for the class of metric spaces, hence by the carrier theorem, β can be extended over $[0,1]$ to a map $\beta\colon [0,1] \to X$ such that $\beta([(m-1)/l, m/l]) \subset \lambda(\operatorname{bst} v(F_{j_m})) \subset F_{j_m}$ for each $0 < m \leq l$. Let $\Phi\colon [-1,1] \to X$ be a map defined by the formula

$$\Phi(x) = \begin{cases} \gamma(x+1) & x \in [-1,0] \\ \beta(x) & x \in [0,1] \end{cases} .$$

This map is a $\circledcirc_{\mathcal{F}}$-contraction of φ directly from the definition. The image of Φ is contained in $\operatorname{st}_{\mathcal{F}} \operatorname{im}\gamma$, $\operatorname{im}\gamma$ is contained in an element of $\mathcal{E}$ and $\mathcal{F}$ refines $\mathcal{E}$, hence the image of Φ is contained in an element of $\operatorname{st}\mathcal{E}$.

We proved that $\mathcal{F}$ is $\circledcirc_0$-contractible in $\operatorname{st}\mathcal{E}$. By lemma 11.10 and by remark 11.11, there exists a 1-regular n-semiregular closed interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$, that refines $\operatorname{st}^{31}\mathcal{E}$ and that is equal to $\mathcal{F}$ on some open neighborhood of Z. We are done.

11.4 $\Cup$-Contractibility

Definition. Let $\mathcal{F}$ and $\mathcal{G}$ be covers of a space X. Let A be a subspace of X. Let φ be a map of a pair (B^k, S^{k-1}) into a pair (X, A). We say that φ is *$\Cup_{\mathcal{F}}$-contractible to A in $\mathcal{G}$* if it admits a homotopy $\Phi\colon (B^k, S^{k-1}) \times [0,1] \to (X, A)$ with a constant map and such that the image of Φ lies in an element of $\mathcal{G}$ and for some integer l, the collection $\{B^k \times [(m-1)/l, m/l]\}_{1\leq m\leq l}$ refines $\Phi^{-1}(\mathcal{F})$. We shall call Φ a *$\Cup_{\mathcal{F}}$-contraction of φ to A in $\mathcal{G}$*.

Definition. Let $0 < k < n$ and let A be a subspace of a space X. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a cover of X. Assume that each non-empty F_J has a basepoint x_J in the intersection $F_J \cap A$. We say that $\mathcal{F}$ is *$\Cup_k$-contractible to A in a cover $\mathcal{G}$* if for each non-empty subset J of I such that F_J is non-empty there exists a set of maps $\{\varphi^J_m\colon (B^k, S^{k-1}) \to (F_J, F_J \cap A)\}_{m\geq 1}$ that are $\Cup_{\mathcal{F}}$-contractible in $\mathcal{G}$ and such that $i_*^{-1}(\{\varphi^J_m\}_{m\geq 1})$ generates $\pi_k(F_J)$, where $i_*\colon \pi_k(F_J) \to \pi_k(F_J, F_J \cap A)$ is a function induced by the inclusion $(F_J, x_J) \subset (F_J, F_J \cap A)$.

Lemma 11.12. *Let $0 < k < n$ and let A be a subspace of a space X. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a cover of X. As usual, let $F_J = \bigcap_{i\in J} F_i$ for each non-empty subset J of I. Assume that each non-empty F_J has a base-point x_J in the intersection $F_J \cap A$. Let $\mathcal{G}$ be a* *closed cover of X. Assume that for each non-empty subset J of I the $(k-1)$th homotopy group of $A \cap F_J$ is trivial and that the inclusion of $A \cap F_J$ into F_J induces a trivial (zero) homomorphism on kth homotopy groups. If $\mathcal{F}$ is $\mathbb{U}_k$-contractible to A in $\mathcal{G}$, then $\mathcal{F}$ is $\circledcirc_k$-contractible in* $\operatorname{st}\mathcal{G}$.

Proof. Fix $m \geq 1$ and fix a non-empty $J \subset I$ such that $F_J \neq \emptyset$. Let $\{\varphi^J_m\}_{m\geq 1}$ be a set of generators of $\pi_k(F_J)$ such that compositions $\varphi^J_m \circ \xi$ are $\mathbb{U}_{\mathcal{F}}$-contractible to A in $\mathcal{G}$. Pick $\varphi = \varphi^J_m$. We will show that φ is $\circledcirc_{\mathcal{F}}$-contractible in $\operatorname{st}\mathcal{G}$, thus proving the assertion directly from the definition of $\circledcirc_k$-contractibility. Let B^k_+ be the north hemisphere of S^k, let B^k_- be the south hemisphere of S^k and let $S^{k-1} = B^k_+ \cap B^k_-$ be the equator of S^k. Without a loss of generality, we may let B^k_+ denote the domain of φ. Let $\varPhi$ be a $\mathbb{U}_{\mathcal{F}}$-contraction to A of $\varphi \circ \xi$ in some element G of $\mathcal{G}$. It is a homotopy of φ with a constant map. For $0 \leq t \leq 1$ we let $\varPhi_t \colon B^k_+ \to X$ denote $\varPhi$ restricted to a time t. By the definition of $\mathbb{U}_{\mathcal{F}}$-contraction to A, for each t, $\varPhi_t(S^{k-1}) \subset A$ and there exists a positive integer l and a sequence of elements $j_1, j_2, \ldots, j_l \in I$ such that

(*) $\varPhi_t \subset F_{j_q}$ for each $0 < q \leq l$ and each t such that $(q-1)/l \leq t \leq q/l$.

We will extend $\varPhi$ over $B^k_- \times [0,1]$ to a $\circledcirc_{\mathcal{F}}$-contraction of φ.

By the definition, $\varPhi_0$ restricted to $S^{k-1} \times \{0\}$ is constant. We extend $\varPhi_0$ onto $B^k_- \times \{0\}$ to be constant as well, with value determined by the value of $\varPhi$ on $S^{k-1} \times \{0\}$. Again by the definition, $\varPhi_1$ restricted to $S^{k-1} \times \{1\}$ is constant and we extend $\varPhi_1$ onto $B^k_- \times \{1\}$ to be a constant map. For each integer $0 < q < l$, $\varPhi_{q/l}$ maps $S^{k-1} \times \{q/l\}$ into $F_{j_{q-1}} \cap F_{j_q} \cap A$. By the assumptions, this restriction is contractible in $F_{j_{q-1}} \cap F_{j_q} \cap A$. We extend $\varPhi$ over $B^k_- \times \{q/l\}$ to be any such contraction. For each $0 < q \leq l$, the (extended) map $\varPhi_{|S^{k-1}\times[(q-1)/l,q/l]\cup B^k_-\times\{(q-1)/l,q/l\}}$ is a map from a k-dimensional sphere into $A \cap F_{j_q}$. By the assumptions, every such map extends over $B^k_- \times [(q-1)/l, q/l]$ to a map into F_{j_q}. We define an extension of $\varPhi$ onto $B^k_- \times [(q-1)/l, q/l]$ to be any such extension.

From the construction, the extended $\varPhi$ is a $\circledcirc_{\mathcal{F}}$-contraction of $\varPhi_0$ and $\varPhi_0$ is homotopic with φ in F_J. Its image lies in $\operatorname{st}_{\mathcal{F}}\mathcal{G}$. Since $\mathcal{F}$ refines $\mathcal{G}$, it is a $\circledcirc_{\mathcal{F}}$-contraction in $\operatorname{st}\mathcal{G}$.

□

Example 11.13. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a cover of a space X. Assume that $\mathcal{F}$ admits an anticanonical map λ that is a closed embedding and such that $\lambda(\operatorname{bst} v(F_i)) = F_i \cap \operatorname{im}\lambda$ for each $i \in I$. By lemma 11.12, if $\mathcal{F}$ is $\mathbb{U}_k$-contractible to $\operatorname{im}\lambda$ in $\mathcal{G}$, then $\mathcal{F}$ is $\circledcirc_k$-contractible in $\operatorname{st}\mathcal{G}$.

11.5 Patching of large holes

Definition 11.14. Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a cover of a space X and let $F_J = \bigcap_{i\in J} F_i$ for each non-empty subset J of I. Let τ be the set of simplices of a triangulated

k-dimensional sphere S^k. Let $\varphi\colon S^k \to F_J$ be a k-hole in F_J. Let

$$C = \left(S^k \times [-1,1] \times \{0\}\right) \cup \left((S^k)^{(k-2)} \times [-1,1] \times [0,1]\right),$$

where $(S^k)^{(k-2)}$ denotes the $(k-2)$-dimensional skeleton of S^k (empty if $k=1$). We say that a map $\Phi\colon C \to X$ is a *patch for* φ if the following conditions are satisfied.

(1) Φ restricted to $S^k = S^k \times \{-1\} \times \{0\}$ is equal to φ,

(2) Φ restricted to $S^k \times \{1\} \times \{0\}$ is a trivial hole in F_J,

(3) there is an integer l such that the collection

$$\{\Phi((\delta \times B_{m,l}) \cap \operatorname{dom}\Phi)\colon \delta \in \tau, 1 \le m \le l\}$$

refines $\mathcal{F}$, where $B_{m,l} = \{x \in [-1,1] \times [0,1]\colon |x|_\infty \in [\frac{m-1}{l}, \frac{m}{l}]\}$ (see figure 11.3).



Figure 11.3: $B_{m,l} = \{x \in [-1,1] \times [0,1]\colon |x|_\infty \in [\frac{m-1}{l}, \frac{m}{l}]\}$.

Remark 11.15. Let $\mathcal{F} = \{F_i\}_{i \in I}$. If Φ is a patch for a k-hole φ if $\mathcal{F}$, then by the definition there exists a triangulation τ of S^k and an integer l such that the collection

$$\left\{\bigcup \{\delta \times B_{m,l}\colon \delta \in \tau, 1 \le m \le l, \Phi(\delta \times B_{m,l}) \subset F_i\}\right\}_{i \in I}$$

is a cover of the domain of Φ by subcomplexes. We can always replace Φ by another patch such that this collection (possibly for a finer triangulation τ and a larger integer l) is an interior cover of the domain of Φ. The new patch can be obtained by a reparametrization of Φ. In particular, we may assume that its image is equal to the image of Φ.

We'll prove below two lemmas. The first of them states that every hole in $\mathcal{F}$ admits a patch the size of which is controlled, while the next lemma enables one to use this patch to patch the hole. Some new holes may be added in the process, but they will be handled by lemma 11.12.

Lemma 11.16. *If $\mathcal{F}$ is a star-finite n-semiregular interior cover that is n-contractible in a cover $\mathcal{E}$ and φ is a k-hole in $\mathcal{F}$ for some $k < n$, then φ has a patch whose image is contained in an element of* $\operatorname{st}\mathcal{E}$*, for some sufficiently fine triangulation of S^k.*

Proof. We denote the underlying space by X. Let $\mathcal{F} = \{F_i\}_{i \in I}$ and let $\varphi\colon S^k \to F_J$ be a k-hole in F_J for some non-empty $J \subset I$. Identify S^k with $S^k \times \{-1\} \times \{0\}$ and let $\varPhi\colon S^k \times \{-1\} \times \{0\} \to X$ to be equal to φ. We'll extend $\varPhi$ consecutively onto $S^k \times [-1, 0] \times \{0\}$, $S^k \times [0, 1] \times \{0\}$ and $(S^k)^{(k-2)} \times [-1, 1] \times [0, 1]$ to obtain a patch for φ. We shall fix a triangulation of S^k after the first extension. In the first step we will use the assumption that $\mathcal{F}$ is n-contractible in $\mathcal{E}$; in the second step that it is n-semiregular; in the third step that it is k-regular.

First step. Let $C_1 = S^k \times [-1, 0] \times \{0\}$. Let λ be an anticanonical map of $\mathcal{F}$. For each element F of $\mathcal{F}$ we let $v(F)$ be the vertex in the nerve of $\mathcal{F}$ that corresponds to F. Let σ_J be the barycenter of a simplex of $N(\mathcal{F})$ spanned by the vertices $v(F_j)$, $j \in J$. We define $\varPhi$ on $S^k \times \{0\} \times \{0\}$ to be a constant map with value $\lambda(\sigma_J)$. Since $\mathcal{F}$ is n-contractible in $\mathcal{E}$ and $\varPhi$ (so far defined) has image contained in F_J and a domain whose dimension is less than n, $\varPhi$ extends over C_1 to a map with image contained in some element of $\mathcal{E}$. We define $\varPhi$ on C_1 to be any such extension.

Second step. Let $C_2 = S^k \times [0, 1] \times \{0\}$. We first fix a triangulation τ of S^k and an integer l such that for every $1 \le m \le l$ and every $\delta \in \tau$ the set $\varPhi(\delta \times [-m/l, (-m + 1)/l]) \times \{0\}$ is a subset of $F_{i_{\delta,m}}$ for some $i_{\delta,m} \in I$. Such a pair exists because $\mathcal{F}$ is an interior cover and C_1 is compact. Without a loss of generality we may assume that $i_{\delta,1} \in J$ and $i_{\delta,l} \in J$ for each $\delta \in \tau$. A function that assigns $\operatorname{bst} v(F_{i_{\delta,m}})$ to a subset $\delta \times [(m-1)/l, m/l] \times \{0\}$ of C_2 is a carrier. A constant map on $S^k \times \{0, 1\} \times \{0\}$ with value σ_J is carried by it. By lemma 5.3, the cover of the nerve of $\mathcal{F}$ by barycentric stars of vertices is n-regular. Therefore by the carrier theorem, the constant map extends over C_2 to a map that maps $\delta \times [(m-1)/l, m/l]) \times \{0\}$ into $\operatorname{bst} v(F_{i_{\delta,m}})$. We define $\varPhi$ on C_2 to be a composition with λ of any such extension.

Third step. Let $C_3 = (S^k)^{(k-2)} \times [-1, 1] \times [0, 1]$. By the construction, $\varPhi((C_1 \cup C_2) \cap (\delta \times B_{m,l})) \subset F_{i_{\delta,m}}$ for each $\delta \in \tau$ and $1 \le m \le l$. Therefore the map $\delta \times B_{m,l} \mapsto F_{i_{\delta,m}}$ is a carrier. By assumptions $\mathcal{F}$ is a k-regular cover and by the definition C_3 is k-dimensional. Hence by the carrier theorem, $\varPhi$ extends over C_3 to a map carried by this carrier. We define $\varPhi$ on C_3 to be any such extension.

By the construction $\varPhi$ is a patch for φ. Observe that $\varPhi(C_1)$ is contained in some element of $\mathcal{E}$. Therefore by condition (3) of the definition of a patch, since $\mathcal{F}$ refines $\mathcal{E}$, the image of $\varPhi$ must be contained in an element of $\operatorname{st}\mathcal{E}$. □

Lemma 11.17. *Let $0 < k < n$. Let $\mathcal{F} = \{F_i\}_{i \in I}$ be a closed countable k-regular star-finite interior $\mathcal{N}_n$-cover of a space X. Let J be a nonempty subset of I and let φ be a closed embedding of S^k into $F_J = \bigcap_{i \in I} F_i$. Let $\varPhi$ be a closed embedding and a patch for φ. Let L be an exact n-core of $\mathcal{F}$ such that* $\operatorname{im}\varPhi$ *is a subcomplex of L. Let A be a subcomplex of L such that* $\operatorname{im}\varPhi$ *is a subcomplex of A. Let $\mathcal{H}$ be a cover of L such that $\mathcal{F}/L$ is $\mathbb{W}_k$-contractible to A in $\mathcal{H}$. For every neighborhood U of* $\operatorname{im}\varPhi$ *there* *exists a cover $\mathcal{G}$ of X and an exact n-core K of $\mathcal{G}$ such that the following conditions are satisfied.*

(1) $\mathcal{G}$ is a k-regular closed interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$ and is equal to $\mathcal{F}$ on $L \cup (X \setminus U)$,

(2) L is a subcomplex of K and $K \setminus L \subset U$,

(3) $\mathcal{G}/K$ is $\mathbb{U}_k$-contractible to A in $\mathcal{H} \cup \{U \cap K\}$,

(4) the element φ of $\pi_k(L \cap F_J) = \pi_k(L \cap G_J)$ becomes 0 in $\pi_k(K \cap G_J)$.

In the next section we apply lemma 11.17 recursively as to kill all generators of $\pi_k(L \cap F_J)$, $J \subset I$, to be able to use lemma 11.12 and reduce the proof of theorem 6.7 to the previously solved case of $\circledcirc_k$-contractible covers.

Proof. Let τ be a set of simplices of S^k and let l be a positive integer such that conditions of definition 11.14 are satisfied for Φ. Order $(k-1)$-dimensional simplices from τ into a sequence $\delta_0, \delta_1, \ldots, \delta_q$. Let $C_{0,0}$ denote the domain of Φ. For each $0 \leq p \leq q$ and for each $1 \leq m \leq l$, let $C_{p,m}$ be a simplicial complex defined recursively by the identity $C_{p,m} = C_{p,m-1} \cup (\delta_p \times B_{m,l})$ with $C_{p,0}$ identified with $C_{p-1,l}$ for $p > 0$. Note that

$$C_{q,l} = (S^k \times [-1,1] \times \{0\}) \cup \left((S^k)^{(k-1)} \times [-1,1] \times [0,1]\right)$$

and compare $C_{q,l}$ with complex C from definition 11.14. Let $D_{p,m} = (\delta_p \times B_{m,l}) \cap C_{p,m-1}$ for each $0 \leq p \leq q$ and each $0 < m \leq l$.



Figure 11.4: Patching of $(k-1)$-holes.

Since Φ is an embedding, we may identify $C_{0,0}$ with the image of Φ and treat it as a subcomplex of L. Below we shall recursively embed $C_{p,m}$'s into X and we will use same symbols to denote images of these embeddings. Let $\mathcal{F}_{0,0} = \mathcal{F}$ and $L_{0,0} = L$. We shall recursively construct a sequence of covers $\mathcal{F}_{p,m} = \{F_i^{p,m}\}_{i \in I}$ and their exact n-cores $L_{p,m}$ in such a way that the following conditions are satisfied ($L_{p,0}$ is identified with $L_{p-1,l}$ for each $p > 0$).

(i) $\mathcal{F}_{p,m}$ is a k-regular closed interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$ and is equal to $\mathcal{F}$ on $L \cup (X \setminus U)$,

(ii) $C_{p,m}$ is embedded into X in such a way that $L_{p,m} = L_{p,m-1} \cup C_{p,m}$ and $C_{p,m}$ is a subset of U,

(iii) $\mathcal{F}_{p,m}/L_{p,m}$ is $\mathbb{U}_k$-contractible to A in $\mathcal{H} \cup \{U \cap L_{p,m}\}$,

(iv) if $D_{p,m} \subset F_i^{p,m-1}$, then $\delta_p \times B_{m,l} \subset F_i^{p,m}$; if $\partial D_{p,m} \subset F_i^{p,m-1}$, then $B_{p,m+1} \subset F_i^{p,m}$.

Identify $\mathcal{F}_{p,0}$ with $\mathcal{F}_{p-1,l}$ for each $p > 0$ and assume that we already contructed $\mathcal{F}_{p,m-1}$. By conditions (i) and (iv), conditions of lemma 11.6 are satisfied with $\mathcal{F} = \mathcal{F}_{p,m-1}$, $K = L_{p,m-1}$ and $B = B_{p,m}$. Hence there exists a closed interior $\mathcal{N}_n$-cover $\mathcal{F}_{p,m}$ with an exact n-core $L_{p,m}$ that is obtained from $\mathcal{F}_{p,m-1}$ by patching of $B_{p,m}$ in U. We may identify $C_{p,m} \setminus C_{p,m-1}$ with $L_{p,m} \setminus L_{p,m-1}$, as these sets are homeomorphic by the definitions. Conditions (i), (ii), (iii) and (iv) are satisfied directly from the definitions.

Order k-dimensional simplices of τ into a sequence $\Delta_0, \Delta_1, \ldots, \Delta_Q$. Let $E_{0,0} = C_{q,l}$. For each $0 \leq p \leq Q$ and for each $1 \leq m \leq l$, let $E_{p,m}$ be a simplicial complex defined recursively by the identity $E_{p,m} = E_{p,m-1} \cup (\Delta_p \times B_{m,l})$ if $k < n-1$ or by the identity $E_{p,m} = E_{p,m-1} \cup (\Delta_p \times \partial B_{m,l})$ if $k = n-1$. In either case we identify $E_{p,0}$ with $E_{p-1,l}$ for each $p > 0$. Let $F_{p,m} = (\Delta_p \times B_{m,l}) \cap E_{p,m-1}$ for each $0 \leq p \leq Q$ and each $0 < m \leq l$.


(a) (b)

Figure 11.5: Patching of k-holes.

By the definition, $E_{0,0}$ is a subset of X. Below we shall recursively embed $E_{p,m}$'s into X and we will use same symbols to denote images of these embeddings. Let $\mathcal{G}_{0,0} = \mathcal{F}_{q,l}$ and let $K_{0,0} = L_{q,l}$. We shall recursively construct a sequence of covers $\mathcal{G}_{p,m} = \{G_i^{p,m}\}_{i \in I}$ and their exact n-cores $K_{p,m}$ in such a way that the following conditions are satisfied ($K_{p,0}$ is identified with $K_{p-1,l}$ for each $p > 0$).

(I) $\mathcal{G}_{p,m}$ is a k-regular closed interior $\mathcal{N}_n$-cover of X that is isomorphic to $\mathcal{F}$ and is equal to $\mathcal{F}$ on $L \cup (X \setminus U)$,

(II) $E_{p,m}$ is embedded into X in such a way that $K_{p,m} = K_{p,m-1} \cup E_{p,m}$ and $E_{p,m}$ is a subset of U.

(III) $\mathcal{G}_{p,m}/K_{p,m}$ is $\mathbb{W}_k$-contractible to A in $\mathcal{H} \cup \{U \cap K_{p,m}\}$,

(IV) if $F_{p,m} \subset G_i^{p,m-1}$, then $\Delta_p \times B_{m,l} \subset G_i^{p,m}$ if $k < n-1$ and $\Delta_p \times \partial E_{p,m} \subset G_i^{p,m}$ if $k = n-1$; if $\partial F_{p,m} \subset G_i^{p,m-1}$, then $F_{p,m+1} \subset G_i^{p,m}$.

Again, we obtain $\mathcal{G}_{p,m}$ from $\mathcal{G}_{p,m-1}$ by patching of $E_{p,m}$ in U. All four conditions are verified easily.

Let $\mathcal{G} = \mathcal{G}_{Q,l}$ and $K = K_{Q,l}$. Observe that if $\operatorname{im}\varphi \subset F_i$ for i in I, then by (iv) and (IV), for every simplex $\delta \in \tau$, $\delta \times \{x \in [-1,1] \times [0,1] \colon |x|_\infty = 1\} \subset G_i^{Q,l}$. Hence φ is contractible in $G_i^{Q,l}$. By (I), (II) and (III), $\mathcal{G}$ satisfies conditions (1), (2) and (3) of the lemma. We are done. ☐

11.6 Proof of theorem 6.7 for $k > 0$

Let $\mathcal{F} = \{F_i\}_{i\in I}$ be a k-regular n-semiregular $(0 < k < n)$ closed countable star-finite interior $\mathcal{N}_n$-cover of a space X. Assume that $\mathcal{F}$ is n-contractible in a closed cover $\mathcal{E}$ of X. Let Z be a $Z(\mathcal{F})$-set. We shall construct a closed $(k+1)$-regular n-semiregular closed countable star-finite interior $\mathcal{N}_n$-cover of X that refines $\operatorname{st}^{472}\mathcal{E}$, that is isomorphic to $\mathcal{F}$ and that is equal to $\mathcal{F}$ on a neighborhood of Z.

Let $\mathcal{J}$ be a set of those non-empty subsets J of I, for which the intersection $F_J = \bigcap_{i\in I} F_i$ is non-empty. The set $\mathcal{J}$ is countable, because $\mathcal{F}$ is countable and locally finite. By proposition 2.5, the set of homotopy classes of maps from S^k into F_J is countable for each J in $\mathcal{J}$. Hence there exists a sequence J_m of elements of $\mathcal{J}$ and a sequence of maps $\varphi_m \colon S^k \to F_{J_m}$ such that for each $J \in \mathcal{J}$, the set $\{\varphi_m\}_{m\geq 1, J_m = J}$ contains elements from all homotopy classes of maps from S^k into F_J. By lemma 11.16, for each φ_m there exists a patch $\varPhi_m$ whose image is contained in an element of $\operatorname{st}\mathcal{E}$. We assume that the assertion of remark 11.15 is true for $\varPhi_m$, i.e. that the collection defined in its statement is an interior cover of $\varPhi_m$. Let $\mathcal{O}$ be an open cover of X obtained via lemma 10.4 applied to $\mathcal{F}$. Let $\varPhi$ be a disjoint union of $\varPhi_m$'s. By theorem 2.10, the domain of $\varPhi$ is Polish. By theorem 3.6, there exists a $\mathcal{O}$-approximation Ψ of $\varPhi$ within $\mathcal{F}$ by a closed embedding. By proposition 3.2 and by corollary 3.7, we may assume that the image of Ψ is disjoint from Z. By a slight abuse of notation, we may let Ψ_m denote the restriction of Ψ to the domain of $\varPhi_m$ and let ψ_m denote the restriction of Ψ to the domain of φ_m. By the construction, the collection of images of Ψ_m's is discrete in X, ψ_m maps S^k into F_{J_m}, for each J in $\mathcal{J}$ the set $\{\psi_m\}_{m\geq 1, J_m = J}$ contains elements from all homotopy classes of maps from S^k into F_J, Ψ_m is a patch for ψ_m and the image of Ψ_m is contained in $\operatorname{st}_{\mathcal{F}}\mathcal{E}$.

Let τ_m be a triangulation of S^k and let l_m be an integer such that $\varPsi_m$ satisfies conditions of the definition of a patch for $\tau = \tau_m$ and $l = l_m$. Let $M = \operatorname{im} \Psi$ and let

$$\mathcal{M} = \left\{ M_i = \bigcup \{ \varPhi(\delta \times B_{m,l_m}) \colon \delta \in \tau_m, 1 \leq m \leq l_m, \varPhi(\delta \times B_{m,l_m}) \subset F_i \} \right\}_{i \in I}.$$

By the assumption we made on $\varPhi_m$'s, $\mathcal{M}$ is an interior cover of M. By the definition, it refines $\mathcal{F}$. Hence by theorem 10.5, there exists an n-core $\langle L_0, \mathcal{L}_0 \rangle$ of $\mathcal{F}$ such that M is a subcomplex of L_0 and such that $\mathcal{L}_0$ is equal to $\mathcal{M}$ on M. Let $\widetilde{L}$ be a disjoint union $L_0 \sqcup \bigsqcup_{m \geq 1} (\operatorname{im} \varPhi_m \times [0,1])$, divided by an equivalence relation that for each $m \geq 1$ identifies $\operatorname{im} \varPhi_m \subset L_0$ with $\operatorname{im} \varPhi_m \times \{0\} \subset \operatorname{im} \varPhi_m \times [0,1]$. Let $\widetilde{p}$ be a restriction to the n-dimensional skeleton of $\widetilde{L}$ of a natural projection from $\widetilde{L}$ onto L_0. By theorem 3.6, there exists an approximation of $\widetilde{p}$ within $\mathcal{F} \operatorname{rel} L_0$ by a closed embedding p. By proposition 3.2 and by corollary 3.7, we may assume that the image of p is disjoint from Z. We let $\mathcal{L} = \{ p(\widetilde{p}^{-1}(L_0^i)) \}_{i \in I}$. We let $L = \operatorname{im} p$. We let $L_m = L_{m-1} \cup p(\widetilde{p}^{-1}(\operatorname{im} \Psi_m))$. By the construction, for each $m \geq 0$, $\langle L_m, \mathcal{L}/L_m \rangle$ is an n-core of $\mathcal{F}$. Let $\widetilde{\Xi}_m$ be the unique map onto $\operatorname{im} \varPhi_m \times \{1\}$ such that $p \circ \Xi_m = \varPhi_m$ and let $\widetilde{\xi}_m$ be the unique map onto $\operatorname{im} \varphi_m \times \{1\}$ such that $p \circ \xi_m = \varphi_m$. Let $\Xi_m = p \circ \widetilde{\Xi}_m$ and let $\xi_m = p \circ \widetilde{\xi}_m$. By the construction, Ξ_m is an approximation of Ψ_m within $\mathcal{F}$. Hence Ξ_m is a patch for ξ_m. The image of Ψ_m lies in an element of $\operatorname{st}_{\mathcal{F}} \mathcal{E}$ and so does the image of Ξ_m. By the construction, ξ_m is homotopic with ψ_m in F_{J_m}. Let $\mathcal{U} = \{U_m\}_{m \geq 1}$ be a discrete collection of open subsets of X such that $\operatorname{im} \Xi_m \subset U_m$, $U_m \cap Z = \emptyset$, $L_{m-1} \cap U_m = \emptyset$ and $(L_{m+1} \setminus L_m) \cap U_m = \emptyset$ for each $m \geq 1$. Since the collection $\{\operatorname{im} \Xi_m\}_{m \geq 1}$ refines $\operatorname{st}_{\mathcal{F}} \mathcal{E}$, we may assume that $\mathcal{U}$ refines $\operatorname{st}^2_{\mathcal{F}} \mathcal{E}$, because $\mathcal{F}$ is an interior cover (which implies that for each set A, $\operatorname{st}_{\mathcal{F}} A$ is a neighborhood of A).


U_1
$\operatorname{im} \Xi_1$
$L_1 \setminus L_0$
$\operatorname{im} \varPhi_1$
L_0
U_2
$\operatorname{im} \Xi_2$
$L_2 \setminus L_1$
$\operatorname{im} \varPhi_2$

Figure 11.6: Some of the sets constructed in the proof of theorem 6.7.

By theorem 10.1, there exists a deformation of $\mathcal{F}$ to a closed interior $\mathcal{N}_n$-cover $\mathcal{F}_{-1} = \{F_i^{-1}\}_{i \in I}$ of X with an exact n-core $\langle L, \mathcal{L} \rangle$. We may assume that $\mathcal{F}_{-1}$ is equal to $\mathcal{F}$ on a neighborhood of Z and that $\mathcal{F}_{-1}$ refines $\operatorname{st} \mathcal{F}$. By the construction, each Ξ_m is a patch for ξ_m and for each $J \in \mathcal{J}$, the set $\{\xi_m\}_{m \geq 1, J_m = J}$ contains elements from all homotopy classes of maps from S^k into F_J.

By lemma 4.1, for each $m \geq 1$ there exists an open cover $\mathcal{U}_m$ of U_m such that every map from U_m into U_m that is $\mathcal{U}_m$-close to the identity on U_m extends continuously by the identity on $X \setminus U_m$. By theorem 2.18, each U_m is an $\mathcal{N}_n$-space and by remark 3.5, $\operatorname{im}\Xi_m$ is a $Z(\mathcal{F}/U_m)$-set in U_m. By corollary 3.7, there exists a $\mathcal{U}_m$-approximation $h_m\colon U_m \to U_m$ within $\mathcal{F}_{-1}$ of the identity on U_m by a closed embedding with image disjoint from $\operatorname{im}\Xi_m$. We let $V_m = U_m \setminus \operatorname{im} h_m$. Note that V_m is an open neighborhood of $\operatorname{im}\Xi_m$. For each $m \geq 0$, let $H_m\colon X \to X \setminus \bigcup_{l>m} V_l$ be a homeomorphism defined by the formula

$$H_m(x) = \begin{cases} h_l(x) & x \in U_l \text{ and } l > m \\ x & x \in X \setminus \bigcup_{l>m} U_l \end{cases} .$$

Let $\mathcal{F}_0 = \{F_i^0 = H_0(F_i^{-1})\}_{i\in I}$. Observe that L_0 is an exact n-core of $\mathcal{F}_0$. We let $K_0 = L_0$. By the construction, H_m is an approximation within $\mathcal{F}_{-1}$ of the identity on X, hence $\mathcal{F}_0$ refines $\mathcal{F}_{-1}$. Hence $\mathcal{F}_0$ refines $\operatorname{st}\mathcal{F}$.

We shall recursively construct a sequence of closed interior $\mathcal{N}_n$-covers $\mathcal{F}_m = \{F_i^m\}_{i\in I}$ of $X \setminus \bigcup_{l>m} V_l$ such that

(i) $\mathcal{F}_m$ is k-regular and is isomorphic to $\mathcal{F}_{m-1}$,

(ii) ξ_m is a trivial hole in $F^m_{J_m}$,

(iii) $\mathcal{F}_m$ is $\mathbb{U}_k$-contractible to L_m in $\operatorname{st}_{\{U_l\}_{l\leq m}} \mathcal{F}_{-1}$,

(iv) $\mathcal{F}_m$ is equal to $\mathcal{F}_{m-1}$ on $X \setminus U_m$ and is its swelling on $U_m \setminus V_m$,

(v) $\mathcal{F}_m$ has an exact n-core K_m such that L_m is a subcomplex of K_m and $K_m \setminus K_{m-1} \subset V_m$.

Let $m > 0$ and assume that we already constructed $\mathcal{F}_{m-1}$ and K_{m-1}. Let $\mathcal{F}_{m-1/2} = \{F_i^{m-1/2}\}_{i\in I}$ with

$$F_i^{m-1/2} = \begin{cases} h_m^{-1}(x) & x \in U_m \\ x & x \in X \setminus U_m \end{cases} .$$

The collection $\mathcal{F}_{m-1/2}$ is a closed interior $\mathcal{N}_n$-cover of $X \setminus \bigcup_{l>m} V_l$. Let $K_{m-1/2} = K_{m-1} \cup L_m$. By the construction, $K_{m-1/2}$ is an exact n-core of $\mathcal{F}_{m-1/2}$. Obviously, $\mathcal{F}_{m-1/2}$ is k-regular, is equal to $\mathcal{F}_{m-1}$ on $X \setminus U_m$ and is its swelling on $U_m \setminus V_m$. By (iii), $\mathcal{F}_{m-1}$ is $\mathbb{U}_k$-contractible to L_{m-1} in $\operatorname{st}_{\{U_l\}_{l<m}} \mathcal{F}_{-1}$. Since K_{m-1} is an exact n-core of $\mathcal{F}_{m-1}$, this is equivalent to $\mathbb{U}_k$-contractibility of $\mathcal{F}_{m-1}/K_{m-1}$ to L_{m-1}. Since K_{m-1} is an exact n-core of $\mathcal{F}_{m-1/2}$ as well, $\mathcal{F}_{m-1/2}$ is $\mathbb{U}_k$-contractible to L_{m-1} (and obviously to L_m as well) in $\operatorname{st}_{\{U_l\}_{l<m}} \mathcal{F}_{-1}$. Hence conditions of lemma 11.17 are satsified with $\mathcal{F} = \mathcal{F}_{m-1/2}$, $X = X \setminus \bigcup_{l>m} V_l$, $J = J_m$, $\varphi = \xi_m$, $\varPhi = \Xi_m$, $L = K_{m-1/2}$, $L_0 = L_m$, $\mathcal{H} = \operatorname{st}_{\{U_l\}_{l<m}} \mathcal{F}_{-1}$ and $U = U_m$. Hence there exists a closed interior $\mathcal{N}_n$-cover $\mathcal{F}_m$ of $X \setminus \bigcup_{l>m} V_l$ and an exact n-core K_m of $\mathcal{F}_m$ that satisfy conditions (i)-(v). The construction of the sequence $\mathcal{F}_m$ is done.

Take $\mathcal{F}_\infty = \lim_{m\to\infty} \mathcal{F}_m$. By the construction $\mathcal{U}$ is a discrete collection in X, hence $\mathcal{F}_\infty$ is a closed interior $\mathcal{N}_n$-cover of X. We shall verify that

(0) $\mathcal{F}_\infty$ is k-regular and is isomorphic to $\mathcal{F}$,

(1) for each J in $\mathcal{J}$, the inclusion of $L \cap F_J^\infty$ into F_J^∞ induces a trivial (zero) homomorphism on kth homotopy groups,

(2) $\mathcal{F}_\infty$ is $\mathbb{U}_k$-contractible to L in $\operatorname{st}_{\mathcal{U}} \mathcal{F}_{-1}$.

If (0), (1) and (2) are satisfied, then by lemma 11.12, $\mathcal{F}_\infty$ is $\circledcirc_k$-contractible in $\operatorname{st} \operatorname{st}_{\mathcal{U}} \mathcal{F}_0$. Observe that $\mathcal{U}$ refines $\operatorname{st}^2_{\mathcal{F}} \mathcal{E}$, $\mathcal{F}_{-1}$ refines $\operatorname{st} \mathcal{F}$ and $\operatorname{st} \mathcal{F}$ refines $\operatorname{st}_{\mathcal{F}} \mathcal{E}$, hence $\operatorname{st} \operatorname{st}_{\mathcal{U}} \mathcal{F}_{-1}$ refines $\operatorname{st} \operatorname{st}_{\operatorname{st}^2_{\mathcal{F}} \mathcal{E}} \operatorname{st}^2_{\mathcal{F}} \mathcal{E} = \operatorname{st} \operatorname{st} \operatorname{st}^2_{\mathcal{F}} \mathcal{E}$, which refines $\operatorname{st} \operatorname{st} \operatorname{st}^2 \mathcal{E} = \operatorname{st} \operatorname{st}^7 \mathcal{E} = \operatorname{st}^{22} \mathcal{E}$, the last equalities by lemma 2.2. Then by lemma 11.10 and by remark 11.11, there exists a $(k+1)$-regular closed interior $\mathcal{N}_n$-cover that is isomorphic to $\mathcal{F}_\infty$, that refines $\operatorname{st} \operatorname{st}^{21 \cdot 22 + 10} \mathcal{E} = \operatorname{st}^{472} \mathcal{E}$ and that is equal to $\mathcal{F}_\infty$ on a neighborhood of Z. By the construction, $\mathcal{F}_\infty$ is isomorphic to $\mathcal{F}$ and equal to $\mathcal{F}$ on a neighborhood of Z. Hence verification of (0), (1) and (2) will finish the proof.

By lemma 10.8, $K_\infty = \bigcup_{m \geq 0} K_m$ is an exact n-core of $\mathcal{F}_\infty$. By the construction, $K_\infty \cap F_J^\infty = \bigcup K_m \cap F_J^m$. Since K_m is an exact n-core of $\mathcal{F}_m$ and $\mathcal{F}_m$ is k-regular, each $K_m \cap F_J^m$ is l-connected for each $l < k$. Therefore $\bigcup_{m \geq 0} K_m \cap F_J^m$ is l-connected for each $l < k$. Since K_∞ is an exact n-core of $\mathcal{F}_\infty$, $\mathcal{F}_\infty$ is k-regular. Condition (0) is satisfied.

Let J in $\mathcal{J}$. By the construction, the inclusion $L_0 \cap F_J^\infty \subset L \cap F_J^\infty$ is an n-homotopy equivalence. Hence to prove (1), it suffices to check that the inclusion of $L_0 \cap F_J^\infty$ into F_J^∞ induces a trivial homomorphism on kth homotopy groups. Let φ be a map from S^k into $L_0 \cap F_J^\infty$. By the construction, φ is homotopic in $L_0 \cap F_J^\infty$ with ξ_m for some m such that $J = J_m$. By (ii), ξ_m is nullhomotopic in $K_m \cap F_J^m$. Hence it is nullhomotopic in F_J^∞, as $K_m \cap F_J^m \subset F_J^\infty$.

Every element of $\pi_k(F_J^\infty)$ is homotopic in F_J^∞ with a map into $F_J^\infty \cap K_m$ for some m. Hence $\pi_k(F_J^\infty)$ is generated by maps into $F_J^m \cap K_m$. Such maps, in turn, are $\mathbb{U}_k$-contractible to L in $\operatorname{st}_{\mathcal{U}} \mathcal{F}_{-1}$, by (iii). Hence (2) is satisfied. We are done!

Bibliography

[1] S. Ageev. Axiomatic method of partitions in the theory of Menger and Nöbeling spaces. Available online from Topology Atlas, 2000.

[2] S. Ageev, Jul 2005. Talk at the Geometric Topology Conference in Będlewo (Poland).

[3] R. D. Anderson. On topological infinite deficiency. *Michigan Math. J.*, 14:365–383, 1967.

[4] R. D. Anderson and John D. McCharen. On extending homeomorphisms to Fréchet manifolds. *Proc. Amer. Math. Soc.*, 25:283–289, 1970.

[5] W. Barit. Small extensions of small homeomorphisms. *Notices Amer. Math. Soc.*, 16:295, 1969.

[6] C. Bessaga and A. Pełczyński. The estimated extension theorem, homogeneous collections and skeletons, and their applications to the topological classification of linear matrix spaces and convex sets. *Fund. Math.*, 69:153–190, 1970.

[7] Czesław Bessaga and Aleksander Pełczyński. *Selected topics in infinite-dimensional topology.* PWN—Polish Scientific Publishers, Warsaw, 1975.

[8] Mladen Bestvina. Characterizing k-dimensional universal Menger compacta. *Mem. Amer. Math. Soc.*, 71(380):vi+110, 1988.

[9] Anders Björner. Nerves, fibers and homotopy groups. *J. Combin. Theory Ser. A*, 102(1):88–93, 2003.

[10] Karol Borsuk. On the imbedding of systems of compacta in simplicial complexes. *Fund. Math.*, 35:217–234, 1948.

[11] Philip L. Bowers. Limitation topologies on function spaces. *Trans. Amer. Math. Soc.*, 314(1):421–431, 1989.

[12] T. A. Chapman. Topological invariance of Whitehead torsion. *Amer. J. Math.*, 96:488–497, 1974.

[13] A. Chigogidze. *Inverse spectra*, volume 53 of *North-Holland Mathematical Library.* North-Holland Publishing Co., Amsterdam, 1996.

[14] Alex Chigogidze and Alexandr Karasev. Topological model categories generated by finite complexes. *Monatsh. Math.*, 139(2):129–150, 2003.

[15] T. Dobrowolski and J. Mogilski. Problems on topological classification of incomplete metric spaces. In *Open problems in topology*, pages 409–429. North-Holland, Amsterdam, 1990.

[16] A. N. Dranishnikov, 1997. Math. Reviews MR1406565 (97g:54001).

[17] J. Dugundji. Absolute neighborhood retracts and local connectedness in arbitrary metric spaces. *Compositio Math.*, 13:229–246 (1958), 1958.

[18] Ryszard Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, 1989.

[19] Ja. N. Haimov. The homotopy type of a space having a brick decomposition. *Dokl. Akad. Nauk Tadzhik. SSR*, 22(1):25–29, 1979.

[20] Allen Hatcher. Cw approximation. Available online at http://www.math.cornell.edu/ hatcher/AT/AT-CWapprox.pdf.

[21] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. Available online at http://www.math.cornell.edu/ hatcher/AT/ATpage.html.

[22] David W. Henderson. Infinite-dimensional manifolds are open subsets of Hilbert space. *Topology*, 9:25–33, 1970.

[23] David W. Henderson and R. Schori. Topological classification of infinite dimensional manifolds by homotopy type. *Bull. Amer. Math. Soc.*, 76:121–124, 1970.

[24] W. Holsztyński. On spaces with regular decomposition. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 12:607–611, 1964.

[25] Sze-tsen Hu. *Theory of retracts*. Wayne State University Press, Detroit, 1965.

[26] K. Kawamura, M. Levin, and E. D. Tymchatyn. A characterization of 1-dimensional Nöbeling spaces. *Topology Proc.*, 22(Summer):155–174, 1997.

[27] Kazuhiro Kawamura. Some topics in geometric topology. In *Recent progress in general topology, II*, pages 287–311. North-Holland, Amsterdam, 2002.

[28] M. Levin. A Z-set unknotting theorem for Nöbeling spaces, 2005. Available online from http://www.arxiv.org/pdf/math.GT/0510571.

[29] M. Levin. Characterizing Nöbeling spaces, 2006. Available online from http://www.arxiv.org/pdf/math.GT/0602361.

[30] Albert T. Lundell and Stephen Weingram. *The Topology of CW Complexes*. The University Series in Higher Mathematics. Van Nostrand Reinhold Company, 1969.

[31] Michael C. McCord. Homotopy type comparison of a space with complexes associated with its open covers. *Proc. Amer. Math. Soc.*, 18:705–708, 1967.

[32] A. Nagórko. Carrier and nerve theorems in the extension theory. *Proc. Amer. Math. Soc.*, 135(2):551–558, 2007. http://arxiv.org/abs/math.GN/0506525.

[33] R. B. Sher. The union of two Hilbert cubes meeting in a Hilbert cube need not be a Hilbert cube. *Proc. Amer. Math. Soc.*, 63(1):150–152, 1977.

[34] Edwin H. Spanier. *Algebraic topology.* Springer-Verlag, New York, 1989. Corrected reprint of the 1966 original.

[35] H. Toruńczyk. On CE-images of the Hilbert cube and characterization of Q-manifolds. *Fund. Math.*, 106(1):31–40, 1980.

[36] H. Toruńczyk. Characterizing Hilbert space topology. *Fund. Math.*, 111(3):247–262, 1981.

[37] André Weil. Sur les théorèmes de de Rham. *Comment. Math. Helv.*, 26:119–145, 1952.

[38] James E. West. Infinite products which are Hilbert cubes. *Trans. Amer. Math. Soc.*, 150:1–25, 1970.

[39] James E. West. Open problems in infinite-dimensional topology. In *Open problems in topology*, pages 523–597. North-Holland, Amsterdam, 1990.

[40] J. H. C. Whitehead. Combinatorial homotopy. I. *Bull. Amer. Math. Soc.*, 55:213–245, 1949.

Index

absolute extensor for a space X ... 37
absolute extensor in dimension n ... 8
absolute neighborhood extensor in dimension n ... 8
abstract Nöbeling manifold ... 4
adjustment ... 32
adjustment rel A ... 32
adjustment with a fixed J ... 32
$AE(\mathcal{C})$ ... 37
$AE(n)$ ... 8
$AE(X)$ ... 37
A_l^m ... 82
$ANE(n)$ ... 8
anticanonical map ... 41
approximable within a cover ... 28
approximation within a cover ... 28

$B(f,\mathcal{U})$ ... 11
bst L ... 10
bst v ... 10
$\mathrm{bst}^m L$ ... 10

$C(X,Y)$ ... 11
$\mathcal{C}$-collection ... 27
canonical map ... 41
carrier ... 38
  invertible ... 39
$\mathrm{Cl}_X A$ ... 6
closed partition ... 51
corresponding elements ... 7
cover ... 6
  kth join of a ... 45
  m-barycentric ... 45
  n-contractible ... 44
  n-regular ... 42
  n-semiregular ... 43
  $\circledcirc_k$-contractible ... 83
  $\mathcal{C}$- ... 27
  equal on a set ... 7
  interior ... 6
  isomorphic ... 7
  locally finite dimensional ... 37
  obtained by patching of B ... 80
  regular for a class $\mathcal{C}$ ... 37
  regular for a space X ... 38
  $\mathbb{U}_k$-contractible ... 88
covers
  compatible with a homeomorphism ... 51

∂X ... 6
dom f ... 6

F/A ... 7
$\mathcal{F}_1 \prec_f \mathcal{F}_2$ ... 49
fits ... 60
F_J ... 6
full subcomplex ... 45

hole
  in a cover ... 78
  small ... 78
  trivial ... 78

im f ... 6
indexed collection ... 6
indexing function ... 6
indexing set ... 6
$\mathrm{Int}_X A$ ... 6
intersecting sets ... 6

- limitation topology 11

- map 6
  - approximable 12
  - carried by a carrier 38
  - $\circledcirc_{\mathcal{F}}$-contractible 83
  - $\mathbb{U}_{\mathcal{F}}$-contractible 88
- maps
  - n-homotopic 15
- mth star 7

- n-core 64
  - exact 64
  - outer 65
- n-homotopy equivalence 17
- n-homotopy equivalent 17
- Nöbeling manifold 3
- Nöbeling space 3
- $\mathcal{N}_n$ 14
- $\mathcal{N}_n$-
  - neighborhood 23
  - space 14

- $\circledcirc_{\mathcal{F}}$-contraction 83
- obtained by patching 78

- patch 90
- Polish space 6
- primary complex 45
- proof
  - Z-set unknotting theorem 55
  - characterization theorem 55
  - local Z-set unknotting theorem 54
  - main theorem 54
  - open embedding theorem 55
  - sum theorem 55

- refinement 7
- restriction of a collection 7

- shrinking 7
- $\mathrm{st}_{\mathcal{F}}\,A$ 7
- $\mathrm{st}_{\mathcal{F}}\,x$ 7
- star of a collection 7
- star of a point 7
- star of a set 7
- strong universality in dimension n 12
- swelling 7

- theorem
  - characterization 3
  - local Z-set unknotting 4
  - main 4
  - open embedding 4
  - sum 4
  - topological rigidity 3
  - Z-set unknotting 4

- $\mathcal{U}$-
  - adjustment 32
  - adjustment rel A 32
  - approximation 8
  - close 8
  - deformation 64
  - homotopic 8
- $\mathbb{U}_{\mathcal{F}}$-contraction 88

- weak n-homotopy equivalence 15
- Whitehead's characterization 17

- Z-
  - collection 30
  - embedding 21
  - set 20
- $Z(\mathcal{F})$-
  - embedding 27
  - embedding into X 27
  - set 27
  - set in X 27
- Z_σ-set 21